

Notes on Coxeter Transformations and the McKay correspondence

Rafael Stekolshchik

Key words and phrases. Coxeter transformations, Cartan matrix, McKay correspondence, Poincaré series

email: rs2@biu.013.co.il.

ABSTRACT. We consider the Coxeter transformation in the context of the McKay correspondence, representations of quivers, and Poincaré series.

We study in detail the Jordan forms of the Coxeter transformations and prove shearing formulas due to Subbotin and Sumin for the characteristic polynomials of the Coxeter transformations. Using shearing formulas we calculate characteristic polynomials of the Coxeter transformation for the diagrams $T_{2,3,r}$, $T_{3,3,r}$, $T_{2,4,r}$, prove J. S. Frame's formulas, and generalize R. Steinberg's theorem on the spectrum of the affine Coxeter transformation for the multiply-laced diagrams. This theorem is the key statement in R. Steinberg's proof of the McKay correspondence. For every extended Dynkin diagram, the spectrum of the Coxeter transformation is easily obtained from R. Steinberg's theorem.

In the study of representations π_n of $SU(2)$, we extend B. Kostant's construction of a vector-valued generating function

$$P_G(t) = ([P_G(t)]_0, [P_G(t)]_1, \dots, [P_G(t)]_r).$$

B. Kostant's construction appears in the context of the McKay correspondence and gives a way to obtain multiplicities of indecomposable representations ρ_i of the binary polyhedral group G in the decomposition of $\pi_n|G$. In the case of multiply-laced graphs, instead of indecomposable representations ρ_i we use restricted representations and induced representations of G introduced by P. Slodowy. Using B. Kostant's construction we generalize to the case of multiply-laced graphs W. Ebeling's theorem which connects the Poincaré series $[P_G(t)]_0$ and the Coxeter transformations. According to W. Ebeling's theorem

$$[P_G(t)]_0 = \frac{\chi(t^2)}{\tilde{\chi}(t^2)},$$

where χ is the characteristic polynomial of the Coxeter transformation and $\tilde{\chi}$ is the characteristic polynomial of the corresponding affine Coxeter transformation.

Using the Jordan form of the Coxeter transformation we prove a criterion of V. Dlab and C. M. Ringel of regularity of quiver representations, consider necessary and sufficient conditions of this criterion for extended Dynkin diagrams and for diagrams with indefinite Tits form.

In memory of V. F. Subbotin

Contents

List of Figures	9
List of Tables	11
Chapter 1. Introduction	13
1.1. The three historical aspects of the Coxeter transformation	13
1.2. A brief review of this work	14
1.3. The Spectrum and the Jordan form	15
1.3.1. The Jordan form and reduction to the golden pair of matrices	15
1.3.2. An explicit construction of eigenvectors. The eigenvalues are roots of unity	16
1.3.3. Study of the Coxeter transformation and the Cartan matrix	17
1.3.4. Monotonicity of the dominant eigenvalue of the golden pair	17
1.4. Shearing formulas and the diagrams $T_{p,q,r}$	18
1.4.1. Shearing formulas for the characteristic polynomial	18
1.4.2. An explicit calculation of characteristic polynomials	18
1.4.3. Formulas for the diagrams $T_{2,3,r}, T_{3,3,r}, T_{2,4,r}$	20
1.5. Coxeter transformations and the McKay correspondence	21
1.5.1. The generalized R. Steinberg theorem	21
1.5.2. The Kostant generating functions and W. Ebeling's theorem	22
1.6. The regular representations of quivers	23
1.6.1. Finite-type, tame and wild quivers	23
1.6.2. The Dlab-Ringel defect δ and Δ -defect ρ	23
1.6.3. Necessary and sufficient conditions of regularity	24
Chapter 2. Preliminaries	27
2.1. The Cartan matrix and the Tits form	27
2.2. The Schwartz inequality	28
2.3. The Dynkin diagrams and the extended Dynkin diagrams	29
2.4. The real and imaginary roots	30
2.5. The hyperbolic Dynkin diagrams and hyperbolic Cartan matrices	32
2.6. Orientations and the associated Coxeter transformations	33
2.7. The Poincaré series	34
2.7.1. Graded algebras, symmetric algebras, algebras of invariants	34
2.7.2. Invariants of finite groups generated by reflections	35
Chapter 3. The Jordan normal form of the Coxeter transformation	39
3.1. The Cartan matrix and the Coxeter transformation	39
3.1.1. A bicolored partition and a bipartite graph	39
3.1.2. Conjugacy of Coxeter transformations	40
3.1.3. The Cartan matrix and the bicolored Coxeter transformation	40
3.1.4. Dual graphs and Dual forms	41
3.1.5. Eigenvalues of the Cartan matrix and the Coxeter transformation	41

3.2.	An application of the Perron-Frobenius theorem	42
3.2.1.	The pair of matrices DD^t and D^tD (resp. DF and FD)	42
3.2.2.	The Perron-Frobenius theorem applied to DD^t and D^tD (resp. DF and FD)	45
3.3.	The basis of eigenvectors and a theorem on the Jordan form	45
3.3.1.	An explicit construction of the eigenvectors	45
3.3.2.	Monotonicity of the dominant eigenvalue	47
3.3.3.	A theorem on the Jordan form	49
Chapter 4.	Eigenvalues, shearing formulas and diagrams $T_{p,q,r}$	51
4.1.	The eigenvalues of the affine Coxeter transformation are roots of unity	51
4.2.	Bibliographical notes on the spectrum of the Coxeter transformation	54
4.3.	Shearing and gluing formulas for the characteristic polynomial	56
4.4.	Formulas of the characteristic polynomials for the diagrams $T_{p,q,r}$	61
4.4.1.	The diagrams $T_{2,3,r}$	61
4.4.2.	The diagrams $T_{3,3,r}$	63
4.4.3.	The diagrams $T_{2,4,r}$	66
4.4.4.	Convergence of the sequence of eigenvalues	68
Chapter 5.	R. Steinberg's theorem, B. Kostant's construction	73
5.1.	R. Steinberg's theorem and a (p, q, r) mystery	73
5.2.	Characteristic polynomials for the Dynkin diagrams	76
5.3.	A generalization of R. Steinberg's theorem	78
5.3.1.	Folded Dynkin diagrams and branch points	78
5.3.2.	R. Steinberg's theorem for the multiply-laced case	79
5.4.	The Kostant generating function and Poincaré series	81
5.4.1.	The generating function	81
5.4.2.	The characters and the McKay operator	84
5.4.3.	The Poincaré series and W. Ebeling's theorem	87
Chapter 6.	Regular representations of quivers	91
6.1.	The regular representations and the Dlab-Ringel criterion	91
6.2.	Necessary conditions of regularity of the extended Dynkin diagrams	91
6.3.	The Dlab-Ringel definition of the defect	93
6.4.	Necessary conditions of regularity for diagrams with indefinite Tits form	94
6.5.	Transforming elements and sufficient conditions of regularity	96
6.5.1.	Sufficient conditions of regularity for the bicolored orientation	96
6.5.2.	A theorem on transforming elements	97
6.5.3.	Sufficient conditions of regularity for an arbitrary orientation	99
6.5.4.	Invariance of the defect	100
6.6.	Examples of necessary conditions of regularity	101
6.6.1.	The three equivalence classes of orientations of $\tilde{D}_4$	101
6.6.2.	The bicolored and central orientations of $\tilde{E}_6$	102
6.6.3.	The multiply-laced case. The two orientations of $\tilde{G}_{21}$ and $\tilde{G}_{22} = \tilde{G}_{21}^\vee$	103
6.6.4.	The case of indefinite $\mathcal{B}$. The oriented star $*_{n+1}$	104
Appendix A.	The McKay correspondence and the Slodowy generalization	107
A.1.	Finite subgroups of $SU(2)$ and $SO(3, \mathbb{R})$	107
A.2.	Generators and relations in polyhedral groups	108
A.3.	Kleinian singularities and the Du Val resolution	109
A.4.	The McKay correspondence	111

A.5. The Slodowy generalization of the McKay correspondence	112
A.5.1. The Slodowy correspondence	112
A.5.2. The binary tetrahedral group and the binary octahedral group	113
A.5.3. Representations of the binary octahedral and tetrahedral groups	122
A.5.4. Induced and restricted representations	128
Appendix B. Miscellaneous	133
B.1. The triangle groups and Hurwitz groups	133
B.2. Algebraic integers	134
B.3. The Perron-Frobenius Theorem	135
B.4. The complex projective line and stereographic projection	136
B.5. Prime spectrum, coordinate ring, orbit space	137
B.5.1. Hilbert's Nullstellensatz (Theorem of zeros)	137
B.5.2. Prime spectrum	139
B.5.3. Coordinate ring	140
B.5.4. Orbit space	141
B.6. The orbit structure of the Coxeter transformation	141
B.7. Fixed and anti-fixed points of the Coxeter transformation	145
B.7.1. Chebyshev polynomials and the McKay-Slodowy matrix	145
B.7.2. A theorem on fixed and anti-fixed points	146
Bibliography	149

List of Figures

2.1	The Dynkin diagrams	29
2.2	The extended Dynkin diagrams	30
2.3	The diagram $T_{p,q,r}$	32
3.1	The star $*_{k+1}$ with k rays	44
3.2	The Jordan normal forms of DD^t and D^tD	46
3.3	The Jordan normal form of the Coxeter transformation	49
3.4	A graph Γ such that $\dim \ker \mathbf{B}$ is however large	50
4.1	A split graph Γ	56
4.2	The asterisks mark the deleted line a_m and column b_1	58
4.3	Splitting along the edge l	60
4.4	Splitting along the edge l , $\Gamma_1 \neq \Gamma_2$	61
4.5	The spectral radius attains its minimum on the graph E_{10}	62
5.1	Splitting the graph $T_{p,q,r}$ along the edge $\{\alpha\beta\}$	73
5.2	Diagrams $\tilde{D}_{r+2}$ and D_{r+2}	75
5.3	Folding operation applied to Dynkin diagrams	79
6.1	For $\tilde{D}_4$, the bicolored and central orientations coincide	101
6.2	For $\tilde{E}_6$, the bicolored and central orientations	102
6.3	The two orientations of $\tilde{G}_{21}$ and $\tilde{G}_{22} = \tilde{G}_{21}^\vee$	103
6.4	A bicolored orientation of the star $*_{n+1}$	104
A.1	The minimal resolution $\pi^{-1} : \tilde{X} \longrightarrow X$ for $X = \mathcal{D}_2$	111
A.2	The induced and restricted representations of $\mathcal{T} \triangleleft \mathcal{O}$	129
B.1	The stereographic projection	137

List of Tables

1.1	Characteristic polynomials, Dynkin diagrams	19
1.2	Characteristic polynomials, extended Dynkin diagrams	20
2.1	The exponents n_{ij}	31
2.2	The Poincaré polynomials for the simple compact Lie groups	36
4.1	The eigenvalues of affine Coxeter transformations	52
4.2	The Coxeter numbers and affine Coxeter numbers	53
4.3	Proposition 4.5 for $r = 2, 3$.	55
4.4	The characteristic polynomials for $T_{2,3,r}$	64
4.5	The characteristic polynomials for $T_{3,3,r}$	66
4.6	The characteristic polynomials for $T_{2,4,r}$	69
4.7	The diagrams $T_{p,q,r}$ and characteristic polynomials $\lambda^n f(\lambda) + g(\lambda)$	70
5.1	The binary polyhedral groups (BPG) and the Kostant numbers a, b	84
5.2	The characteristic polynomials $\chi, \tilde{\chi}$ and the Poincaré series	88
A.1	The polyhedral groups in $\mathbb{R}^3$	108
A.2	The finite subgroups of $SU(2)$	109
A.3	The relations $R(x, y, z)$ describing the algebra of invariants $\mathbb{C}[z_1, z_2]^G$	110
A.4	The pairs $H \triangleleft G$ of binary polyhedral groups	113
A.5	The pairs $H \triangleleft G$ and folded extended Dynkin diagrams	114
A.6	The elements of the binary octahedral group	115
A.7	The conjugacy classes in the binary octahedral group	116
A.8	The conjugacy classes in the binary tetrahedral group	121
A.9	The characters of the binary octahedral group	122
A.10	The 3-dimensional characters of S_4	124
A.11	The character of the 4-dimensional representation ρ_7	125
A.12	The characters of the binary tetrahedral group	126
A.13	The 3-dimensional character of A_4	127
A.14	The restricted characters of the binary octahedral group	128
A.15	The inner products $\langle \psi^\downarrow, \chi \rangle$	130
B.1	Orbit of the Coxeter transformation on the highest root	143
B.2	Assembling vectors $z_n = \tau^{(n-1)}\beta - \tau^{(n)}\beta$	143

CHAPTER 1

Introduction

...A second empirical procedure for finding the exponents was discovered by H. S. M. Coxeter. He recognized that the exponents can be obtained from a particular transformation γ in the Weyl group, which he had been studying, and which we take the liberty of calling a Coxeter-Killing transformation...

B. Kostant, [Kos59, p.974], 1959.

1.1. The three historical aspects of the Coxeter transformation

The three areas, where the Coxeter transformation plays a dramatic role, are:

- the theory of Lie algebras of the compact simple Lie groups;
- the representation theory of algebras and quivers;
- the McKay correspondence.

A *Coxeter transformation* or a *Coxeter element* is defined as the product of all the reflections in the root system of a compact Lie group. Neither the choice of simple roots nor the ordering of reflections in the product affects its conjugacy class, see [Bo, Ch.5,§6], see also Remark 6.10. H. S. M. Coxeter studied these elements and their eigenvalues in [Cox51].

Let h be the order of the Coxeter transformation (called the *Coxeter number*), $|\Delta|$ the number of roots in the corresponding root system Δ , and l the number of eigenvalues of the Coxeter transformation, i.e., the rank of the Cartan subalgebra. Then

$$hl = |\Delta|. \tag{1.1}$$

This fact was empirically observed by H. S. M. Coxeter in [Cox51] and proved by A. J. Coleman in [Col58]. One can also find a proof of this fact in B. Kostant's work [Kos59, p.1021].

H. S. M. Coxeter also observed that the order of the Weyl group is equal to

$$(m_1 + 1)(m_2 + 1) \dots (m_l + 1), \tag{1.2}$$

where the m_i are the exponents of the eigenvalues of the Coxeter transformation, and the factors $m_i + 1$ are the degrees of l basic polynomial invariants of the Weyl group. Proofs of these facts were obtained by C. Chevalley [Ch55] and other authors; for historical notes, see [Bo], [Kos59].

Let $\Delta_+ \subset \Delta$ be the subset of positive roots, let

$$\beta = n_1\alpha_1 + \dots + n_l\alpha_l,$$

where $\alpha_i \in \Delta_+$, be the highest root in the root system Δ . The Coxeter number h from (1.1) and coordinates n_i of β are related as follows:

$$h = n_1 + n_2 + \dots + n_l + 1. \tag{1.3}$$

Observation (1.3) is due to H. S. M. Coxeter [Cox49, p.234], see also [Stb59, Th.1.4.], [Bo, Ch.6, 1, §11, Prop.31].

The Coxeter transformation is important in the study of representations of algebras, quivers, partially ordered sets (posets) and lattices. Their distinguished role in this area is related to the construction of the Coxeter functors given by I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev in [BGP73]. Further revelation of the role of Coxeter functors for representations of algebras is due to V. Dlab and C. M. Ringel [DR76]; for a construction of the functor $D\text{Tr}$, see M. Auslander, M. I. Platzeck and I. Reiten [AuPR79], [AuRS95]. For an application of the Coxeter functors in the representations of posets, see [Drz74]; for their applications in the representations of the modular lattices, see [GP74], [GP76].

Another area where the affine Coxeter transformations appeared is the McKay correspondence — a one-to-one correspondence between finite subgroups of $SU(2)$ and simply-laced extended Dynkin diagrams. Affine Coxeter transformations play the principal role in R. Steinberg's work [Stb85] on the proof of the McKay correspondence. B. Kostant ([Kos84]) obtains multiplicities of the representations related to the concrete nodes of the extended Dynkin diagram from the orbit structure of the affine Coxeter transformation, see §A.4, §B.6.

In this work we only consider two areas of application of the Coxeter transformation: representations of quivers and the McKay correspondence. We do not consider other areas where the Coxeter transformation plays an important role, such as the theory of singularities of differentiable maps (the monodromy operator coincides with a Coxeter transformation, see, e.g., [A'C75], [Gu76], [ArGV86], [EbGu99]), Alexander polynomials, pretzel knots, Lehmer's problem, growth series of Coxeter groups (see, e.g., [Lev66], [Hir02], [GH01], [McM02]).

1.2. A brief review of this work

In Ch.2, we recall some common definitions and notions.

In Ch.3, we establish general results about the Jordan form and the spectrum of the Coxeter transformation.

In Ch.4, we give the eigenvalues of the affine Coxeter transformation. After that we prove some shearing formulas concerning the characteristic polynomials of the Coxeter transformation. The main shearing formula is the Subbotin-Sumin *splitting along the edge* formula which is extended in this chapter to multiply-laced diagrams. One of applications of shearing formulas is a construction of characteristic polynomials of the Coxeter transformation for the hyperbolic Dynkin diagrams $T_{2,3,r}$, $T_{3,3,r}$ and $T_{2,4,r}$. Two *Frame formulas* from [Fr51] (see Remark 4.13 and Proposition 5.13) are easily obtained from shearing formulas.

In Ch.5, we generalize a number of results appearing in a context of the McKay correspondence to multiply-laced diagrams. First, we consider R. Steinberg's theorem playing the key role in his proof of the McKay correspondence. Essentially, R. Steinberg observed that the orders of eigenvalues of the affine Coxeter transformation corresponding to the *extended* Dynkin diagram $\tilde{\Gamma}$ coincide with the lengths of branches of the corresponding Dynkin diagram Γ , [Stb85, p.591, (*)].

Further, in Ch.5, we move on to B. Kostant's construction of a vector-valued *generating function* $P_G(t)$ [Kos84]. Let G be a binary polyhedral group, let ρ_i , where $i = 0, \dots, r$, be irreducible representations of G corresponding by the McKay correspondence to simple roots α_i of the extended Dynkin diagram, let π_n , where $n = 0, 1, \dots$, be indecomposable representations of $SU(2)$ in the symmetric algebra $\text{Sym}^n(\mathbb{C}^2)$, where $\text{Sym}^n(V)$ is the n th

symmetric power of V . Let $m_i(n)$ be *multiplicities* in the decomposition

$$\pi_n|G = \sum_{i=0}^r m_i(n)\rho_i; \quad (1.4)$$

set

$$v_n = \sum_{i=0}^r m_i(n)\alpha_i. \quad (1.5)$$

Then the Kostant generating function is defined as the following vector-valued function:

$$P_G(t) = ([P_G(t)]_0, [P_G(t)]_1, \dots, [P_G(t)]_r)^t := \sum_{n=0}^{\infty} v_n t^n. \quad (1.6)$$

In particular, $[P_G(t)]_0$ is the Poincaré series of the algebra of invariants $\text{Sym}(\mathbb{C}^2)^G$, i.e.,

$$[P_G(t)]_0 = P(\text{Sym}(\mathbb{C}^2)^G, t). \quad (1.7)$$

B. Kostant obtained explicit expression for the series $[P_G(t)]_0$, and therefore found a way to calculate the multiplicities $m_i(n)$. In Ch.5, we extend B. Kostant's construction to the case of multiply-laced graphs. For this purpose, we use P. Slodowy's generalization [Sl80, App.III] of the McKay correspondence to the multiply-laced case. The main idea of P. Slodowy is to consider the pair of binary polyhedral groups $H \triangleleft G$ and their *restricted representations* $\rho \downarrow_H^G$ and *induced representations* $\tau \uparrow_H^G$ instead of the representations ρ_i .

In Appendix A, we study in detail P. Slodowy's generalization for the case of the binary octahedral group $\mathcal{O}$ and the binary tetrahedral group $\mathcal{T}$, where $\mathcal{T} \triangleleft \mathcal{O}$. The generalization of the McKay correspondence to the multiply-laced case is said to be the *Slodowy correspondence*.

Generally, one can speak about the *McKay-Slodowy correspondence*.

Finally, in Ch.5 we generalize to the multiply-laced case W. Ebeling's theorem [Ebl02]: it relates the Poincaré series $[P_G(t)]_0$ and the Coxeter transformations. According to W. Ebeling's theorem,

$$[P_G(t)]_0 = \frac{\chi(t^2)}{\tilde{\chi}(t^2)}, \quad (1.8)$$

where χ is the characteristic polynomial of the Coxeter transformation $\mathbf{C}$ and $\tilde{\chi}$ is the characteristic polynomial of the corresponding affine Coxeter transformation $\mathbf{C}_a$, see Theorem 5.12.

The results of Ch.4 and Ch.5 are published for the first time.

In Ch.6, using results on the Jordan form of the Coxeter transformation we prove a criterion of V. Dlab and C.M. Ringel of regularity of quiver representations, we consider necessary and sufficient conditions of this criterion for extended Dynkin diagrams and for diagrams with indefinite Tits form.

1.3. The Spectrum and the Jordan form

1.3.1. The Jordan form and reduction to the golden pair of matrices. In this work we review the research started 30 years ago in the teamwork with V.F. Subbotin and discuss the Jordan form, and the spectrum of the Coxeter transformation. We show that the study of eigenvalues of the Coxeter transformation reduces to the study of the *golden pair* of matrices satisfying conditions of the Perron-Frobenius theorem [MM64], [Ga90] and having other nice properties.

In §3 we recall (see (3.2), (3.4), (3.6)) that Cartan matrix K can be represented in the form

$$K = \begin{cases} 2\mathbf{B} & \text{for } K \text{ symmetric, see (3.2)} \\ T\mathbf{B} & \text{for } K \text{ symmetrizable, see (3.4),} \end{cases} \quad (1.9)$$

where T is a diagonal matrix, $\mathbf{B}$ is a symmetric matrix. For simply-laced (resp. multiply-laced) graphs, the *golden pair* of matrices is

$$DD^t \text{ and } D^tD \text{ (resp. } DF \text{ and } FD),$$

where the matrices D and F are found from the expressions

$$\mathbf{B} = \begin{pmatrix} I_m & D \\ D^t & I_k \end{pmatrix} \quad (1.10)$$

for the simply-laced case and

$$K = \begin{pmatrix} 2I_m & 2D \\ 2F & 2I_k \end{pmatrix} \quad (1.11)$$

for the multiply-laced case.

The mentioned reduction method works only for trees (they have symmetrizable Cartan matrices) and for graphs with even cycles. In §4.2 we give some bibliographical remarks regarding generalized Cartan matrices.

According to Proposition 3.3, relation (3.17), the corresponding eigenvalues $\lambda_{1,2}^{\varphi_i}$ of Coxeter transformations are

$$\lambda_{1,2}^{\varphi_i} = 2\varphi_i - 1 \pm 2\sqrt{\varphi_i(\varphi_i - 1)},$$

where the φ_i are the eigenvalues of DD^t and D^tD .

One of the central results of this work is Theorem 3.14 (§3, Fig. 3.3) on Jordan form [SuSt75, SuSt78, St85]:

1) *The Jordan form of the Coxeter transformation is diagonal if and only if the Tits form is non-degenerate.*

2) *If $\mathcal{B}$ is nonnegative definite, $\mathcal{B} \geq 0$ (Γ is an extended Dynkin diagram), then the Jordan form of the Coxeter transformation contains only one 2×2 Jordan block. All eigenvalues λ_i lie on the unit circle.*

3) *If $\mathcal{B}$ is indefinite and degenerate, then the number of two-dimensional Jordan blocks coincides with $\dim \ker \mathbf{B}$. All other Jordan blocks are 1×1 . There is simple maximal $\lambda_1^{\varphi_1}$ and simple minimal $\lambda_2^{\varphi_1}$ eigenvalues such that*

$$\lambda_1^{\varphi_1} > 1, \quad \lambda_2^{\varphi_1} < 1.$$

C. M. Ringel [Rin94] generalized the result of Theorem 3.14 for non-symmetrizable Cartan matrices, see §4.2.

1.3.2. An explicit construction of eigenvectors. The eigenvalues are roots of unity. An important point of this work is an explicit construction of eigenvectors and adjoint vectors of the Coxeter transformation — the vectors that form a Jordan basis, see Proposition 3.9, [SuSt75, SuSt78]. This construction is used to obtain the necessary and sufficient condition for regularity of representations [SuSt75, SuSt78]. This condition was also found by V. Dlab and C.M. Ringel, [DR76], see §6.1.

The eigenvalues for all cases of extended Dynkin diagrams are easily calculated using a generalized R. Steinberg theorem (Theorem 5.5) and Table 1.2.

Theorem 4.1 and Table 4.1 summarize this calculation as follows ([SuSt79, St81, St82a, St85]):

the eigenvalues of the Coxeter transformation for any extended Dynkin diagram are roots of unity.

The case of $\tilde{A}_n$ is considered in the §4.2. According to formula (4.5), the characteristic polynomial of the Coxeter transformation for $\tilde{A}_n$ is ([MeSu82], [Men85], [Col89], [Shi00], [BT97]):

$$(\lambda^{n-k+1} - 1)(\lambda^k - 1),$$

where k is the number characterizing the conjugacy class of the Coxeter transformation. For $\tilde{A}_n$, there are $[n/2]$ characteristic polynomials [Col89], (see §4.2), but, for all these cases, the eigenvalues of the Coxeter transformation are roots of unity.

1.3.3. Study of the Coxeter transformation and the Cartan matrix. To study of the Coxeter transformation is almost the same as to study of the Cartan matrix. The Cartan matrix and the matrix of the Coxeter transformation (more precisely, the bicolored representative of the conjugacy class of the Coxeter transformation, §3.1.3) are constructed from the same blocks, see relations (3.2), (3.4).

By Proposition 3.1, the eigenvalues λ of the Coxeter transformation and the eigenvalues γ of the matrix $\mathbf{B}$ of the quadratic Tits form are related as follows:

$$\frac{(\lambda + 1)^2}{4\lambda} = (\gamma - 1)^2.$$

By Corollary 3.10, the Jordan form of the Coxeter transformation is diagonal if and only if the Tits form is nondegenerate.

It seems that the Coxeter transformation contains more information than the Cartan matrix, namely, the Coxeter transformation contains additional information about orientation, see §2.6 and considerations on the graphs containing cycles in §4.2.

The Coxeter transformation and the Cartan matrix are also related to the *fixed points* and *anti-fixed points* of the powers of the Coxeter transformation. This connection is given by means of the Chebyshev polynomials in Theorem B.19 ([SuSt82]).

1.3.4. Monotonicity of the dominant eigenvalue of the golden pair. According to Corollary 3.7, the matrices DD^t and D^tD have a common simple positive eigenvalue φ_1 , the maximal eigenvalue. This eigenvalue is said to be the *dominant* eigenvalue. The dominant eigenvalue φ_1 is a certain characteristic of the graph Γ . In Proposition 3.11, we show that if any edge is added to Γ , then this characteristic only grows. The same is true for the maximal eigenvalue $\lambda_1^{\varphi_1}$, see Proposition 3.3, relation (3.17).

Problem. Is the dominant eigenvalue an invariant of the graph in the class of trees, $\mathbb{T}$, i.e., is there one-to-one correspondence between the dominant eigenvalue and the graph if the Tits form of the graph is indefinite?

If there exist two graphs Γ_1 and Γ_2 with the same dominant eigenvalue φ_1 , and θ is the assignment of the dominant eigenvalue to a graph, then what class of graphs $\mathbb{T}/\theta$ we obtain modulo the relation given by θ ?

This problem is solved in this work for the diagrams $T_{2,3,r}$, $T_{3,3,r}$, $T_{2,4,r}$, see Propositions 4.15, 4.17, 4.19 and Tables 4.4, 4.5, 4.6 in §4.4.

1.4. Shearing formulas and the diagrams $T_{p,q,r}$

1.4.1. Shearing formulas for the characteristic polynomial. There is a number of recurrent formulas used to calculate the characteristic polynomial of the Coxeter transformation of a given graph in terms of characteristic polynomials of the Coxeter transformation of the graph's components. Subbotin and Sumin proved the formula of *splitting along the edge* [SuSum82]:

$$\mathcal{X}(\Gamma, \lambda) = \mathcal{X}(\Gamma_1, \lambda)\mathcal{X}(\Gamma_2, \lambda) - \lambda\mathcal{X}(\Gamma_1 \setminus \alpha, \lambda)\mathcal{X}(\Gamma_2 \setminus \beta, \lambda). \quad (1.12)$$

The proof of the Subbotin-Sumin formula is given in Proposition 4.7.

Another formula ([KMSS83]) is given in Proposition 4.10.

V. Kolmykov kindly informed me that the following statement holds:

PROPOSITION 1.1. *If λ is the eigenvalue of Coxeter transformations for graphs Γ_1 and Γ_2 , then λ is also the eigenvalue of the graph Γ obtained by gluing Γ_1 and Γ_2 .*

For details, see Proposition 4.11.

In Proposition 4.8 we generalize formula (1.12) to the multiply-laced case. The formula of *splitting along the weighted edge* holds:

$$\mathcal{X}(\Gamma, \lambda) = \mathcal{X}(\Gamma_1, \lambda)\mathcal{X}(\Gamma_2, \lambda) - \rho\lambda\mathcal{X}(\Gamma_1 \setminus \alpha, \lambda)\mathcal{X}(\Gamma_2 \setminus \beta, \lambda), \quad (1.13)$$

where ρ is the ratio of the lengths of the roots, corresponding to the endpoints of the split edge. Corollary 4.9 deals with the case where Γ_2 contains only one point. In this case, we have

$$\mathcal{X}(\Gamma, \lambda) = -(\lambda + 1)\mathcal{X}(\Gamma_1, \lambda) - \rho\lambda\mathcal{X}(\Gamma_1 \setminus \alpha, \lambda). \quad (1.14)$$

A formula similar to (1.13) can be proved not only for the Coxeter transformation, but, for example, for the Cartan matrix, see È. B. Vinberg's paper [Vin85, Lemma 5.1]. See also Remark 4.13 concerning the works of J. S. Frame and S. M. Gussein-Zade.

1.4.2. An explicit calculation of characteristic polynomials. We use recurrent formulas (1.12) – (1.14) to calculate the characteristic polynomials of the Coxeter transformation for the Dynkin diagrams and extended Dynkin diagrams.

The characteristic polynomials of the Coxeter transformations for the Dynkin diagrams are presented in Table 1.1. For calculations, see §5.2.

The characteristic polynomials of the Coxeter transformations for the extended Dynkin diagrams are presented in Table 1.2. The polynomials χ_n from Table 1.2 are characteristic polynomials of the Coxeter transformation for the Dynkin diagram A_n :

$$\chi_n = (-1)^n \mathcal{X}(A_n),$$

defined (see §4.3 and Remark 4.12) to be

$$\chi_n = \frac{\lambda^{n+1} - 1}{\lambda - 1} = \lambda^n + \lambda^{n-1} + \dots + \lambda^2 + \lambda + 1.$$

For calculations of characteristic polynomials for the extended Dynkin diagrams, see §5.1 and §5.3.2.

TABLE 1.1. Characteristic polynomials, Dynkin diagrams

Dynkin diagram	Characteristic polynomial	Form with denominator
A_n	$\lambda^n + \lambda^{n-1} + \cdots + \lambda + 1$	$\frac{\lambda^{n+1} - 1}{\lambda - 1}$
B_n, C_n	$\lambda^n + 1$	
D_n	$\lambda^n + \lambda^{n-1} + \lambda + 1$	$(\lambda + 1)(\lambda^{n-1} + 1)$
E_6	$\lambda^6 + \lambda^5 - \lambda^3 + \lambda + 1$	$\frac{(\lambda^6 + 1)(\lambda^3 - 1)}{(\lambda^2 + 1)(\lambda - 1)}$
E_7	$\lambda^7 + \lambda^6 - \lambda^4 - \lambda^3 + \lambda + 1$	$\frac{(\lambda + 1)(\lambda^9 + 1)}{(\lambda^3 + 1)}$
E_8	$\lambda^8 + \lambda^7 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	$\frac{(\lambda^{15} + 1)(\lambda + 1)}{(\lambda^5 + 1)(\lambda^3 + 1)}$
F_4	$\lambda^4 - \lambda^2 + 1$	$\frac{\lambda^6 + 1}{\lambda^2 + 1}$
G_2	$\lambda^2 - \lambda + 1$	$\frac{\lambda^3 + 1}{\lambda + 1}$

TABLE 1.2. Characteristic polynomials, extended Dynkin diagrams

Extended Dynkin diagram	Characteristic polynomial	Form with χ_i	Class g
$\tilde{D}_4$	$(\lambda - 1)^2(\lambda + 1)^3$	$(\lambda - 1)^2\chi_1^3$	0
$\tilde{D}_n$	$(\lambda^{n-2} - 1)(\lambda - 1)(\lambda + 1)^2$	$(\lambda - 1)^2\chi_{n-3}\chi_1^2$	0
$\tilde{E}_6$	$(\lambda^3 - 1)^2(\lambda + 1)$	$(\lambda - 1)^2\chi_2^2\chi_1$	0
$\tilde{E}_7$	$(\lambda^4 - 1)(\lambda^3 - 1)(\lambda + 1)$	$(\lambda - 1)^2\chi_3\chi_2\chi_1$	0
$\tilde{E}_8$	$(\lambda^5 - 1)(\lambda^3 - 1)(\lambda + 1)$	$(\lambda - 1)^2\chi_4\chi_2\chi_1$	0
$\widetilde{CD}_n, \widetilde{DD}_n$	$(\lambda^{n-1} - 1)(\lambda^2 - 1)$	$(\lambda - 1)^2\chi_{n-2}\chi_1$	1
$\tilde{F}_{41}, \tilde{F}_{42}$	$(\lambda^2 - 1)(\lambda^3 - 1)$	$(\lambda - 1)^2\chi_2\chi_1$	1
$\tilde{B}_n, \tilde{C}_n, \widetilde{BC}_n$	$(\lambda^n - 1)(\lambda - 1)$	$(\lambda - 1)^2\chi_{n-1}$	2
$\tilde{G}_{21}, \tilde{G}_{22}$	$(\lambda - 1)^2(\lambda + 1)$	$(\lambda - 1)^2\chi_1$	2
A_{11}, A_{12}	$(\lambda - 1)^2$	$(\lambda - 1)^2$	3

1.4.3. Formulas for the diagrams $T_{2,3,r}, T_{3,3,r}, T_{2,4,r}$. For the three classes of diagrams — $T_{2,3,r}, T_{3,3,r}, T_{2,4,r}$ — explicit formulas of characteristic polynomials of the Coxeter transformations are obtained, see §2.5, Fig. 2.3.

The case of $T_{2,3,r}$, where $r \geq 2$, contains diagrams $D_5, E_6, E_7, E_8, \tilde{E}_8, E_{10}$, and so we call these diagrams the E_n -series, where $n = r + 3$. The diagram $T_{2,3,7}$ is *hyperbolic*, (see §2.5) and, for all $r \geq 3$, we have

$$\chi(T_{2,3,r}) = \lambda^{r+3} + \lambda^{r+2} - \sum_{i=3}^r \lambda^i + \lambda + 1, \quad (1.15)$$

see (4.22) and Table 4.4. The spectral radius of $\chi(T_{2,3,r})$ converges to the *Zhang number*¹

$$\sqrt[3]{\frac{1}{2} + \sqrt{\frac{23}{108}}} + \sqrt[3]{\frac{1}{2} - \sqrt{\frac{23}{108}}} \approx 1.324717..., \quad (1.16)$$

¹Hereafter we give all such numbers with six decimal points.

as $r \rightarrow \infty$, see Proposition 4.15 and Remark 4.16. The number (1.16) is also the smallest *Pisot number*, see §B.2.

The case of $T_{3,3,r}$, where $r \geq 2$, contains diagrams $E_6, \tilde{E}_6$, and so we call these diagrams the $E_{6,n}$ -series, where $n = r - 2$. The diagram $T_{3,3,4}$ is *hyperbolic*, (see §2.5) and, for all $r \geq 3$, we have

$$\chi(T_{3,3,r}) = \lambda^{r+4} + \lambda^{r+3} - 2\lambda^{r+1} - 3 \sum_{i=4}^r \lambda^i - 2\lambda^3 + \lambda + 1, \quad (1.17)$$

see (4.27) and Table 4.5. The spectral radius of $\chi(T_{3,3,r})$ converges to

$$\frac{\sqrt{5} + 1}{2} \approx 1.618034\dots,$$

as $r \rightarrow \infty$, see Proposition 4.17.

The case of $T_{2,4,r}$, where $r \geq 2$, contains diagrams $D_6, E_7, \tilde{E}_7$, and so we call these diagrams the $E_{7,n}$ -series, where $n = r - 3$. The diagram $T_{2,4,5}$ is *hyperbolic*, (see §2.5) and, for all $r \geq 4$, we have

$$\chi(T_{2,4,r}) = \lambda^{r+4} + \lambda^{r+3} - \lambda^{r+1} - 2 \sum_{i=4}^r \lambda^i - \lambda^3 + \lambda + 1, \quad (1.18)$$

see (4.35) and Table 4.6. The spectral radius of $\chi(T_{2,4,r})$ converges to

$$\frac{1}{3} + \sqrt[3]{\frac{58}{108} + \sqrt{\frac{31}{108}}} + \sqrt[3]{\frac{58}{108} - \sqrt{\frac{31}{108}}} \approx 1.465571\dots$$

as $r \rightarrow \infty$, see Proposition 4.19.

1.5. Coxeter transformations and the McKay correspondence

1.5.1. The generalized R. Steinberg theorem. Here we generalize R. Steinberg's theorem concerning the mysterious connection between lengths of branches of any Dynkin diagram and orders of eigenvalues of the affine Coxeter transformation. R. Steinberg proved this theorem for the simply-laced case in [Stb85, p.591,(*)]; it was a key statement in his explanation of the phenomena of the McKay correspondence.

We prove the R. Steinberg theorem for the simply-laced case in §5.1, Theorem 5.1. The multiply-laced case (generalized R. Steinberg's theorem) is proved in §5.3.2, Theorem 5.5. Essentially, the generalized R. Steinberg theorem immediately follows from Table 1.2.

THEOREM 1.2 (The generalized R. Steinberg theorem). *The affine Coxeter transformation with the extended Dynkin diagram $\tilde{\Gamma}$ has the same eigenvalues as the product of some Coxeter transformations of types A_i , where $i \in \{p-1, q-1, r-1\}$ and matches to $(3-g)$ branches of the corresponding Dynkin diagram Γ , where g is the class number (5.33) of $\tilde{\Gamma}$. In other words,*

For $g = 0$, the product $\chi_{p-1}\chi_{q-1}\chi_{r-1}$, is taken.

For $g = 1$, the product $\chi_{p-1}\chi_{q-1}$, is taken.

For $g = 2$, the product consists of only from χ_{p-1} .

For $g = 3$, the product is trivial ($= 1$).

For details, see Theorem 5.5.

1.5.2. The Kostant generating functions and W. Ebeling's theorem. Now we consider B. Kostant's construction of the vector-valued *generating function* $P_G(t)$ [Kos84]. Let G be a binary polyhedral group, and ρ_i , where $i = 0, \dots, r$, irreducible representations of G corresponding due to the McKay correspondence to simple roots α_i of the extended Dynkin diagram; let π_n , where $n = 0, 1, \dots$, be irreducible representations of $SU(2)$ in $\text{Sym}^n(\mathbb{C}^2)$. Let $m_i(n)$ be *multiplicities* in the decomposition

$$\pi_n|G = \sum_{i=0}^r m_i(n)\rho_i, \quad (1.19)$$

and so $m_i(n) = \langle \pi_n|G, \rho_i \rangle$; set

$$v_n = \sum_{i=0}^r m_i(n)\alpha_i = \begin{pmatrix} m_0(n) \\ \dots \\ m_r(n) \end{pmatrix} \quad (1.20)$$

Then

$$P_G(t) = \sum_{n=0}^{\infty} v_n t^n = \begin{pmatrix} \sum_{n=0}^{\infty} m_0(n)t^n \\ \dots \\ \sum_{n=0}^{\infty} m_r(n)t^n \end{pmatrix}. \quad (1.21)$$

Thus, $P_G(t) = ([P_G(t)]_0, [P_G(t)]_1, \dots, [P_G(t)]_r)^t$ is a vector-valued series. In particular, $[P_G(t)]_0$ is the Poincaré series of the algebra of invariants $\text{Sym}(\mathbb{C}^2)^G$, i.e.,

$$[P_G(t)]_0 = P(\text{Sym}(\mathbb{C}^2)^G, t). \quad (1.22)$$

B. Kostant obtained explicit formulas for the series $[P_G(t)]_0$, and therefore a way to calculate the multiplicities $m_i(n)$. B. Kostant's construction is generalized in Ch. 5 to the multiply-laced case. For this purpose, we use the P. Slodowy generalization [Sl80, App.III] of the McKay correspondence to the multiply-laced case.

The main idea of P. Slodowy is to consider a pair of binary polyhedral groups $H \triangleleft G$ and their *restricted representations* $\rho \downarrow_H^G$ and *induced representations* $\tau \uparrow_H^G$ instead of representations ρ_i . In Appendix A, we study in detail P. Slodowy's generalization for the pair $\mathcal{T} \triangleleft \mathcal{O}$, where $\mathcal{O}$ is the binary octahedral group and $\mathcal{T}$ is the binary tetrahedral group. We call the generalization of the McKay correspondence to the multiply-laced case the *Slodowy correspondence*. Finally, in Ch. 5, we generalize to the multiply-laced case W. Ebeling's theorem [Ebl02] which relates the Poincaré series $[P_G(t)]_0$ and the Coxeter transformations.

First, we prove the following proposition due to B. Kostant [Kos84]. It holds for the McKay operator and also for the Slodowy operator.

PROPOSITION 1.3 (B. Kostant [Kos84]). *If B is either the McKay operator A or the Slodowy operator $\tilde{A}$ or $\tilde{A}^\vee$, then*

$$Bv_n = v_{n-1} + v_{n+1}.$$

For details, see Proposition 5.10 in §5.4.2.

THEOREM 1.4 (generalized W. Ebeling theorem, [Ebl02]). *Let G be a binary polyhedral group and $[P_G(t)]_0$ the Poincaré series (5.55) of the algebra of invariants $\text{Sym}(\mathbb{C}^2)^G$. Then*

$$[P_G(t)]_0 = \frac{\det M_0(t)}{\det M(t)},$$

where

$$\det M(t) = \det |t^2 I - \mathbf{C}_a|, \quad \det M_0(t) = \det |t^2 I - \mathbf{C}|,$$

and where $\mathbf{C}$ is the Coxeter transformation and $\mathbf{C}_a$ is the corresponding affine Coxeter transformation.

In Theorem 1.4 the Coxeter transformation $\mathbf{C}$ and the affine Coxeter transformation $\mathbf{C}_a$ are related to the binary polyhedral group G . For the multiply-laced case, we consider a pair of binary polyhedral groups $H \triangleleft G$ and $\mathbf{C}$, $\mathbf{C}_a$ are related again to the group G . We generalize W. Ebeling's theorem for the multiply-laced case, see Theorem 5.12. For definition of the Poincaré series for the multiply-laced case, see (5.56) from §5.4.1 and Remark 5.8.

1.6. The regular representations of quivers

1.6.1. Finite-type, tame and wild quivers. In [BGP73], Bernshtein, Gelfand and Ponomarev introduced *regular representations* of quivers, i.e., representations that never vanish under the Coxeter transformations. More precisely, let V be any representation in the category of representations $\mathcal{L}(\Gamma, \Delta)$ of the graph Γ with an orientation Δ , and let Φ^+ , Φ^- be the Coxeter functors ([BGP73], [DR76]). The representation V is said to be *regular* if

$$(\Phi^+)^k (\Phi^-)^k (\dim V) = (\Phi^+)^k (\Phi^-)^k (\dim V) = \dim V \text{ for all } k \in \mathbb{Z}. \quad (1.23)$$

The Dynkin graphs do not have regular representations; in the category of graph representations, only a finite set of irreducible representations is associated to any Dynkin diagram. Graphs with such property are called *finite type quivers*. According to P. Gabriel's theorem [Gab72], *a quiver is of finite type if and only if it is a (simply laced) Dynkin diagram*.

In the category of all representations of a given quiver, the regular representations are the most complicated ones; they have been completely described only for the extended Dynkin diagrams which for this reason, in the representation theory of quivers, were called *tame quivers* ([Naz73], [DR76]).

Any quiver with indefinite Tits form $\mathcal{B}$ is *wild*, i.e., the description of its representations contains the problem of classifying pairs of matrices up to simultaneous similarity; this classification is hopeless in a certain sense, see [GP69], [Drz80].

By (1.23), the representation V is regular if and only if

$$\mathbf{C}^k (\dim V) > 0 \text{ for all } k \in \mathbb{Z}, \quad (1.24)$$

see [DR76], [St75], [SuSt75], [SuSt78].

1.6.2. The Dlab-Ringel defect δ and Δ -defect ρ . In [St75], [SuSt75], [SuSt78], the linear form $\rho_{\Delta'}(z)$ was considered, see Definition 6.4. For the simply-laced case, it is defined as

$$\rho_{\Delta'}(z) = \langle Tz, \tilde{z}^1 \rangle, \quad (1.25)$$

and for the multiply-laced case as

$$\rho_{\Delta'}(z) = \langle Tz, \tilde{z}^{1\nu} \rangle. \quad (1.26)$$

Here z^1 is the eigenvector of the Coxeter transformation corresponding to eigenvalue 1, and $z^{1\nu}$ is the eigenvector corresponding to eigenvalue 1 of the Coxeter transformation for the dual diagram $\Gamma^\vee$; $\tilde{v}$ denotes the *dual* vector to v obtained from v by changing the sign of the $\mathbb{Y}$ -component (see Definition 6.2).

The linear form $\rho_{\Delta'}(z)$ is said to be the Δ' -defect of the vector z , or the *defect of the vector z in the orientation Δ'* .

In [DR76], V. Dlab and C. M. Ringel introduced the *defect* $\delta_{\Delta'}$ obtained as the solution to the following equation, see (6.20):

$$\mathbf{C}_{\Delta'}^* \delta_{\Delta'} = \delta_{\Delta'}.$$

In Proposition 6.6, [St85] we show that the Dlab-Ringel defect $\delta_{\Delta'}$ coincides with the Δ' -defect $\rho_{\Delta'}$.

1.6.3. Necessary and sufficient conditions of regularity. In Theorem 6.5 proved by Dlab-Ringel [DR76] and by Subbotin-Stekolshchik [SuSt75], [SuSt78], we show the necessary condition of regularity of the representation V :

If $\dim V$ is the regular vector for the extended Dynkin diagram Γ in the orientation Δ' , then

$$\rho_{\Delta'}(\dim V) = 0. \quad (1.27)$$

In Proposition 6.8 (for the bicolored orientation) and in Proposition 6.13 (for an arbitrary orientation Δ') we show that (see [St82])

the condition (1.27) is also sufficient if $\dim V$ is a positive root in the root system associated with the extended Dynkin diagram Γ .

To prove the sufficient condition of regularity, we study the transforming elements interrelating Coxeter transformations for different orientations. Here, Theorem 6.11, [St82], plays a key role. Proposition 6.9 and Theorem 6.11 yield the following:

1) Let Δ', Δ'' be two arbitrary orientations of the graph Γ that differ by the direction of k edges. Consider a chain of orientations, in which every two adjacent orientations differ by the direction of one edge:

$$\Delta' = \Delta_0, \Delta_1, \Delta_2, \dots, \Delta_{k-1}, \Delta_k = \Delta''.$$

Then, in the Weyl group, there exist elements P_i and S_i , where $i = 1, 2, \dots, k$, such that

$$\begin{aligned} \mathbf{C}_{\Delta_0} &= P_1 S_1, \\ \mathbf{C}_{\Delta_1} &= S_1 P_1 = P_2 S_2, \end{aligned}$$

$$\begin{aligned} \mathbf{C}_{\Delta_{k-1}} &= S_{k-1} P_{k-1} = P_k S_k, \\ \mathbf{C}_{\Delta_k} &= S_k P_k. \end{aligned}$$

2) $T^{-1} \mathbf{C}_{\Delta'} T = \mathbf{C}_{\Delta''}$ for the following $k+1$ transforming elements $T := T_i$:

$$\begin{aligned} T_1 &= P_1 P_2 P_3 \dots P_{k-2} P_{k-1} P_k, \\ T_2 &= P_1 P_2 P_3 \dots P_{k-2} P_{k-1} S_k^{-1}, \\ T_3 &= P_1 P_2 P_3 \dots P_{k-2} S_{k-1}^{-1} S_k^{-1}, \end{aligned}$$

$$\begin{aligned} T_{k-1} &= P_1 P_2 S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}, \\ T_k &= P_1 S_2^{-1} S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}, \\ T_{k+1} &= S_1^{-1} S_2^{-1} S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}. \end{aligned}$$

In addition, for each reflection σ_α , there exists a T_i whose decomposition does not contain this reflection.

3) The following relation holds:

$$T_p T_q^{-1} = \mathbf{C}_{\Delta'}^{q-p}.$$

For the graph Γ with indefinite Tits form $\mathcal{B}$, we prove the following *necessary condition of regularity*, see Theorem 6.7 [SuSt75], [SuSt78]:

If z is the regular vector in the orientation Δ' , then

$$\langle Tz, \tilde{z}_1^m \rangle \leq 0, \quad \langle Tz, \tilde{z}_2^m \rangle \geq 0,$$

where $\tilde{z}_1^m$ and $\tilde{z}_2^m$ are the *dual vectors* (Definition 6.2) to the vectors z_1^m and z_2^m corresponding to the maximal eigenvalue $\varphi^m = \varphi^{max}$ of DD^t and D^tD , respectively, see §6.4.

Similar results were obtained by Y. Zhang in [Zh89, Prop.1.5], and by J.A. de la Peña and M. Takane in [PT90, Th.2.3].

For an application of this necessary condition to the star graph, see §6.6.4.

CHAPTER 2

Preliminaries

...Having computed the m 's several years earlier ..., I recognized them¹ in the Poincaré polynomials while listening to Chevalley's address at the International Congress in 1950. I am grateful to A. J. Coleman for drawing my attention to the relevant work of Racah, which helps to explain the "coincidence"; also, to J. S. Frame for many helpful suggestions...

H. S. M. Coxeter, [**Cox51**, p.765], 1951

2.1. The Cartan matrix and the Tits form

Let K be a matrix with the following properties: [**Mo68**], [**Kac80**]

- (C1) $k_{ii} = 2$ for $i = 1, \dots, n$,
- (C2) $-k_{ij} \in \mathbb{Z}_+ = \{0, 1, 2, \dots\}$ for $i \neq j$,
- (C3) $k_{ij} = 0$ implies $k_{ji} = 0$ for $i, j = 1, \dots, n$.

Such a matrix is called a *generalized Cartan matrix*. A matrix M is said to be *symmetrizable* if there exists an invertible diagonal matrix T with positive integer coefficients and a symmetric matrix $\mathbf{B}$ such that $M = T\mathbf{B}$. For example, any generalized Cartan matrix K whose diagram contains no cycles is symmetrizable ([**Mo68**, §3]). Any Cartan matrix whose diagram is a simply-laced diagram (even with cycles) is symmetrizable because it is symmetric. In particular, $\tilde{A}_n$ has a symmetrizable Cartan matrix. In this work we consider only diagrams without cycles, so the diagrams we consider have symmetrizable Cartan matrices.

Let Γ_0 (resp. Γ_1) be the set of vertices (resp. edges) of the graph Γ . A *valued graph* (Γ, d) ([**DR76**], [**AuRS95**]) is a finite set Γ_1 (of edges) rigged with non-negative integers d_{ij} for all pairs $i, j \in \partial\Gamma_1$ of the endpoints of the edges in such a way that $d_{ii} = 0$ and there exist numbers $f_i \in \mathbb{N}$ satisfying

$$d_{ij}f_j = d_{ji}f_i \quad \text{for all } i, j \in \partial\Gamma_1. \quad (2.1)$$

The rigging of the edges of Γ_1 is depicted by symbols

$$i \xrightarrow{(d_{ij}, d_{ji})} j$$

If $d_{ij} = 1 = d_{ji}$, we simply write

$$i \text{ --- } j$$

There is, clearly, a one-to-one correspondence between valued graphs and symmetrizable Cartan matrices, see [**AuRS95**].

¹The numbers $m_1, \dots, m_n$ are said to be *exponents* of the Weyl group. See §2.7.2.

A non-symmetric bilinear form $\mathcal{Q}$ defined on vectors–dimensions of the quiver representation was independently introduced by Ringel [Rin76] and Ovsienko and Roiter [OR77]. The form $\mathcal{Q}$ has the following property important for the representations of quivers [Rin76]:

$$\mathcal{Q}(\dim X, \dim Y) = \dim \operatorname{Hom}(X, Y) - \dim \operatorname{Ext}^1(X, Y). \quad (2.2)$$

The *Tits form* is the quadratic form $\mathcal{B}$ given by the relation

$$\mathcal{B}(\alpha) = \mathcal{Q}(\alpha, \alpha). \quad (2.3)$$

The corresponding symmetric bilinear form is

$$(\alpha, \beta) = \frac{1}{2}(\mathcal{Q}(\alpha, \beta) + \mathcal{Q}(\beta, \alpha)). \quad (2.4)$$

Let $\mathbf{B}$ be the matrix of the quadratic form $\mathcal{B}$. In the simply-laced case,

$$K = 2\mathbf{B} \quad (2.5)$$

with the symmetric Cartan matrix K . So, in the simply-laced case, the Tits form is the Cartan-Tits form. In the multiply-laced case, the symmetrizable matrix K factorizes

$$K = T\mathbf{B},$$

$$k_{ij} = \frac{2(a_i, b_j)}{(a_i, a_i)}, \quad k_{ji} = \frac{2(a_i, b_j)}{(b_j, b_j)}, \quad (2.6)$$

where T is a diagonal matrix with positive integers on the diagonal, the vectors a_i and b_j are defined in §3.1.2, see also §3.1.3.

2.2. The Schwartz inequality

Let $\mathcal{B}$ be the quadratic Tits form associated with an arbitrary tree graph (simply or multiply laced) and $\mathbf{B}$ the matrix of $\mathcal{B}$. Let

$$\ker \mathbf{B} = \{x \mid \mathbf{B}x = 0\}. \quad (2.7)$$

Since $(x, y) = \langle x, \mathbf{B}y \rangle = \langle \mathbf{B}x, y \rangle$, it follows that

$$\ker \mathbf{B} = \{x \mid (x, y) = 0 \text{ for all } y \in \mathcal{E}_F\} \subseteq \{x \mid \mathcal{B}(x) = 0\}. \quad (2.8)$$

We will write $\mathcal{B} > 0$ (resp. $\mathcal{B} \geq 0$) if $\mathcal{B}$ is positively (resp. nonnegative) definite. Let $x, y \in \mathbb{R}^n$, where n is the number of vertices in Γ_0 . Then the following *Schwartz inequality* is true:

$$(x, y)^2 \leq \mathcal{B}(x)\mathcal{B}(y). \quad (2.9)$$

To prove (2.9), it suffices to consider the inequality $(x + \alpha y, x + \alpha y) \geq 0$ which is true for all $\alpha \in \mathbb{R}$. Then the discriminant of the polynomial

$$(x, x) + 2\alpha(x, y) + \alpha^2(y, y)$$

should be non-positive, whence (2.9).

Over $\mathbb{C}$, there exists $x \notin \ker \mathbf{B}$ such that $\mathcal{B}(x) = 0$. For example, the eigenvectors of the Coxeter transformation with eigenvalues $\lambda \neq \pm 1$ satisfy this condition because

$$\mathcal{B}(x) = \mathcal{B}(Cx) = \lambda^2 \mathcal{B}(x).$$

If $\mathcal{B}(x) = 0$ and $x \in \mathbb{R}^n$, then from (2.9) we get $(x, y) \leq 0$ and $(x, -y) \leq 0$, i.e., $(x, y) = 0$ for all $y \in \mathbb{R}^n$. In other words, $x \in \ker \mathbf{B}$. Taking (2.8) into account we see that if $\mathcal{B} \geq 0$, then

$$\ker \mathbf{B} = \{x \mid \mathcal{B}(x) = 0\}. \quad (2.10)$$

2.3. The Dynkin diagrams and the extended Dynkin diagrams

For the proof of the following well-known characterization of the Dynkin diagrams and the extended Dynkin diagrams, see, e.g., [Bo], [DR76]:

1) (Γ, d) is a Dynkin diagram if and only if its quadratic form is positive definite, see Fig. 2.1.

2) (Γ, d) is an extended Dynkin diagram if and only if its quadratic form is nonnegative definite, see Fig. 2.2.

REMARK 2.1. 1) All quadratic forms $\mathcal{B}$ fall into 3 non-intersecting sets

- (a) $\mathcal{B}$ is positive definite, $\mathcal{B} > 0$ (the Dynkin diagrams),
- (b) $\mathcal{B}$ is nonnegative definite, $\mathcal{B} \geq 0$ (the extended Dynkin diagrams),
- (c) $\mathcal{B}$ is indefinite.

2) Consider two operations:

2.1) “Add Edge”: add a vertex and connect it with Γ by only one edge. We denote the new graph $\overset{\wedge}{\Gamma}$.

2.2) “Remove Vertex”: remove a vertex (the new graph may contain more than one component). We denote the new graph $\overset{\vee}{\Gamma}$.

3) The set (b) ($\mathcal{B} \geq 0$) is not stable under these operations:

$$\begin{aligned} \wedge : \mathcal{B} \geq 0 &\implies \mathcal{B} \text{ is indefinite,} \\ \vee : \mathcal{B} \geq 0 &\implies \mathcal{B} > 0. \end{aligned}$$

The set (a) is invariant under $\vee$ and sometimes is invariant under $\wedge$. The set (c) is invariant under $\wedge$ and sometimes is invariant under $\vee$.

4) If the graph Γ with indefinite form $\mathcal{B}$ is obtained from any Dynkin diagram by adding an edge, then the same graph Γ can be obtained by adding an edge (or maybe several edges) to an extended Dynkin diagram.




FIGURE 2.1. The Dynkin diagrams




FIGURE 2.2. The extended Dynkin diagrams

2.4. The real and imaginary roots

Recall now the definitions of *imaginary* and *real* roots in the infinite root system associated with infinite dimensional Kac-Moody Lie algebras. We mostly follow V. Kac's definitions [Kac80], [Kac82], [Kac93].

We consider the vector space $\mathcal{E}_\Gamma$ over $\mathbb{Q}$; set

$$\dim \mathcal{E}_\Gamma = |\Gamma_0|.$$

Let

$$\alpha_i = \{0, 0, \dots, 0, \overset{i}{1}, 0, \dots, 0\} \in \mathcal{E}_\Gamma$$

be the basis vector corresponding to the vertex $v_i \in \Gamma_0$. The space $\mathcal{E}_\Gamma$ is spanned by the vectors $\{\alpha_i \mid i \in \Gamma_0\}$; the vectors $\alpha_1, \dots, \alpha_n$ form the basis in $\mathcal{E}_\Gamma$. Let

$$\mathcal{E}_+ = \{\alpha = \sum k_i \alpha_i \in \mathcal{E}_\Gamma \mid k_i \in \mathbb{Z}_+, \sum k_i > 0\}$$

be the set of all non-zero elements in $\mathcal{E}_\Gamma$ with non-negative integer coordinates in the basis $\{\alpha_1, \dots, \alpha_n\}$.

Define the linear functions $\phi_1, \dots, \phi_n$ on $\mathcal{E}_\Gamma$ by means of the elements of the Cartan matrix (2.6):

$$\phi_i(\alpha_j) = k_{ij}. \quad (2.11)$$

The *positive root system* Δ_+ associated to the Cartan matrix K is a subset in $\mathcal{E}_+$ defined by the following properties (R1)–(R3):

(R1) $\alpha_i \in \Delta_+$ and $2\alpha_i \notin \Delta_+$ for $i = 1, \dots, n$.

(R2) If $\alpha \in \Delta_+$ and $\alpha \neq \alpha_i$, then $\alpha + k\alpha_i \in \Delta_+$ for $k \in \mathbb{Z}$ if and only if $-p \leq k \leq q$, where p and q are some non-negative integers such that $p - q = \phi_i(\alpha)$.

(R3) Any $\alpha \in \Delta_+$ has a connected support.

We define endomorphisms $\sigma_1, \dots, \sigma_n$ of $\mathcal{E}_\Gamma$ by the formula

$$\sigma_i(x) = x - \phi_i(x)\alpha_i. \quad (2.12)$$

Each endomorphism σ_i is the reflection in the hyperplane $\phi_i = 0$ such that $\sigma_i(\alpha_i) = -\alpha_i$. These reflections satisfy the following relations:

$$\sigma_i^2 = 1, \quad (\sigma_i \sigma_j)^{n_{ij}} = 1,$$

where n_{ij} are given in the Table 2.1.

TABLE 2.1. The exponents n_{ij}

$k_{ij}k_{ji}$	0	1	2	3	≥ 4
n_{ij}	2	3	4	6	∞

The group W generated by the reflections $\sigma_1, \dots, \sigma_n$ is called the *Weyl group*. The vectors $\alpha_1, \dots, \alpha_n$ are called *simple roots*; we denote by Π the set of all simple roots. Let $W(\Pi)$ be the orbit of Π under the W -action.

Set

$$M = \{\alpha \in \mathcal{E}_+ \mid \phi_i(\alpha) \leq 0 \text{ for } i = 1, \dots, n, \text{ and } \alpha \text{ has a connected support}\}.$$

The set M is called the *fundamental set*. Let $W(M)$ be the orbit of M under the W -action. We set

$$\Delta_+^{re} = \bigcup_{w \in W} (w(\Pi) \cap \mathcal{E}_+), \quad \Delta_+^{im} = \bigcup_{w \in W} (w(M)).$$

The elements of the set Δ_+^{re} are called *real roots* and the elements of the set Δ_+^{im} are called *imaginary roots*. By [Kac80], the *system of positive roots* Δ_+ is the disjoint union of the sets Δ_+^{re} and Δ_+^{im} :

$$\Delta_+^{re} = \Delta_+^{re} \cup \Delta_+^{im}.$$

If the Tits form $\mathcal{B}$ (or, which is the same, the Cartan matrix K) is positive definite, then the root system is finite, it corresponds to a simple finite dimensional Lie algebra. In this case, the root system consists of real roots.

If the Tits form $\mathcal{B}$ is nonnegative definite, $\mathcal{B} \geq 0$, we have an infinite root system whose imaginary root system is one-dimensional:

$$\Delta_+^{im} = \{\delta, 2\delta, 3\delta, \dots\}, \quad \text{where } \delta = \sum_i k_i \alpha_i,$$

the coefficients k_i being the labels of the vertices from Fig. 2.2.

The numerical labels at the vertices are the coefficients of the imaginary root which coincides with the fixed point z^1 of the Coxeter transformation, see §3.3.1.

The elements $k\delta$ ($k \in \mathbb{N}$) are called *nil-roots*. Every nil-root is a fixed point for the Weyl group.

2.5. The hyperbolic Dynkin diagrams and hyperbolic Cartan matrices

A connected graph Γ with indefinite Tits form is said to be a *hyperbolic Dynkin diagram* (resp. *strictly hyperbolic Dynkin diagram*) if every subgraph $\Gamma' \subset \Gamma$ is a Dynkin diagram or an extended Dynkin diagram (resp. Dynkin diagram). The corresponding Cartan matrix K is said to be the *hyperbolic Cartan matrix* (resp. *strictly hyperbolic Cartan matrix*), see [Kac93, exs. of §4.10]. The corresponding Weyl group is said to be a *hyperbolic Weyl group* (resp. *compact hyperbolic Weyl group*), see [Bo, Ch.5, exs. of §4].




FIGURE 2.3. The diagram $T_{p,q,r}$

Consider the Dynkin diagram $T_{p,q,r}$ depicted on Fig. 2.3. By eq.(3.3) the quadratic Tits form $\mathcal{B}$ can be expressed in the following form

$$\begin{aligned} \frac{1}{2}\mathcal{B}(z) = & x_1^2 + x_2^2 + \cdots + x_{p-1}^2 - x_1x_2 - \cdots - x_{p-2}x_{p-1} - x_{p-1}u_0 + \\ & y_1^2 + y_2^2 + \cdots + y_{q-1}^2 - y_1y_2 - \cdots - y_{q-2}y_{q-1} - y_{q-1}u_0 + \\ & z_1^2 + z_2^2 + \cdots + z_{r-1}^2 - z_1z_2 - \cdots - z_{r-2}z_{r-1} - z_{r-1}u_0 + u_0^2, \end{aligned} \quad (2.13)$$

where u_0 is the coordinate of the vector z corresponding to the branch point of the diagram. Let

$$\begin{aligned} V(x_1, \dots, x_{p-1}, u_0) = & x_1^2 + x_2^2 + \cdots + x_{p-1}^2 - x_1x_2 - \cdots - x_{p-2}x_{p-1} - x_{p-1}u_0, \\ V(y_1, \dots, y_{q-1}, u_0) = & y_1^2 + y_2^2 + \cdots + y_{q-1}^2 - y_1y_2 - \cdots - y_{q-2}y_{q-1} - y_{q-1}u_0, \\ V(z_1, \dots, z_{r-1}, u_0) = & z_1^2 + z_2^2 + \cdots + z_{r-1}^2 - z_1z_2 - \cdots - z_{r-2}z_{r-1} - z_{r-1}u_0. \end{aligned} \quad (2.14)$$

Then

$$\frac{1}{2}\mathcal{B}(z) = V(x_1, \dots, x_{p-1}) + V(y_1, \dots, y_{q-1}) + V(z_1, \dots, z_{r-1}) + u_0^2. \quad (2.15)$$

It is easy to check that

$$\begin{aligned} 2V(x_1, \dots, x_{p-1}, u_0) &= \sum_{i=1}^{p-1} \frac{i+1}{i} \left(x_i - \frac{i}{i+1}x_{i+1}\right)^2 - \frac{p-1}{p}u_0^2, \\ 2V(y_1, \dots, y_{q-1}, u_0) &= \sum_{i=1}^{q-1} \frac{i+1}{i} \left(y_i - \frac{i}{i+1}y_{i+1}\right)^2 - \frac{q-1}{q}u_0^2, \\ 2V(z_1, \dots, z_{r-1}, u_0) &= \sum_{i=1}^{r-1} \frac{i+1}{i} \left(z_i - \frac{i}{i+1}z_{i+1}\right)^2 - \frac{r-1}{r}u_0^2. \end{aligned} \quad (2.16)$$

Denote by $U(x)$, $U(y)$, $U(z)$ the corresponding sums in (2.16):

$$\begin{aligned} 2V(x_1, \dots, x_{p-1}, u_0) &= U(x) - \frac{p-1}{p}u_0^2, \\ 2V(y_1, \dots, y_{p-1}, u_0) &= U(y) - \frac{q-1}{q}u_0^2, \\ 2V(z_1, \dots, z_{p-1}, u_0) &= U(z) - \frac{r-1}{r}u_0^2. \end{aligned} \quad (2.17)$$

Thus,

$$\mathcal{B}(z) = U(x) + U(y) + U(z) - \frac{p-1}{p}u_0^2 - \frac{q-1}{q}u_0^2 - \frac{r-1}{r}u_0^2 + 2u_0^2, \quad (2.18)$$

or

$$\mathcal{B}(z) = U(x) + U(y) + U(z) + \left(\frac{1}{p} + \frac{1}{q} + \frac{1}{r} - 1 \right) u_0^2. \quad (2.19)$$

Set

$$\mu = \frac{1}{p} + \frac{1}{q} + \frac{1}{r}; \quad (2.20)$$

then we have

$$\mathcal{B}(z) = U(x) + U(y) + U(z) + (\mu - 1)u_0^2. \quad (2.21)$$

We see that $\mathcal{B} > 0$ for $\mu > 1$, i.e., the triples p, q, r are as follows:

- 1) $(1, q, r)$ for any $q, r \in \mathbb{Z}$, i.e., the diagrams A_n ,
- 2) $(2, 2, r)$, the diagrams D_n ,
- 3) $(2, 3, r)$, where $r = 3, 4, 5$, i.e., the diagrams E_6, E_7, E_8 .

Similarly, $\mathcal{B} \geq 0$ for $\mu = 1$, i.e., for the following triples p, q, r :

- 1) $(3, 3, 3)$, the diagram $\tilde{E}_6$,
- 2) $(2, 4, 4)$, the diagram $\tilde{E}_7$,
- 3) $(2, 3, 6)$, the diagram $\tilde{E}_8$.

The Tits form $\mathcal{B}$ is indefinite for $\mu < 1$. Since $U(x)$, $U(y)$, $U(z)$ in (2.21) are positive, we see that, in this case, the signature of $\mathcal{B}$ is equal to $(n-1, 1)$, see [Kac93, exs.4.2]. It is easy to check that the triples p, q, r corresponding to hyperbolic graphs $T_{p,q,r}$ are only the following ones:

- 1) $(2, 3, 7)$, the diagram E_{10} , $1 - \mu = \frac{1}{42}$,
- 2) $(2, 4, 5)$, $1 - \mu = \frac{1}{20}$,
- 3) $(3, 3, 4)$, $1 - \mu = \frac{1}{12}$.

2.6. Orientations and the associated Coxeter transformations

Here, we follow the definitions of Bernstein-Gelfand-Ponomarev [BGP73]. Let us consider the graph Γ endowed with an orientation Δ .

The vertex v_i is called *sink-admissible* (resp. *source-admissible*) in the orientation Δ if every arrow containing v_i ends in (resp. starts from) this vertex. The reflection σ_i can be applied only to vertices which are either sink-admissible or source-admissible. The reflection σ_i acts on the orientation Δ by reversing all arrows containing the vertex v_i .

Consider now a sequence of vertices and the corresponding reflections. A sequence of vertices

$$\{v_{i_n}, v_{i_{n-1}}, \dots, v_{i_3}, v_{i_2}, v_{i_1}\}$$

is called *sink-admissible*, if the vertex v_{i_1} is sink-admissible in the orientation Δ , the vertex v_{i_2} is sink-admissible in the orientation $\sigma_{i_1}(\Delta)$, the vertex v_{i_3} is sink-admissible in the orientation $\sigma_{i_2}\sigma_{i_1}(\Delta)$, and so on. *Source-admissible sequences* are similarly defined.

A sink-admissible (resp. source-admissible) sequence

$$\mathcal{S} = \{v_{i_n}, v_{i_{n-1}}, \dots, v_{i_3}, v_{i_2}, v_{i_1}\}$$

is called *fully sink-admissible* (resp. *fully source-admissible*) if $\mathcal{S}$ contains every vertex $v \in \Gamma_0$ exactly once. Evidently, the inverse sequence $\mathcal{S}^{-1}$ of a sink-admissible sequence $\mathcal{S}$ is source-admissible and vice versa.

Every tree has a fully sink-admissible sequence $\mathcal{S}$. To every sink-admissible sequence $\mathcal{S}$, we assign the Coxeter transformation depending on the order of vertices in $\mathcal{S}$:

$$\mathbf{C} = \sigma_{i_n}\sigma_{i_{n-1}}\dots\sigma_{i_2}\sigma_{i_1}.$$

For every orientation Δ of the tree, every fully sink-admissible sequence gives rise to the same Coxeter transformation $\mathbf{C}_\Delta$, and every fully source-admissible sequence gives rise to $\mathbf{C}_\Delta^{-1}$.

2.7. The Poincaré series

2.7.1. Graded algebras, symmetric algebras, algebras of invariants. Let k be a field. We define a *graded k -algebra* to be a finitely generated k -algebra A (associative, commutative, and with identity), together with a direct sum decomposition (as vector space)

$$A = A_0 \oplus A_1 \oplus A_2 \oplus \dots, \quad (2.22)$$

such that $A_0 = k$ and $A_i A_j \subset A_{i+j}$. The component A_n is called the *n th homogeneous part of A* and the element $x \in A_n$ is said to be *homogeneous of degree n* , notation: $\deg x = n$.

We define a *graded A -module* to be a finitely generated A -module, together with a direct sum decomposition

$$M = M_0 \oplus M_1 \oplus M_2 \oplus \dots, \quad (2.23)$$

such that $A_i M_j \subset M_{i+j}$.

The *Poincaré series* of a graded algebra $A = \bigoplus_{n=1}^{\infty} A_n$ is the formal series

$$P(A, t) = \sum_{n=1}^{\infty} (\dim A_n) t^n. \quad (2.24)$$

The *Poincaré series* of a graded A -module $M = \bigoplus_{n=1}^{\infty} M_n$ is the formal series

$$P(M, t) = \sum_{n=1}^{\infty} (\dim M_n) t^n, \quad (2.25)$$

see [Sp77], [PV94].

THEOREM 2.2. (Hilbert, Serre, see [AtMa69, p.117, Th.11.1]) *The Poincaré series $P(M, t)$ of a finitely generated graded A -module is a rational function in t of the form*

$$P(M, t) = \frac{f(t)}{\prod_{i=1}^s (1 - t^{k_i})}, \text{ where } f(t) \in \mathbb{Z}[t]. \quad (2.26)$$

In what follows in this section, any algebraically closed field k can be considered instead of $\mathbb{C}$. Set

$$R = \mathbb{C}[x_1, \dots, x_n]. \quad (2.27)$$

The set R_d of homogeneous polynomials of degree d is a finite dimensional subspace of R , and $R_0 = \mathbb{C}$. Moreover, $R_d R_e \subset R_{d+e}$, and R is a graded $\mathbb{C}$ -algebra with a direct sum decomposition (as a vector space)

$$R = R_0 \oplus R_1 \oplus R_2 \oplus \dots \quad (2.28)$$

Let $f_1, \dots, f_n \in V^*$ be the linear forms defined by $f_i(x_j) = \delta_{ij}$. Then the f_i , where $i = 1, \dots, n$, generate the *symmetric algebra* $\text{Sym}(V^*)$ isomorphic to R .

For any $a \in GL(V)$ and $f \in R$, define $af \in R$ by the rule

$$(af)(v) = f(a^{-1}v) \text{ for any } v \in V; \quad (2.29)$$

then, for any $a, b \in G$ and $f \in R$, we have

$$a(bf) = bf(a^{-1}v) = f(b^{-1}a^{-1}v) = f((ab)^{-1}v) = ((ab)f)(v),$$

and $aR_d = R_d$.

Let G be a subgroup in $GL(V)$. We say that $f \in R$ is a G -invariant if $af = f$ for all $a \in G$. The G -invariant polynomial functions form a subalgebra R^G of R , which is a graded subalgebra, i.e.,

$$R^G = \bigoplus R^G \cap R_i. \quad (2.30)$$

The algebra R^G is said to be the *algebra of invariants* of the group G . If G is finite, then

$$P(R^G, t) = \frac{1}{|G|} \frac{1}{\sum_{g \in G} \det(1 - gt)}. \quad (2.31)$$

Eq. (2.31) is a classical theorem of Moilen (1897), see [PV94, §3.11], or [Bo, Ch.5, §5.3].

2.7.2. Invariants of finite groups generated by reflections. Let G be a finite subgroup of $GL(V)$ and $g \in G$. Then g is called a *pseudo-reflection* if precisely one eigenvalue of g is not equal to 1. Any pseudo-reflection with determinant -1 is called a *reflection*.

THEOREM 2.3. STCS (Sheppard and Todd, Chevalley, Serre) *Let G be a finite subgroup of $GL(V)$. There exist $n = \dim V$ algebraically independent homogeneous invariants $\theta_1, \dots, \theta_n$ such that*

$$R^G = \mathbb{C}[\theta_1, \dots, \theta_n]$$

if and only if G is generated by pseudo-reflections.

For references, see [Stn79, Th.4.1], [Ch55, Th.A, p.778].

Sheppard and Todd [ShT54] explicitly determined all finite subgroups of $GL(V)$ generated by pseudo-reflections and verified the sufficient condition of Theorem STCS. Chevalley [Ch55] found the classification-free proof of this sufficient condition for a particular case where G is generated by reflections. Serre observed that Chevalley's proof is also valid for groups generated by pseudo-reflections. Sheppard and Todd ([ShT54]) proved the necessary condition of the theorem by a strong combinatorial method; see also Stanley [Stn79, p.487].

The coefficients of the Poincaré series are called the *Betti numbers*. For some bibliographical notes on Betti numbers, see Remark 2.5. The Poincaré polynomial of the algebra R^G is

$$(1 + t^{2p_1+1})(1 + t^{2p_2+1}) \dots (1 + t^{2p_n+1}), \quad (2.32)$$

where $p_i + 1$ are the degrees of homogeneous basis elements of R^G , see [Cox51], and [Ch50]. See Table 2.2 taken from [Cox51, p.781, Tab.4].

TABLE 2.2. The Poincaré polynomials for the simple compact Lie groups

Dynkin diagram	Poincaré polynomial
A_n	$(1 + t^3)(1 + t^5) \dots (1 + t^{2n+1})$
B_n or C_n	$(1 + t^3)(1 + t^7) \dots (1 + t^{4n-1})$
D_n	$(1 + t^3)(1 + t^7) \dots (1 + t^{4n-5})(1 + t^{2n-1})$
E_6	$(1 + t^3)(1 + t^9)(1 + t^{11})(1 + t^{15})(1 + t^{17})(1 + t^{23})$
E_7	$(1 + t^3)(1 + t^{11})(1 + t^{15})(1 + t^{19})(1 + t^{23})(1 + t^{27})(1 + t^{35})$
E_8	$(1 + t^3)(1 + t^{15})(1 + t^{23})(1 + t^{27})(1 + t^{35})(1 + t^{39})(1 + t^{47})(1 + t^{59})$
F_4	$(1 + t^3)(1 + t^{11})(1 + t^{15})(1 + t^{23})$
G_2	$(1 + t^3)(1 + t^{11})$

Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of a Coxeter transformation in a finite Weyl group. These eigenvalues can be given in the form

$$\omega^{m_1}, \dots, \omega^{m_n},$$

where $\omega = \exp^{2\pi i/h}$ is a primitive root of unity. The numbers $m_1, \dots, m_n$ are called the *exponents* of the Weyl group. H. S. M. Coxeter observed that the *exponents* m_i and numbers p_i in (2.32) coincide (see the epigraph to this chapter).

THEOREM 2.4. CCKS (Coxeter, Chevalley, Coleman, Steinberg) *Let*

$$u_1, \dots, u_n$$

be homogeneous elements generating the algebra of invariants R^G , where G is the Weyl group corresponding to a simple compact Lie group. Let $m_i + 1 = \deg u_i$, where $i = 1, 2, \dots, n$. Then the exponents of the group G are

$$m_1, \dots, m_n.$$

For more details, see [Bo, Ch.5, §6.2, Prop.3] and historical notes in [Bo], and [Ch50], [Cox51], [Col58], [Stb85].

REMARK 2.5. *On the Betti numbers.* Here, we mostly follow the works of C. Chevalley [Ch50], and A. J. Coleman [Col58] and review of C. A. Weibel [Weib].

Let X be a topological space, and $H_k(X, \mathbb{Z})$ be the k -th homology group of X . If $H_k(X, \mathbb{Z})$ is finitely generated, then its rank is called the *k -th Betti number* of X .

L. Pontrjagin [**Pon35**] computed the homology of the four classical Lie groups by means of combinatorial arguments; R. Brauer [**Br35**] used de Rham homology. H. Hopf [**Ho41**] and H. Samelson [**Sam41**] showed that, for a compact Lie group $\mathfrak{G}$, the Poincaré series $\sum B_k t^k$ of $\mathfrak{G}$, (where B_k is the k -th Betti number) is of the form

$$P(t) = \prod_{i=1}^n (1 + t^{k_i}). \quad (2.33)$$

The relation (2.33) is called Hopf's theorem [**Ch50**].¹

For the simple compact Lie groups, the Poincaré series $P(t)$ are given in Table 2.2.

J. L. Koszul [**Kosz50**] proved Hopf's theorem in a purely algebraic manner. C. Chevalley [**Ch50**] proved that $k_i = 2p_i + 1$ in (2.33), where $p_i + 1$ is the degree of a minimal homogeneous invariant of the group $\mathfrak{G}$. A. Borel and C. Chevalley [**BC55**] have simplified the calculations for the exceptional Lie groups.

¹As C. A. Weibel [**Weib**, p.6] writes: "...today we would say that Hopf's result amounted to an early classification of finite dimensional graded "Hopf algebras" over $\mathbb{Q}$ ".

CHAPTER 3

The Jordan normal form of the Coxeter transformation

It turned out that most of the classical concepts of the Killing-Cartan-Weyl theory can be carried over to the entire class of Kac-Moody algebras, such as the Cartan subalgebra, the root system, the Weyl group, etc. ... I shall only point out that $\mathfrak{g}'(K)$ ¹ does not always possess a nonzero invariant bilinear form. This is the case if and only if the matrix K is *symmetrizable* ...

V. Kac, [Kac93, p.XI], 1993

3.1. The Cartan matrix and the Coxeter transformation

In this subsection a graph Γ and a partition $S = S_1 \coprod S_2$ of its vertices are fixed.

3.1.1. A bicolored partition and a bipartite graph. A partition $S = S_1 \coprod S_2$ of the vertices of the graph Γ is said to be *bicolored* if all edges of Γ lead from S_1 to S_2 . A *bicolored partition* exists if and only if all cycles in Γ are of even order. The graph Γ admitting a bicolored partition is said to be *bipartite* [McM02]. An orientation Δ is said to be *bicolored*, if there is the corresponding sink-admissible sequence

$$\{v_1, v_2, \dots, v_m, v_{m+1}, v_{m+2}, \dots, v_{m+k}\}$$

of vertices in this orientation Δ , such that subsequences

$$S_1 = \{v_1, v_2, \dots, v_m\} \text{ and } S_2 = \{v_{m+1}, v_{m+2}, \dots, v_{m+k}\}$$

form a bicolored partition. Any two generators g' and g'' of the Coxeter group $W(S)$ (contained in the same subpartition) commute and the subgroups $W(S_1)$ and $W(S_2)$ are abelian. So, the products $w_i \in W(S_i)$ for $i = 1, 2$ of generators of $W(S_i)$ are involutions, i.e.,

$$w_1^2 = 1, \quad w_2^2 = 1.$$

For the first time (as far as I know), the technique of bipartite graphs was used by R. Steinberg in [Stb59], where he gave classification-free proofs for some results of H. S. M. Coxeter [Cox34] concerning properties of the order of the Coxeter transformation; see also R. Carter's paper [Car70].

¹Here, $\mathfrak{g}'(K)$ is the subalgebra $[\mathfrak{g}(K), \mathfrak{g}(K)]$ of the Kac-Moody algebra $\mathfrak{g}(K)$ associated with the generalized Cartan matrix K . One has

$$\mathfrak{g}(K) = \mathfrak{g}'(K) + \mathfrak{h},$$

where $\mathfrak{h}$ is the Cartan subalgebra, $\mathfrak{g}(K) = \mathfrak{g}'(K)$ if and only if $\det K \neq 0$, [Kac93, §1.3 and Th.2.2].

3.1.2. Conjugacy of Coxeter transformations. All Coxeter transformations are conjugate for any tree or forest Γ [Bo, Ch.5, §6]; see also Proposition 6.9 and Remark 6.10. The Coxeter transformations for the graphs with cycles are studied in [Col89], [Rin94], [Shi00], [BT97], and in the works by Menshikh and Subbotin of 1982–1985, see §4.2. Here, we consider only trees.

Let us select one Coxeter transformation most simple to study. It is one of the two *bicolored Coxeter transformations* corresponding to a bipartite graph:

$$\mathbf{C} = w_1 w_2 \quad \text{or} \quad \mathbf{C}^{-1} = w_2 w_1. \quad (3.1)$$

From now on we assume that S_1 contains m elements and S_2 contains k elements, we denote by $a_1, \dots, a_m$ (resp. $b_1, \dots, b_k$) basis vectors corresponding to vertices $v_1, \dots, v_m$ of S_1 (resp. vertices $v_{m+1}, \dots, v_{m+k}$ of S_2). We denote by $\mathcal{E}_{\Gamma_a}$ (resp. $\mathcal{E}_{\Gamma_b}$) the vector space generated by the a_i , where $i = 1, \dots, m$ (resp. by the b_i , where $i = 1, \dots, k$). So,

$$\dim \mathcal{E}_{\Gamma_a} = m, \quad \dim \mathcal{E}_{\Gamma_b} = k.$$

3.1.3. The Cartan matrix and the bicolored Coxeter transformation. The Cartan matrix and the bicolored Coxeter transformation are constructed from the same blocks. More exactly, the matrix $\mathbf{B}$ and involutions w_i , where $i = 1, 2$, are constructed from the same blocks.

In the simply-laced case (i.e., for the symmetric Cartan matrix), we have

$$\begin{aligned} K = 2\mathbf{B}, \quad \text{where} \quad \mathbf{B} &= \begin{pmatrix} I_m & D \\ D^t & I_k \end{pmatrix}, \\ w_1 &= \begin{pmatrix} -I_m & -2D \\ 0 & I_k \end{pmatrix}, \quad w_2 = \begin{pmatrix} I_m & 0 \\ -2D^t & -I_k \end{pmatrix}, \end{aligned} \quad (3.2)$$

where the elements d_{ij} that constitute matrix D are given by the formula

$$d_{ij} = (a_i, b_j) = \begin{cases} -\frac{1}{2} & \text{if } |a_i - b_j| = 1, \\ 0 & \text{if } |a_i - b_j| > 1. \end{cases} \quad (3.3)$$

In the multiply-laced case (i.e., for the symmetrizable and non-symmetric Cartan matrix K), we have

$$\begin{aligned} K = T\mathbf{B}, \quad \text{where} \quad K &= \begin{pmatrix} 2I_m & 2D \\ 2F & 2I_k \end{pmatrix}, \\ w_1 &= \begin{pmatrix} -I_m & -2D \\ 0 & I_k \end{pmatrix}, \quad w_2 = \begin{pmatrix} I_m & 0 \\ -2F & -I_k \end{pmatrix}, \end{aligned} \quad (3.4)$$

and where

$$d_{ij} = \frac{(a_i, b_j)}{(a_i, a_i)}, \quad f_{pq} = \frac{(b_p, a_q)}{(b_p, b_p)}, \quad (3.5)$$

where the a_i and b_j are simple roots in the root systems of the corresponding to S_1 and S_2 Kac-Moody Lie algebras [Kac80]. Let $T = (t_{ij})$ be the diagonal matrix (2.6). Then

$$(t_{ii}) = \frac{1}{(a_i, a_i)} = \frac{1}{\mathcal{B}(a_i)}.$$

Dividing T and $\mathbf{B}$ into blocks of size $m \times m$ and $k \times k$, we see that

$$T = \begin{pmatrix} 2T_1 & 0 \\ 0 & 2T_2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} T_1^{-1} & A \\ A^t & T_2^{-1} \end{pmatrix}, \quad T_1 A = D, \quad T_2 A^t = F. \quad (3.6)$$

3.1.4. Dual graphs and Dual forms. Every valued graph Γ has a dual graph denoted by $\Gamma^\vee$. The dual graph is obtained by means of transposition $d_{ij} \leftrightarrow d_{ji}$. In other words, if K is the Cartan matrix for Γ , then the Cartan matrix $K^\vee$ for $\Gamma^\vee$ is K^t ,

$$K^\vee = K^t,$$

i.e.,

$$F^\vee = D^t, \quad D^\vee = F^t. \quad (3.7)$$

Therefore,

$$F^\vee D^\vee = (FD)^t, \quad D^\vee F^\vee = (DF)^t. \quad (3.8)$$

For any simply-laced graph, the Cartan matrix is symmetric and $F = D^t = F^\vee$. In this case the graph Γ is dual to itself. Among extended Dynkin diagrams the following pairs of diagram are dual:

$$(\tilde{B}_n, \tilde{C}_n), \quad (\widetilde{CD}_n, \widetilde{DC}_n), \quad (\tilde{G}_{21}, \tilde{G}_{12}), \quad (\tilde{F}_{41}, \tilde{F}_{42}).$$

Let dual Cartan matrices be factorized by means of the diagonal matrices T and $T^\vee$:

$$K = T\mathbf{B}, \quad K^\vee = T^\vee\mathbf{B}^\vee.$$

Then according to [Kac93, Ch.3, exs.3.1] we have

$$T^\vee = T^{-1}. \quad (3.9)$$

Since $K^\vee = K^t = \mathbf{B}T = T^{-1}\mathbf{B}^\vee$, we see that dual Tits forms are related as follows:

$$\mathbf{B}^\vee = T\mathbf{B}T. \quad (3.10)$$

3.1.5. Eigenvalues of the Cartan matrix and the Coxeter transformation.

There is a simple relation between eigenvalues of the Cartan matrix and the Coxeter transformation. Let vector $z \in \mathcal{E}_\Gamma$ be given in two-component form

$$z = \begin{pmatrix} x \\ y \end{pmatrix}, \quad (3.11)$$

where $\dim \mathcal{E}_\Gamma = k + m = n$, $x \in \mathcal{E}_{\Gamma_a}$ and $y \in \mathcal{E}_{\Gamma_b}$. Consider the relation

$$\mathbf{C}z = \lambda z \quad \text{or} \quad w_2 z = \lambda w_1 z.$$

In the simply-laced case we deduce from (3.2) that

$$\mathbf{C}z = \lambda z \iff \begin{cases} \frac{\lambda+1}{2\lambda}x = Dy \\ \frac{\lambda+1}{2}y = D^t x \end{cases} \iff \mathbf{B}z = \frac{\lambda-1}{2} \begin{pmatrix} \frac{1}{\lambda}x \\ -y \end{pmatrix}. \quad (3.12)$$

In the multiply-laced case we deduce that

$$\mathbf{C}z = \lambda z \iff \begin{cases} \frac{\lambda+1}{2\lambda}x = Dy \\ \frac{\lambda+1}{2}y = Fx \end{cases} \iff \mathbf{B}z = \frac{\lambda-1}{2} \begin{pmatrix} \frac{1}{\lambda}x \\ -y \end{pmatrix}. \quad (3.13)$$

From (3.12) and (3.13) we have in the simply-laced and multiply-laced cases, respectively:

$$\begin{cases} DD^t x = \frac{(\lambda+1)^2}{4\lambda}x \\ D^t Dy = \frac{(\lambda+1)^2}{4\lambda}y \end{cases} \iff \begin{cases} DFx = \frac{(\lambda+1)^2}{4\lambda}x \\ FDy = \frac{(\lambda+1)^2}{4\lambda}y \end{cases} \quad (3.14)$$

Similarly,

$$\mathbf{B}z = \gamma z \iff \begin{cases} (\gamma - 1)x = Dy \\ (\gamma - 1)y = D^t x \end{cases} \text{ resp. } \begin{cases} (\gamma - 1)x = Dy \\ (\gamma - 1)y = Fx; \end{cases} \quad (3.15)$$

and from (3.15) we have

$$\begin{cases} DD^t x = (\gamma - 1)^2 x \\ D^t Dy = (\gamma - 1)^2 y \end{cases} \text{ resp. } \begin{cases} DFx = (\gamma - 1)^2 x \\ FDy = (\gamma - 1)^2 y. \end{cases} \quad (3.16)$$

By (2.6) we have

PROPOSITION 3.1 (Fixed Points). *1) The eigenvalues λ of the Coxeter transformation and the eigenvalues γ of the matrix $\mathbf{B}$ of the Tits form are related as follows*

$$\frac{(\lambda + 1)^2}{4\lambda} = (\gamma - 1)^2.$$

2) The kernel of the matrix $\mathbf{B}$ coincides with the kernel of the Cartan matrix K and coincides with the space of fixed points of the Coxeter transformation

$$\ker K = \ker \mathbf{B} = \{z \mid \mathbf{C}z = z\}.$$

3) The space of fixed points of matrix $\mathbf{B}$ coincides with the space of anti-fixed points of the Coxeter transformation

$$\{z \mid \mathbf{B}z = z\} = \{z \mid \mathbf{C}z = -z\}.$$

For more information about fixed and anti-fixed points of the powers of the Coxeter transformation, see Appendix B.7.1.

3.2. An application of the Perron-Frobenius theorem

3.2.1. The pair of matrices DD^t and D^tD (resp. DF and FD).

REMARK 3.2. The matrices DD^t and D^tD have some nice properties.

(1) They give us all information about the eigenvalues of Coxeter transformations and Cartan matrices. Our results hold for an arbitrary tree Γ .

(2) The eigenvectors of Coxeter transformations are combinations of eigenvectors of the matrix DD^t and eigenvectors of the matrix D^tD , see §3.3.1, relations (3.29), (3.30), (3.31).

(3) They satisfy the Perron-Frobenius theorem.

More exactly, properties (1)–(3) will be considered below in Proposition 3.3, Proposition 3.5, and in §3.2.2, §3.3.1.

PROPOSITION 3.3. *1) The matrices DD^t and D^tD (resp. DF and FD) have the same non-zero eigenvalues with equal multiplicities.*

2) The eigenvalues φ_i of matrices DD^t and D^tD (resp. DF and FD) are non-negative:

$$\varphi_i \geq 0.$$

3) The corresponding eigenvalues $\lambda_{1,2}^{\varphi_i}$ of Coxeter transformations are

$$\lambda_{1,2}^{\varphi_i} = 2\varphi_i - 1 \pm 2\sqrt{\varphi_i(\varphi_i - 1)}. \quad (3.17)$$

The eigenvalues $\lambda_{1,2}^{\varphi_i}$ either lie on the unit circle or are real positive numbers. In the latter case $\lambda_1^{\varphi_i}$ and $\lambda_2^{\varphi_i}$ are mutually inverse:

$$\lambda_1^{\varphi_i} \lambda_2^{\varphi_i} = 1.$$

Proof. 1) If $DD^t z = \mu z$, where $\mu \neq 0$, then $D^t z \neq 0$ and $D^t D(D^t z) = \mu(D^t z)$. We argue similarly for $D^t D$, DF , FD . The multiplicities of non-zero eigenvalues coincide since

$$x \neq 0, \quad DD^t x = \mu x \implies D^t D(D^t x) = \mu(D^t x), \text{ and } D^t x \neq 0,$$

and

$$y \neq 0, \quad D^t D y = \mu y \implies DD^t(Dy) = \mu(Dy), \text{ and } Dy \neq 0.$$

REMARK 3.4. The multiplicities of the zero eigenvalue are not equal. If $DD^t x = 0$ and x is the eigenvector, $x \neq 0$, then it is possible that $D^t x = 0$ and $D^t x$ is not an eigenvector of $D^t D$, see Remark 3.6, 4) below.

2) The matrices DD^t and $D^t D$ are symmetric and nonnegative definite. For example,

$$\langle DD^t x, x \rangle = \langle D^t x, D^t x \rangle \geq 0.$$

So, if $DD^t x = \varphi_i x$, then

$$\langle DD^t x, x \rangle = \varphi_i \langle x, x \rangle$$

and

$$\varphi_i = \frac{\langle D^t x, D^t x \rangle}{\langle x, x \rangle} \geq 0.$$

In the multiply-laced case, we deduce from (3.6) that the matrix DF is

$$DF = T_1 A T_2 A^t. \quad (3.18)$$

Let φ be a non-zero eigenvalue for $DF = T_1 A T_2 A^t$ with eigenvector x :

$$T_1 A T_2 A^t x = \varphi x.$$

Since T is a positive diagonal matrix, see §2.1, we have

$$(\sqrt{T_1} A T_2 A^t \sqrt{T_1})((\sqrt{T_1})^{-1} x) = \varphi(\sqrt{T_1})^{-1} x, \quad (3.19)$$

and φ is also a non-zero eigenvalue with eigenvector $(\sqrt{T_1})^{-1} x$ for the matrix $\sqrt{T_1} A T_2 A^t \sqrt{T_1}$ which already is symmetric, so $\varphi \geq 0$.

3) From (3.17) if $0 \leq \varphi \leq 1$ we deduce that

$$|\lambda_{1,2}^{\varphi_i}|^2 = (2\varphi_i - 1)^2 + 4\varphi_i(1 - \varphi_i) = 1.$$

If $\varphi_i > 1$, then

$$2\varphi_i - 1 > 2\sqrt{\varphi_i(\varphi_i - 1)} \implies \lambda_{1,2}^{\varphi_i} \geq 0.$$

Thus,

$$\begin{aligned} \lambda_1^{\varphi_i} &= 2\varphi_i - 1 + 2\sqrt{\varphi_i(\varphi_i - 1)} > 1, \\ \lambda_2^{\varphi_i} &= 2\varphi_i - 1 - 2\sqrt{\varphi_i(\varphi_i - 1)} < 1. \quad \square \end{aligned} \quad (3.20)$$

The pair of matrices (A, B) is said to be a *PF-pair* if matrices A and B satisfy the Perron-Frobenius theorem¹.

PROPOSITION 3.5. *The matrix pair $(DD^t, D^t D)$ (resp. (DF, FD)) is a PF-pair, i.e.,*

- 1) *DD^t and $D^t D$ (resp. DF and FD) are non-negative;*
- 2) *DD^t and $D^t D$ (resp. DF and FD) are indecomposable.*

¹The Perron-Frobenius theorem is well known in the matrix theory, see §B.3 and [MM64], [Ga90].

Proof. 1) Indeed, in the simply-laced case, the following relation holds

$$4(DD^t)_{ij} = 4 \sum_{p=1}^k (a_i, b_p)(b_p, a_j) = \begin{cases} s_i & \text{the number of edges with a vertex } v_i \text{ if } i = j, \\ 1 & \text{if } |v_i - v_j| = 2, \\ 0 & \text{if } |v_i - v_j| > 2. \end{cases} \quad (3.21)$$

In the multiply-laced case, we have

$$4(DF)_{ij} = 4 \sum_{p=1}^k \frac{(a_i, b_p)(b_p, a_j)}{(a_i, a_i)(b_p, b_p)} = \begin{cases} 4 \sum_{p=1}^k \frac{(a_i, b_p)^2}{(a_i, a_i)(b_p, b_p)} = 4 \sum_{p=1}^k \cos^2\{a_i, b_p\} & \text{if } i = j, \\ 4 \frac{(a_i, b_p)(b_p, a_j)}{(a_i, a_i)(b_p, b_p)} & \text{if } |v_i - v_j| = 2, \\ 0 & \text{if } |v_i - v_j| > 2. \end{cases} \quad (3.22)$$

2) Define the distance between two sets of vertices $A = \{a_i\}_{i \in I}$ and $B = \{b_j\}_{j \in J}$ to be

$$\min_{i,j} |a_i - b_j|.$$

If the matrix DD^t is decomposable, then the set of vertices $\{v_1, \dots, v_n\}$ can be divided into two subsets such that distance between these two subsets is > 2 . This contradicts the assumption that Γ is connected. $\square$

REMARK 3.6. 1) Formula (3.21) is a particular case of (3.22), since the angles between the adjacent simple roots a_i and b_j are $\frac{2\pi}{3}$, so $\cos\{a_i, b_j\} = \frac{1}{2}$. Of course, the angles and the lengths of vectors are considered in the sense of the bilinear form $(\cdot, \cdot)$ from (2.4).

2) The case $|v_i - v_j| = 2$ from (3.22) can be expressed in the following form:

$$4 \frac{(a_i, b_p)(b_p, a_j)}{(a_i, a_i)(b_p, b_p)} = 4 \frac{(a_i, b_p)(b_p, a_j)}{|a_i|^2 |b_p|^2} = 4 \frac{(a_i, b_p)}{|a_i|} \frac{(b_p, a_j)}{|b_p|} \frac{|a_j|}{|a_i|} = 4 \frac{|a_j|}{|a_i|} \cos\{a_i, b_p\} \cos\{a_j, b_p\}. \quad (3.23)$$




FIGURE 3.1. The star $*_{k+1}$ with k rays

3) One can easily calculate the matrices DD^t, D^tD, DF, FD by means of (3.21) and (3.22).

4) Consider a simple star $*_{k+1}$, Fig. 3.1. It is bipartite with respect to the following bicolored partition. One part of the graph consists of only one vertex a_1 , ($m = 1$), the other one consists of k vertices $\{b_1, \dots, b_k\}$, $n = k + 1$. According to (3.21) the 1×1 matrix DD^t is

$$DD^t = k = n - 1,$$

and the $k \times k$ matrix D^tD is

$$D^tD = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & 1 & 1 & \dots & 1 \\ 1 & 1 & 1 & \dots & 1 \\ & & & \dots & \\ 1 & 1 & 1 & \dots & 1 \end{pmatrix}.$$

By Proposition 3.3, heading 2) the matrices DD^t and D^tD have only one non-zero eigenvalue $\varphi_1 = n - 1$. All the other eigenvalues of D^tD are zeros and the characteristic polynomial of the D^tD is

$$\varphi^{n-1}(\varphi - (n - 1)).$$

3.2.2. The Perron-Frobenius theorem applied to DD^t and D^tD (resp. DF and FD). By Proposition 3.5 the pairs (DD^t, D^tD) (resp. (DF, FD)) are PF-pairs, so we can apply the Perron-Frobenius theorem, see §B.3.

COROLLARY 3.7. *The matrices DD^t and D^tD (resp. DF and FD) have a common simple (i.e., with multiplicity one) positive eigenvalue φ_1 . This eigenvalue is maximal (called dominant eigenvalue):*

$$0 \leq \varphi_i \leq \varphi_1, \quad \varphi_1 = \begin{cases} \max_{x \geq 0} \min_{1 \leq i \leq m} \frac{(DD^t x)_i}{x_i} & \text{in the simply-laced case,} \\ \max_{x \geq 0} \min_{1 \leq i \leq m} \frac{(DF x)_i}{x_i} & \text{in the multiply-laced case.} \end{cases} \quad (3.24)$$

There are positive eigenvectors $\mathbb{X}^{\varphi_1}$, $\mathbb{Y}^{\varphi_1}$ (i.e., non-zero vectors with non-negative coordinates) corresponding to the eigenvalue φ_1 :

$$\begin{aligned} DD^t \mathbb{X}^{\varphi_1} &= \varphi_1 \mathbb{X}^{\varphi_1}, & D^tD \mathbb{Y}^{\varphi_1} &= \varphi_1 \mathbb{Y}^{\varphi_1}, \\ DF \mathbb{X}^{\varphi_1} &= \varphi_1 \mathbb{X}^{\varphi_1}, & FD \mathbb{Y}^{\varphi_1} &= \varphi_1 \mathbb{Y}^{\varphi_1}. \end{aligned} \quad (3.25)$$


The matrices DD^t (resp. D^tD) are symmetric and can be diagonalized in the some orthonormal basis of the eigenvectors from $\mathcal{E}_{\Gamma_a} = \mathbb{R}^h$ (resp. $\mathcal{E}_{\Gamma_b} = \mathbb{R}^k$). The Jordan normal forms of these matrices are shown on Fig. 3.2. The normal forms of DF and FD are the same, since by (3.18), (3.19) the non-zero eigenvalues of the matrix DF and symmetric matrix $\sqrt{T_1}AT_2A^t\sqrt{T_1}$ coincide. However, the normal bases (i.e., bases which consist of eigenvectors) for DF and FD are not necessarily orthonormal, since the eigenvectors of DF are obtained from eigenvectors of $\sqrt{T_1}AT_2A^t\sqrt{T_1}$ by means of the matrix $\sqrt{T_1}$ which does not preserve orthogonality.

3.3. The basis of eigenvectors and a theorem on the Jordan form

3.3.1. An explicit construction of the eigenvectors.

PROPOSITION 3.8. *Let*

$$\mathbb{X}_1^{\varphi_1}, \mathbb{X}_1^{\varphi_2}, \dots, \mathbb{X}_{t_2}^{\varphi_2}, \dots, \mathbb{X}_1^{\varphi_s}, \dots, \mathbb{X}_{t_s}^{\varphi_s}, \mathbb{X}_1^0, \dots, \mathbb{X}_{m-p}^0 \quad (3.26)$$

FIGURE 3.2. The Jordan normal forms of DD^t and D^tD

be all the orthonormal eigenvectors for DD^t and

$$\mathbb{Y}_1^0, \dots, \mathbb{Y}_{k-p}^0 \quad (3.27)$$

be all the orthonormal eigenvectors for D^tD corresponding to the zero eigenvalue. Then

$$D^t\mathbb{X}_1^{\varphi_1}, D^t\mathbb{X}_1^{\varphi_2}, \dots, D^t\mathbb{X}_{t_2}^{\varphi_2}, \dots, D^t\mathbb{X}_1^{\varphi_s}, \dots, D^t\mathbb{X}_{t_s}^{\varphi_s}, \mathbb{Y}_1^0, \dots, \mathbb{Y}_{k-p}^0 \quad (3.28)$$

is the set of all orthonormal eigenvectors for DD^t .

Bases for DF, FD (not orthonormal) are similarly constructed.

Proof. Indeed, if $\mathbb{X}^{\varphi_i}$ and $\mathbb{X}^{\varphi_j}$ are the eigenvectors corresponding to the φ_i and φ_j , then the vectors $\mathbb{Y}^{\varphi_i} = D^t\mathbb{X}^{\varphi_i}$ and $\mathbb{Y}^{\varphi_j} = D^t\mathbb{X}^{\varphi_j}$ are eigenvectors of D^tD and

$$\langle \mathbb{Y}^{\varphi_i}, \mathbb{Y}^{\varphi_j} \rangle = \langle \mathbb{X}^{\varphi_i}, DD^t\mathbb{X}^{\varphi_j} \rangle = \varphi_j \langle \mathbb{X}^{\varphi_i}, \mathbb{X}^{\varphi_j} \rangle = 0. \quad \square$$

Let us construct the eigenvectors for the Coxeter transformation. We set:

Case $\varphi_i \neq 0, 1$:

$$z_{r,\nu}^{\varphi_i} = \begin{pmatrix} \mathbb{X}_r^{\varphi_i} \\ -\frac{2}{\lambda_{\nu}^{\varphi_i} + 1} D^t\mathbb{X}_r^{\varphi_i} \end{pmatrix}, \quad 1 \leq i \leq s, \quad 1 \leq r \leq t_i, \quad \nu = 1, 2. \quad (3.29)$$

Here $\lambda_{1,2}^{\varphi_i}$ is obtained by formula (3.17).

Case $\varphi_i = 1$:

$$z_r^1 = \begin{pmatrix} \mathbb{X}_r^1 \\ -D^t\mathbb{X}_r^1 \end{pmatrix}, \quad \tilde{z}_r^1 = \frac{1}{4} \begin{pmatrix} \mathbb{X}_r^1 \\ D^t\mathbb{X}_r^1 \end{pmatrix}, \quad 1 \leq r \leq t_i. \quad (3.30)$$

Case $\varphi_i = 0$:

$$z_{x_\eta}^0 = \begin{pmatrix} \mathbb{X}_\eta^0 \\ 0 \end{pmatrix}, \quad 1 \leq \eta \leq m-p, \quad z_{y_\xi}^0 = \begin{pmatrix} 0 \\ \mathbb{Y}_\xi^0 \end{pmatrix}, \quad 1 \leq \xi \leq k-p. \quad (3.31)$$

PROPOSITION 3.9 ([SuSt75, SuSt78]). 1) The vectors (3.29), (3.30), (3.31) constitute a basis in $\mathcal{E}_\Gamma$ over $\mathbb{C}$.

2) The vectors (3.29) are eigenvectors of the Coxeter transformation corresponding to the eigenvalue λ_{φ_i} :

$$\mathbf{C}z_{r,\nu}^{\varphi_i} = \lambda_{\varphi_i} z_{r,\nu}^{\varphi_i}. \quad (3.32)$$

The vectors (3.30) are eigenvectors and adjoint vectors of the Coxeter transformation corresponding to the eigenvalue 1:

$$\mathbf{C}z_r^1 = z_r^1, \quad \mathbf{C}\tilde{z}_r^1 = z_r^1 + \tilde{z}_r^1. \quad (3.33)$$

The vectors (3.31) are eigenvectors of the Coxeter transformation corresponding to the eigenvalue -1 :

$$\mathbf{C}z_{x_\eta}^0 = -z_{x_\eta}^0, \quad \mathbf{C}z_{y_\xi}^0 = -z_{y_\xi}^0. \quad (3.34)$$

In other words, vectors (3.29), (3.30), (3.31) constitute an orthogonal basis which consists of eigenvectors and adjoint vectors of the Coxeter transformation. The number of the adjoint vectors $\tilde{z}_r^1$ is equal to the multiplicity of eigenvalue $\varphi = 1$.

Proof. 1) The number of vectors (3.29), (3.30), (3.31) is $2p + (m - p) + (k - p) = n$ and it suffices to prove that these vectors are linearly independent. Let us write down the condition of linear dependence. It splits into 2 conditions: for the $\mathbb{X}$ -component and for the $\mathbb{Y}$ -component. The linear independence of vectors (3.29), (3.30), (3.31) follows from the linear independence of vectors (3.26), (3.28). $\square$

2) To prove relation (3.32), the first relation from (3.33) and relation (3.34), it suffices to check (3.12). Let us check that

$$\mathbf{C}\tilde{z}_r^1 = z_r^1 + \tilde{z}_r^1.$$

We consider the multiply-laced case. Making use of (3.4) we see that

$$\mathbf{C} = \begin{pmatrix} 4DF - I_m & 2D \\ -2F & I_k \end{pmatrix}.$$

Then

$$\mathbf{C}\tilde{z}_r^1 = \frac{1}{4} \begin{pmatrix} 4DF\mathbb{X}_r^1 - \mathbb{X}_r^1 + 2DF\mathbb{X}_r^1 \\ -2F\mathbb{X}_r^1 - F\mathbb{X}_r^1 \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 5\mathbb{X}_r^1 \\ -3\mathbb{X}_r^1 \end{pmatrix} = z_r^1 + \tilde{z}_r^1. \quad \square$$

By heading 2) of Proposition 3.1 and formula (3.30) from Proposition 3.8 we have

COROLLARY 3.10. *The Jordan normal form of the Coxeter transformation is diagonal if and only if the Tits form is nondegenerate*². $\square$

3.3.2. Monotonicity of the dominant eigenvalue. The following proposition is important for calculation of the number of 2×2 Jordan blocks in the Jordan normal form of the Coxeter transformation.

PROPOSITION 3.11 ([SuSt75, SuSt78]). *Let us add an edge to some tree Γ and let $\hat{\Gamma}$ be the new graph (Remark 2.1). Then:*

1) *The dominant eigenvalue φ_1 may only grow:*

$$\varphi_1(\hat{\Gamma}) \geq \varphi_1(\Gamma).$$

2) *Let Γ be an extended Dynkin diagram, i.e., $\mathcal{B} \geq 0$. Then the spectra of $DD^t(\hat{\Gamma})$ and $D^tD(\hat{\Gamma})$ (resp. $DF(\hat{\Gamma})$ and $FD(\hat{\Gamma})$) do not contain 1, i.e.,*

$$\varphi_i(\hat{\Gamma}) \neq 1 \quad \text{for all } \varphi_i \in \text{Spec}(DD^t(\hat{\Gamma})).$$

²For the Jordan canonical (normal) form, see, for example, [Ga90, Ch.VI] or [Pr94, Ch.III].

3) Let $\mathcal{B}$ be indefinite. Then

$$\varphi_1(\hat{\Gamma}) > 1.$$

Proof. 1) Adding an edge to the vertex a_i we see, according to (3.21), that only one element of DD^t changes: namely, $(DD^t)_{ii}$ changes from $\frac{s_i}{4}$ to $\frac{s_i+1}{4}$. For the multiply-laced case, $(DF)_{ii}$ changes by $\cos^2(a_i, b_s)$, where b_s is the new vertex connected with the vertex a_i . By Corollary 3.7 we have

$$\begin{aligned} \varphi_1(\hat{\Gamma}) &= \max_{x \geq 0} \min_{1 \leq i \leq m} \frac{(DF(\hat{\Gamma})x)_i}{x_i} \geq \\ &\min_{1 \leq i \leq m} \frac{(DF(\hat{\Gamma})\mathbb{X}_i^{\varphi_1})_i}{\mathbb{X}_i^{\varphi_1}} \geq \min_{1 \leq i \leq m} \frac{(DF\mathbb{X}_i^{\varphi_1})_i}{\mathbb{X}_i^{\varphi_1}} = \varphi_1. \end{aligned}$$

2) The characteristic polynomial of $DF(\hat{\Gamma})$ is

$$\det |DF(\hat{\Gamma}) - \mu I| = \det |DF - \mu I| + \cos^2\{a_i, b_s\} \det |A_i(\mu)|, \quad (3.35)$$

where $A_i(\mu)$ is obtained by deleting the i^{th} row and i^{th} column from the matrix $DF - \mu I$. It corresponds to the operation “*Remove Vertex*” from the graph Γ (Remark 2.1), the graph obtained by removing vertex is $\check{\Gamma}$, i.e.,

$$\det |DF(\hat{\Gamma}) - \mu I| = \det |DF - \mu I| + \cos^2\{a_i, b_s\} \det |DF(\check{\Gamma}) - \mu I|. \quad (3.36)$$

According to Remark 2.1 the quadratic form $\mathcal{B}(\check{\Gamma})$ is positive, $\check{\Gamma}$ is the Dynkin diagram, i.e., $\ker \mathbf{B} = 0$. By Proposition 3.1 the Coxeter transformation for $\check{\Gamma}$ does not have eigenvalue 1. Then by (3.17) the corresponding matrix $DF(\check{\Gamma})$ does not have eigenvalue 1. Thus, in (3.36), $\mu = 1$ is a root of $\det |DF - \mu I|$ and is not a root of $\det |DF(\check{\Gamma}) - \mu I|$, and therefore $\mu = 1$ is not a root of $\det |DF(\hat{\Gamma}) - \mu I|$. The case $\cos^2\{a_i, b_s\} = 0$ is not possible since $\hat{\Gamma}$ is connected.

3) By heading 1) adding only one edge to an extended Dynkin diagram we get $\varphi_1(\hat{\Gamma}) \geq \varphi_1$. By heading 2) $\varphi_1(\hat{\Gamma}) > \varphi_1$. But the form $\mathcal{B}$ becomes indefinite after we add some edges to the extended Dynkin diagram (Remark 2.1, 4). Thus, φ_1 is only grows. $\square$

PROPOSITION 3.12. *The common dominant eigenvalue of DD^t and D^tD (resp. DF and FD) is equal to 1 if and only if Γ is an extended Dynkin diagram.*

Proof. 1) Let Γ be an extended Dynkin diagram, then $\mathcal{B} \geq 0$. Since $\ker \mathbf{B} \neq 0$, we see by Proposition 3.1 that the eigenvalue φ_1 is the eigenvalue of the matrices DF and FD . By (3.20) we have $\lambda_1^{\varphi_1} \geq 1$. Further, since the Weyl group preserves the quadratic form $\mathcal{B}$, we have

$$\mathcal{B}(z^{\varphi_1}) = \mathcal{B}(\mathbf{C}z^{\varphi_1}) = (\lambda_1^{\varphi_1})^2 \mathcal{B}(\mathbf{C}z^{\varphi_1}).$$

Therefore, either $\lambda_1^{\varphi_1} = 1$, i.e., $\varphi_1 = 1$, or $\mathcal{B}(z^{\varphi_1}) = 0$. We will show that in the latter case $\varphi_1 = 1$, too. By Proposition 3.8 the vectors $\mathbb{X}^{\varphi_1}$ and $F\mathbb{X}^{\varphi_1}$ have real coordinates. By (3.20) the eigenvalue $\lambda_1^{\varphi_1}$ is also real because $\varphi_1 \geq 1$. So, the vector z^{φ_1} from (3.30) is real. Then from $\mathcal{B}(z^{\varphi_1}) = 0$ we have $z^{\varphi_1} \in \ker \mathbf{B}$ and again, $\lambda_1^{\varphi_1} = 1$.

2) Conversely, let $\lambda_1^{\varphi_1} = 1$. Then, by Proposition 3.1, $\ker \mathbf{B} \neq 0$, i.e., the form $\mathcal{B}$ is degenerate. Let us find whether $\mathcal{B}$ is definite ($\mathcal{B} \geq 0$) or indefinite. By heading 3) of

Proposition 3.11 if $\mathcal{B}$ is indefinite, then $\lambda_1^{\varphi_1} > 1$. Thus, $\mathcal{B} \geq 0$ and Γ is an extended Dynkin diagram. $\square$

COROLLARY 3.13. *Let $\mathcal{B} \geq 0$, i.e., let Γ be an extended Dynkin diagram. Then*

- 1) *The kernel of the quadratic form $\mathcal{B}$ is one-dimensional.*
- 2) *The vector $z^{\varphi_1} \in \ker \mathbf{B}$ can be chosen so that all its coordinates are positive.*

Proof. 1) φ_1 is a simple eigenvalue of DD^t and D^tD (see Corollary 3.7).

2) $\mathbb{X}^{\varphi_1}$ is a positive vector (Corollary 3.7), $D^t\mathbb{X}^{\varphi_1}$ is a negative vector since D^t is a nonpositive matrix (3.3). So, the vector z^{φ_1} from (3.30) is positive. $\square$

The monotonicity of the dominant value φ_1 and the corresponding maximal eigenvalue λ^{φ_1} of the Coxeter transformation is clearly demonstrated for the diagrams $T_{2,3,r}$, $T_{3,3,r}$, $T_{2,4,r}$, see Propositions 4.15, 4.17, 4.19 and Tables 4.4, 4.5, 4.6 in §4.4.

3.3.3. A theorem on the Jordan form.

Now we can summarize.

THEOREM 3.14 ([SuSt75, SuSt78, St85]). 1) *The Jordan form of the Coxeter transformation is diagonal if and only if the Tits form is non-degenerate.*

2) *If $\mathcal{B} \geq 0$ (Γ is an extended Dynkin diagram), then the Jordan form of the Coxeter transformation contains only one 2×2 Jordan block. All eigenvalues λ_i lie on the unit circle.*

3) *If $\mathcal{B}$ is indefinite and degenerate, then the number of 2×2 Jordan blocks coincides with $\dim \ker \mathbf{B}$. The remaining Jordan blocks are 1×1 . There is a simple maximal eigenvalue $\lambda_1^{\varphi_1}$ and a simple minimal eigenvalue $\lambda_2^{\varphi_1}$, and*

$$\lambda_1^{\varphi_1} > 1, \quad \lambda_2^{\varphi_1} < 1.$$




FIGURE 3.3. The Jordan normal form of the Coxeter transformation

EXAMPLE 3.15 (V. Kolmykov). The example shows that there is a graph Γ with indefinite and degenerate quadratic form $\mathcal{B}$ such that $\dim \ker \mathbf{B}$ is however large (see Fig. 3.4) and the Coxeter transformation has however large number of 2×2 Jordan blocks. Consider n instances of the extended Dynkin diagram $\tilde{D}_4$ with centers $b_1, \dots, b_n$ and marked vertices $a_1, \dots, a_n$. We add new vertex b_{n+1} and connect it with marked vertices a_i for $i = 1, 2, \dots, n$. The new graph is bipartite: one part consists of the vertices $b_1, \dots, b_n, b_{n+1}$; the matrix D^tD is of size $(n+1) \times (n+1)$ and has the form




FIGURE 3.4. A graph Γ such that $\dim \ker \mathbf{B}$ is however large

$$4D^t D = \begin{pmatrix} n & 1 & 1 & 1 & \dots & 1 & 1 \\ 1 & 4 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 4 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & 4 & \dots & 0 & 0 \\ & & & & \dots & & \\ 1 & 0 & 0 & 0 & \dots & 4 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 4 \end{pmatrix}.$$

It is easy to show that

$$|4D^t D - \mu I| = (n - \mu)(4 - \mu)^n - n(4 - \mu)^{n-1}.$$

Thus, $\varphi_i = \frac{\mu_i}{4} = 1$ is of multiplicity $n - 1$.

In Proposition 4.10 of §4.3 we will see more examples of graphs $\Gamma(n)$ obtained by gluing n copies of some graph Γ . By this proposition every eigenvalue λ of Γ is also an eigenvalue of $\Gamma(n)$ and the multiplicity of this eigenvalue for the graph $\Gamma(n)$ is $(n - 1) \times m$, where m is the multiplicity of this eigenvalue in Γ .

CHAPTER 4

Eigenvalues, shearing formulas and diagrams $T_{p,q,r}$

... most of the fundamental results about simple Lie algebras, which were discovered by Killing are usually attributed to É. Cartan. This, despite the meticulousness with which Cartan noted his indebtedness to Killing. In Cartan's thesis there are 28 references to Lie and 60 to Killing!

A. J. Coleman, [Col89, p.447], 1989.

4.1. The eigenvalues of the affine Coxeter transformation are roots of unity

The Coxeter transformation corresponding to the extended Dynkin diagram, i.e., corresponding to affine Kac-Moody algebra is called the *affine Coxeter transformation*.

THEOREM 4.1 ([SuSt79], [St82a], [St85]). *The eigenvalues of the affine Coxeter transformation are roots of unity. The proper eigenvalues are collected in Table 4.1.*

Proof. The eigenvalues for all cases of extended Dynkin diagrams are easily calculated by means of the generalized R. Steinberg's theorem (Theorem 5.5) and Table 1.2. See Remark 4.2. $\square$

REMARK 4.2. According to generalized R. Steinberg's theorem (Theorem 5.5) orders (not always different) of eigenvalues (column 3, Table 4.1) coincide with lengths of branches of the corresponding Dynkin diagram. If g is the class number (5.33) of the extended Dynkin diagram, the number of branches of the corresponding Dynkin diagram is $3 - g$. For $g = 0$, we have a simply-laced case of the extended Dynkin diagrams, i.e., $\tilde{E}_6$, $\tilde{E}_7$, $\tilde{E}_8$, $\tilde{D}_n (n \geq 4)$. In this case number of branches is 3. For $g = 1$, we have extended Dynkin diagrams $\tilde{F}_{41}$, $\tilde{F}_{42}$, $\tilde{CD}_n$, $\widetilde{CD}_n$, there exists 2 groups of eigenvalues, see (5.34). For $g = 2$, we have extended Dynkin diagrams $\tilde{G}_{21}$, $\tilde{G}_{22}$, $\tilde{B}_n$, $\tilde{C}_n$, $\widetilde{BC}_n$ and there exists only one group of eigenvalues. For $g = 3$, we have extended Dynkin diagrams $\tilde{A}_{11}$, $\tilde{A}_{12}$, in this case there is only one trivial eigenvalues 1.

TABLE 4.1. The eigenvalues of affine Coxeter transformations

Diagram	Eigenvalues λ	Orders of eigenvalues ¹	Affine Coxeter number h_a ²
$\tilde{E}_6$	$\lambda_{1,2}^1 = 1,$ $\lambda_{1,2}^2 = \lambda_{1,2}^3 = e^{\pm 2\pi i/3},$ $\lambda_7 = -1$	3, 3, 2	6
$\tilde{E}_7$	$\lambda_{1,2}^1 = 1,$ $\lambda_{1,2}^2 = e^{\pm 2\pi i/3},$ $\lambda_{1,2}^3 = e^{\pm \pi i/2},$ $\lambda_{7,8} = -1$	3, 4, 2	12
$\tilde{E}_8$	$\lambda_{1,2}^1 = 1,$ $\lambda_{1,2}^2 = e^{\pm 2\pi i/3},$ $\lambda_{1,2}^3 = e^{\pm 2\pi i/5}, \lambda_{1,2}^4 = e^{\pm 4\pi i/5},$ $\lambda_9 = -1$	3, 5, 2	30
$\tilde{D}_n$	$\lambda_{1,2}^1 = 1,$ $\lambda_s^2 = e^{2s\pi i/(n-2)}, s = 1, 2, \dots, n-3,$ $\lambda_n = \lambda_{n+1} = -1$	$n-2,$ 2, 2	$n-2$ for $n = 2k$; $2(n-2)$ for $n = 2k+1$
$\tilde{G}_{21}, \tilde{G}_{22}$	$\lambda_{1,2}^1 = 1,$ $\lambda_3 = -1$	2	2
$\tilde{F}_{41}, \tilde{F}_{42}$	$\lambda_{1,2}^1 = 1,$ $\lambda_{1,2}^2 = e^{2\pi i/3},$ $\lambda_3 = -1$	3, 2	6
$\tilde{A}_{11}, \tilde{A}_{12}$	$\lambda_{1,2}^1 = 1$	1	1
$\tilde{B}_n, \tilde{C}_n,$ $\widetilde{BC}_n$	$\lambda_{1,2}^1 = 1,$ $\lambda_s^2 = e^{2s\pi i/n}, s = 1, 2, \dots, n-1$	n	n
$\widetilde{CD}_n,$ $\widetilde{DD}_n$	$\lambda_{1,2}^1 = 1,$ $\lambda_s^2 = e^{2s\pi i/(n-1)}, s = 1, 2, \dots, n-2$ $\lambda_{n+1} = -1$	$n-1,$ 2	$n-1$ for $n = 2k-1$; $2(n-1)$ for $n = 2k$

REMARK 4.3. Let H be the hyperplane orthogonal to the adjoint vector $\tilde{z}^1$ from Proposition 3.9. The vector $\tilde{z}^1$ is responsible for a 2×2 block in the Jordan form of the affine

¹See Remark 4.2.

²See Remark 4.3.

TABLE 4.2. The Coxeter numbers and affine Coxeter numbers

Extended Dynkin Diagram	Notation in context of twisted affine Lie algebra ¹	Affine Coxeter number h_a	Coxeter number h	Dual ² Coxeter number $h^\vee$
$\widetilde{E}_6$	$E_6^{(1)}$	6	12	12
$\widetilde{E}_7$	$E_7^{(1)}$	12	18	18
$\widetilde{E}_8$	$E_8^{(1)}$	30	30	30
$\widetilde{D}_n$	$D_n^{(1)}$	$n - 2$ for $n = 2k$; $2(n - 2)$ for $n = 2k + 1$	$2(n - 1)$	$2(n - 1)$
$\widetilde{A}_{11}$	$A_2^{(2)}$	1	3	3
$\widetilde{A}_{12}$	$A_1^{(1)}$	1	2	2
$\widetilde{BC}_n$	$A_{2n}^{(2)}$	n	$2n + 1$	$2n + 1$
$\widetilde{G}_{22} = \widetilde{G}_{21}^\vee$	$G_2^{(1)}$	2	6	$\boxed{4}$
$\widetilde{G}_{21} = \widetilde{G}_{22}^\vee$	$D_4^{(3)}$	2	$\boxed{4}$	6
$\widetilde{F}_{42} = \widetilde{F}_{41}^\vee$	$F_4^{(1)}$	6	12	$\boxed{9}$
$\widetilde{F}_{41} = \widetilde{F}_{42}^\vee$	$E_6^{(2)}$	6	$\boxed{9}$	12
$\widetilde{C}_n = \widetilde{B}_n^\vee$	$C_n^{(1)}$	n	$2n$	$\boxed{n+1}$
$\widetilde{B}_n = \widetilde{C}_n^\vee$	$D_{n+1}^{(2)}$	n	$\boxed{n+1}$	$2n$
$\widetilde{CD}_n = \widetilde{DD}_n^\vee$	$B_n^{(1)}$	$n - 1$ for $n = 2k - 1$; $2(n - 1)$ for $n = 2k$	$2n$	$\boxed{2n-1}$
$\widetilde{DD}_n = \widetilde{CD}_n^\vee$	$A_{2n-1}^{(2)}$	$n - 1$ for $n = 2k - 1$; $2(n - 1)$ for $n = 2k$	$\boxed{2n-1}$	$2n$

Coxeter transformation. Due to the presence of a 2×2 block, the affine Coxeter transformation is of infinite order in the Weyl group. The restriction of the Coxeter transformation on the hyperplane H is, however, of a finite order h_a . We call this number the *affine Coxeter number*. The affine Coxeter number h_a is the least common multiple of orders of eigenvalues ($\neq 1$) of the affine Coxeter transformation (Table 4.1). We denote by h the *Coxeter number* of the Dynkin diagram. The value $h - 1$ is the sum of coordinates of the highest root β of the corresponding root system, see (1.3). The imaginary vector z^1 depicted on Fig. 2.2 coincides with the highest root β extended to the vector with 1 at the additional vertex, the one that extends the Dynkin diagram to the extended Dynkin diagram. Thus,

$$h = n_1 + \cdots + n_l, \quad (4.1)$$

¹See, Remark 4.4.

²For emphasis and to distinguish h from $h^\vee$, we frame the value $\min(h, h^\vee)$ in the case $h \neq h^\vee$.

where n_i coordinates of the imaginary vector from Fig. 2.2. Similarly to (4.1), the *dual Coxeter number* is defined as

$$h^\vee = n_1^\vee + \cdots + n_l^\vee, \quad (4.2)$$

where $\{n_1^\vee, \dots, n_l^\vee\}$ are coordinates of the imaginary vector $h^\vee$ of the dual Dynkin diagram, Fig. 2.2. For values of the dual Coxeter numbers, see Table 4.2.

REMARK 4.4. For the first time, the notation of *twisted affine Lie algebras* from col. 2, Table 4.2 appeared in [Kac69] in the description of finite order automorphisms; see also [Kac80], [GorOnVi94, p.123], [OnVi90], [Kac93]. The upper index r in the notation of twisted affine Lie algebras has an invariant sense: is the order of the diagram automorphism μ of $\mathfrak{g}$, where $\mathfrak{g}$ is a complex simple finite dimensional Lie algebra of type $X_N = A_l, D_{l+1}, E_6, D_4$, [Kac93, Th.8.3].

The affine Lie algebra associated to a generalized Cartan matrix of type $X_l^{(1)}$ is called a *non-twisted affine Lie algebra*, [Kac93, Ch.7].

The affine Lie algebras associated to a generalized Cartan matrix of type $X_l^{(2)}$ and $X_l^{(3)}$ are called *twisted affine Lie algebras*. [Kac93, Ch.8].

The corresponding $\mathbb{Z}/r\mathbb{Z}$ -gradings are (here $\bar{i} \in \mathbb{Z}/r\mathbb{Z}$):

$$\begin{aligned} \mathfrak{g} &= \mathfrak{g}_{\bar{0}} + \mathfrak{g}_{\bar{1}}, \quad \text{for } r = 2, \\ \mathfrak{g} &= \mathfrak{g}_{\bar{0}} + \mathfrak{g}_{\bar{1}} + \mathfrak{g}_{\bar{2}}, \quad \text{for } r = 3, \end{aligned} \quad (4.3)$$

see [Kac93, 8.3.1, 8.3.2], [GorOnVi94, Ch.3, §3]

PROPOSITION 4.5. ([Kac93, exs.6.3]) *Let A be a Cartan matrix of type $X_N^{(r)}$ from col.2, Table 4.2, let $l = \text{rank of } A$, and let h be the Coxeter number. Let Δ be the finite root system of type X_N . Then*

$$rlh = |\Delta|. \quad (4.4)$$

Proof. For $r = 1$, we have H. S. M. Coxeter's proposition (1.1), §1.1. We consider only remaining cases:

$$A_2^{(2)}, \quad A_{2n}^{(2)} (l \geq 2), \quad A_{2n-1}^{(2)} (l \geq 3), \quad D_{n+1}^{(2)} (l \geq 2), \quad E_6^{(2)}, \quad D_4^{(3)}.$$

(see col. 5, Table 4.2).

(a) $A_2^{(2)}$, $r = 2$; rank $l = 1$; $h = 3$; $X_N = A_2$, $|\Delta(A_2)| = 6$, see [Bo, Tab.I].

(b) $A_{2n}^{(2)}$, $r = 2$; rank $l = n$; $h = 2n + 1$, $X_N = A_{2n}$, $|\Delta(A_{2n})| = 2n(2n + 1)$, see [Bo, Tab.I].

(c) $A_{2n-1}^{(2)}$, $r = 2$; rank $l = n$; $h = 2n - 1$; $X_N = A_{2n-1}$, $|\Delta(A_{2n-1})| = 2n(2n - 1)$, see [Bo, Tab.I].

(d) $D_{n+1}^{(2)}$, $r = 2$; rank $l = n$; $h = n + 1$, $X_N = D_{n+1}$, $|\Delta(D_{n+1})| = 2n(n + 1)$, see [Bo, Tab.IV].

(e) $E_6^{(2)}$, $r = 2$; rank $l = 4$; $h = 9$, $X_N = E_6$, $|\Delta(E_6)| = 72$, see [Bo, Tab.V].

(f) $D_4^{(3)}$, $r = 3$; rank $l = 2$; $h = 4$, $X_N = D_4$, $|\Delta(D_4)| = 24$, see [Bo, Tab.IV].

Cases (a)-(f) are collected in Table 4.3. □

4.2. Bibliographical notes on the spectrum of the Coxeter transformation

The eigenvalues of affine Coxeter transformations were also calculated by S. Berman, Y.S. Lee and R. Moody in [BLM89]. Theorem 3.14 was also proved by N. A'Campo [A'C76] and R. Howlett [How82].

TABLE 4.3. Proposition 4.5 for $r = 2, 3$.

Extended Dynkin Diagram	Index r	Rank of A	Coxeter number h	Root System X_N	$ \Delta = \Delta(X_N) $
$A_2^{(2)}$	2	1	3	A_2	6
$A_{2n}^{(2)}$	2	n	$2n + 1$	A_{2n}	$2n(2n + 1)$
$A_{2n-1}^{(2)}$	2	n	$2n - 1$	A_{2n-1}	$2n(2n - 1)$
$D_{n+1}^{(2)}$	2	n	$2n + 1$	D_{n+1}	$2n(2n + 1)$
$E_6^{(2)}$	2	4	9	E_6	72
$D_4^{(3)}$	3	2	4	D_4	24

Natural difficulties in the study of Cartan matrices and Coxeter transformations for the graphs containing cycles are connected with the following two facts:

- 1) these graphs have non-symmetrizable Cartan matrices,
- 2) in general, there are several conjugacy classes of the Coxeter transformation.

We would like to distinguish several works related to the Coxeter transformation for the graphs with cycles: C. M. Ringel [Rin94], A. J. Coleman [Col89], Shi Jian-yi [Shi00], Menshikh and Subbotin [MeSu82], [Men85], Boldt and Takane [BT97].

For generalized Cartan matrices (see §2.1), i.e., for graphs with cycles, C. M. Ringel [Rin94] showed that the *spectral radius* $\rho(\mathbf{C})$ of the Coxeter transformation lies out the unit circle, and is an eigenvalue of multiplicity one. This result generalizes Theorem 3.14, 3), proved in [SuSt75, SuSt78, St85] only for trees. The spectral radius $\rho(\mathbf{C})$ is used by V. Dlab and C. M. Ringel to determine the Gelfand-Kirillov dimension of the preprojective algebras, [DR81].

A. J. Coleman [Col89] computed characteristic polynomials for the Coxeter transformation for all extended Dynkin diagrams, including the case with cycles $\tilde{A}_n$. He baptized these polynomials *Killing polynomials*, (see the epigraph to this chapter). Coleman also shows that $\tilde{A}_n$ has $\frac{n}{2}$ spectral conjugacy classes.

V. V. Menshikh and V. F. Subbotin in [MeSu82], and V. V. Menshikh in [Men85] established a connection between an orientation Δ of the graph Γ and spectral classes of conjugacy of the Coxeter transformation. For any orientation Δ of a given graph Γ containing several cycles, they consider an invariant R_Δ equal to the number of arrows directed in a clockwise direction. For any graph Γ containing disjoint cycles, they show that $R_{\Delta_1} = R_{\Delta_2}$ if and only if orientations Δ_1 and Δ_2 can be obtained from each other by applying a sink-admissible or a source-admissible sequence of reflections σ_i , see §2.6, i.e.,

$$R_{\Delta_1} = R_{\Delta_2} \iff \Delta_1 = \sigma_{i_n} \dots \sigma_{i_2} \sigma_{i_1}(\Delta_2)$$

for any sink-admissible or source-admissible sequence $i_1, i_2, \dots, i_n$. Menshikh and Subbotin also showed that two Coxeter transformations $\mathbf{C}_{\Delta_1}$ and $\mathbf{C}_{\Delta_2}$ are conjugate if and only if $R_{\Delta_1} = R_{\Delta_2}$. The number R_Δ is called the *index* of the conjugacy class of the Coxeter transformation. Menshikh and Subbotin also calculated the characteristic polynomial of the

Coxeter transformation for every class equivalent to Δ for the extended Dynkin diagram $\tilde{A}_n$; this polynomial is

$$\det |\mathbf{C} - \lambda I| = \lambda^{n+1} - \lambda^{n-k+1} - \lambda^k + 1, \quad (4.5)$$

where $k = R_\Delta$ is the index of the conjugacy class of the Coxeter transformation.

Shi Jian-yi [Shi00] considers conjugacy relation on Coxeter transformations for the case where Γ is just a cycle. Different equivalence relations on Coxeter transformations are considered in more difficult cases. In [Shi00], Shi also obtained an explicit formula (4.5).

In [Bol96], [BT97], Boldt and Takane considered unicyclic graphs.

DEFINITION 4.6 ([Bol96]). An *essential cycle* of Γ is a full subgraph S of Γ having $m \geq 3$ vertices $x_1, \dots, x_m$ such that there are edges between x_i and x_{i+1} for $i = 1, \dots, m-1$ and also between x_m and x_1 (thus, Γ coincides with $\tilde{A}_n$). A graph Γ is called the *unicyclic* if it contains precisely one *essential cycle*.

Boldt and Takane showed how the characteristic polynomial of the Coxeter transformation for the unicyclic graph Γ can be reduced to the characteristic polynomial for the essential cycle. They also arrive at the explicit formula (4.5).

4.3. Shearing and gluing formulas for the characteristic polynomial

The purpose of this section is to prove the Subbotin-Sumin *splitting along the edge* formula for the characteristic polynomial of the Coxeter transformation [SuSum82], to prove its generalization for the multiply-laced case and to get some of its corollaries that will be used in the following sections.

Let us consider a characteristic polynomial of the Coxeter transformation

$$\mathcal{X}(\Gamma, \lambda) = \det |\Gamma - \lambda I|$$

for the graph Γ with a *splitting edge*. Recall that an edge l is said to be *splitting* if deleting it we split the graph Γ into two graphs Γ_1 and Γ_2 , see Fig. 4.1.




FIGURE 4.1. A split graph Γ

PROPOSITION 4.7 ([SuSum82]). For a given graph Γ with a *splitting edge* l , we have

$$\mathcal{X}(\Gamma, \lambda) = \mathcal{X}(\Gamma_1, \lambda) \mathcal{X}(\Gamma_2, \lambda) - \lambda \mathcal{X}(\Gamma_1 \setminus \alpha, \lambda) \mathcal{X}(\Gamma_2 \setminus \beta, \lambda), \quad (4.6)$$

where α and β are the endpoints of the deleted edge l .

Proof. We select a basis in $\mathcal{E}_\Gamma$ such that

$$\begin{aligned}\mathcal{E}_\Gamma &= \mathcal{E}_{\Gamma_1} + \mathcal{E}_{\Gamma_2}, \\ \mathcal{E}_{\Gamma_1} &= \{a_1, \dots, a_m\}, \quad a_m = \alpha, \quad \dim \mathcal{E}_{\Gamma_1} = m, \\ \mathcal{E}_{\Gamma_2} &= \{b_1, \dots, b_k\}, \quad b_1 = \beta, \quad \dim \mathcal{E}_{\Gamma_2} = k,\end{aligned}$$

and a Coxeter transformation $\mathbf{C}$ such that

$$\mathbf{C} = w_{\Gamma_1} w_{\Gamma_2}, \quad w_{\Gamma_1} = \sigma_{a_m} \dots \sigma_{a_1}, \quad w_{\Gamma_2} = \sigma_{b_1} \dots \sigma_{b_k}.$$

An ordering of reflections σ_{a_i} (resp. σ_{b_j}) in w_{Γ_1} (resp. w_{Γ_2}) is selected so that σ_{a_m} (resp. σ_{b_1}) is the last to be executed. Then the reflections σ_{a_i} ($i \neq m$) (resp. σ_{b_j} ($j \neq 1$)) do not act on the coordinates b_j (resp. a_i). Only σ_{a_m} acts on the coordinate b_1 and σ_{b_1} acts on the coordinate a_m . We have

$$\begin{aligned}w_{\Gamma_1} &= \sigma_{a_m} \dots \sigma_{a_1} = \begin{pmatrix} \mathbf{C}_{\Gamma_1} & \delta_{a_m b_1} \\ 0 & I \end{pmatrix} \begin{matrix} \} m \\ \} k \end{matrix}, \\ w_{\Gamma_2} &= \sigma_{b_1} \dots \sigma_{b_k} = \begin{pmatrix} I & 0 \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} \end{pmatrix} \begin{matrix} \} m \\ \} k \end{matrix},\end{aligned}\tag{4.7}$$

where $\mathbf{C}_{\Gamma_1}$ (resp. $\mathbf{C}_{\Gamma_2}$) is the Coxeter transformation for the graph Γ_1 (resp. Γ_2) and $\delta_{a_m b_1}$ (resp. $\delta_{b_1 a_m}$) means the matrix with zeros everywhere except 1 in the $(a_m b_1)$ th (resp. $(b_1 a_m)$ th) slot.

Then we have

$$\begin{aligned}\mathbf{C} &= w_{\Gamma_1} w_{\Gamma_2} = \begin{pmatrix} \mathbf{C}_{\Gamma_1} + \delta_{a_m a_m} & \delta_{a_m b_1} \mathbf{C}_{\Gamma_2} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} \end{pmatrix}, \\ \det |\mathbf{C} - \lambda I| &= \begin{vmatrix} \mathbf{C}_{\Gamma_1} + \delta_{a_m a_m} - \lambda I & \delta_{a_m b_1} \mathbf{C}_{\Gamma_2} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} - \lambda I \end{vmatrix}.\end{aligned}\tag{4.8}$$

Here, $\delta_{a_m b_1} \mathbf{C}_{\Gamma_2}$ is the b_1 st line (the first line in $\mathbf{C}_{\Gamma_2}$ for the selected basis) of the matrix $\mathbf{C}_{\Gamma_2}$ and we subtract now this line from the line a_m . Then

$$\det |\mathbf{C} - \lambda I| = \begin{vmatrix} \mathbf{C}_{\Gamma_1} - \lambda I & \lambda \delta_{a_m b_1} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} - \lambda I \end{vmatrix}$$

or

$$\det |\mathbf{C} - \lambda I| = \begin{vmatrix} & & 0 & 0 & \dots & 0 \\ & & 0 & 0 & \dots & 0 \\ & \mathbf{C}_{\Gamma_1} - \lambda I & & & \dots & \\ & & 0 & 0 & \dots & 0 \\ & & \lambda & 0 & \dots & 0 \\ 0 & \dots & 0 & 1 & & \\ 0 & \dots & 0 & 0 & & \\ & \dots & & & \mathbf{C}_{\Gamma_2} - \lambda I & \\ 0 & \dots & 0 & 0 & & \\ 0 & \dots & 0 & 0 & & \end{vmatrix}.\tag{4.9}$$

Let us expand the determinant $\det |\mathbf{C} - \lambda I|$ in (4.9) with respect to the minors corresponding to the line b_1 (the line containing λ in the column b_1 , which is the next after a_m). We see that

$$\det |\mathbf{C} - \lambda I| = \det |\mathbf{C}_{\Gamma_1} - \lambda I| \det |\mathbf{C}_{\Gamma_2} - \lambda I| + (-1)^{a_m + (a_m + 1)} \lambda R, \quad (4.10)$$

where R is the determinant shown on the Fig. 4.2. Expanding the determinant R we get the Subbotin-Sumin formula (4.6). $\square$




FIGURE 4.2. The asterisks mark the deleted line a_m and column b_1

Now we will extend the Subbotin-Sumin formula to the multiply-laced case where the endpoints α and β correspond to simple roots of different lengths. In this case, formula (4.6) is modified to (4.11).

PROPOSITION 4.8. *For a given graph Γ with a splitting weighted edge l corresponding to roots with different lengths, we have*

$$\mathcal{X}(\Gamma, \lambda) = \mathcal{X}(\Gamma_1, \lambda) \mathcal{X}(\Gamma_2, \lambda) - \rho \lambda \mathcal{X}(\Gamma_1 \setminus \alpha, \lambda) \mathcal{X}(\Gamma_2 \setminus \beta, \lambda), \quad (4.11)$$

where α and β are the endpoints of the deleted edge l and ρ is the ratio of lengths of the roots corresponding to these endpoints.

Proof. Relation (4.8) is modified to the following

$$\mathbf{C} = \begin{pmatrix} \mathbf{C}_{\Gamma_1} + \rho \delta_{a_m a_m} & \rho \delta_{a_m b_1} \mathbf{C}_{\Gamma_2} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} \end{pmatrix}.$$

Multiply the line b_1 by ρ and subtract this product from the line a_m , then we obtain

$$\det |\mathbf{C} - \lambda I| = \begin{vmatrix} \mathbf{C}_{\Gamma_1} + \rho \delta_{a_m a_m} - \lambda I & \rho \delta_{a_m b_1} \mathbf{C}_{\Gamma_2} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} - \lambda I \end{vmatrix} = \begin{vmatrix} \mathbf{C}_{\Gamma_1} - \lambda I & \rho \lambda \delta_{a_m b_1} \\ \delta_{b_1 a_m} & \mathbf{C}_{\Gamma_2} - \lambda I \end{vmatrix} \quad (4.12)$$

Further, as in Proposition 4.7, we obtain (4.11). $\square$

COROLLARY 4.9. *Let Γ_2 in Proposition 4.8 be a component containing only one point. Then, the following formula holds*

$$\mathcal{X}(\Gamma, \lambda) = -(\lambda + 1)\mathcal{X}(\Gamma_1, \lambda) - \rho\lambda\mathcal{X}(\Gamma_1 \setminus \alpha, \lambda), \quad (4.13)$$

Proof. Relation (4.7) is modified to the following

$$\begin{aligned} w_{\Gamma_1} &= \left(\begin{array}{cc} 0 & \\ 0 & \\ \mathbf{C}_{\Gamma_1} & \cdots \\ & 0 \\ & \rho \\ 0 & 1 \end{array} \right) \begin{array}{l} \\ \\ \\ \\ \\ \end{array} \left. \vphantom{\begin{array}{c} \\ \\ \\ \\ \\ \end{array}} \right\}^m, \\ w_{\Gamma_2} &= \left(\begin{array}{cc} I & 0 \\ 0 & 0 \dots 1 & -1 \end{array} \right) \begin{array}{l} \\ \\ \end{array} \left. \vphantom{\begin{array}{c} \\ \\ \end{array}} \right\}^m, \end{aligned} \quad (4.14)$$

and (4.12) is modified to

$$\begin{aligned} \mathbf{C} &= \left(\begin{array}{cc} 0 & \\ 0 & \\ \mathbf{C}_{\Gamma_1} + \rho\delta_{a_m a_m} & \cdots \\ & 0 \\ & -\rho \\ 0 & 0 \dots 1 & -1 \end{array} \right), \\ \det |\mathbf{C} - \lambda I| &= \left| \begin{array}{cc} 0 & \\ 0 & \\ \mathbf{C}_{\Gamma_1} + \rho\delta_{a_m a_m} - \lambda I & \cdots \\ & 0 \\ & -\rho \\ 0 & 0 \dots 1 & -1 - \lambda \end{array} \right| = \left| \begin{array}{cc} 0 & \\ 0 & \\ \mathbf{C}_{\Gamma_1} - \lambda I & \cdots \\ & 0 \\ & \rho\lambda \\ 0 & 0 \dots 1 & -1 - \lambda \end{array} \right| \end{aligned} \quad (4.15)$$

Further, as in Proposition 4.7, we obtain (4.13). $\square$

The next proposition follows from formula (4.6) and allows one to calculate the spectrum of the graph $\Gamma(n)$ obtained by gluing n copies of the graph Γ .

PROPOSITION 4.10 ([SuSum82], [KMSS83], [KMSS83a]). *Let $*_n$ be a star with n rays coming from a vertex. Let $\Gamma(n)$ be the graph obtained from $*_n$ by gluing to the endpoints of its rays n copies of the graph Γ . Then*

$$\mathcal{X}(\Gamma(n), \lambda) = \mathcal{X}(\Gamma, \lambda)^{n-1} \varphi_{n-1}(\lambda) \quad (4.16)$$

for some polynomial $\varphi_{n-1}(\lambda)$. The polynomials $\varphi_{n-1}(\lambda)$ are recursively calculated.

Proof. By splitting along the edge l (4.6) we get, for $n = 2$, (Fig. 4.3)

$$\begin{aligned}
\mathcal{X}(\Gamma(2), \lambda) &= \\
\mathcal{X}(\Gamma, \lambda)\mathcal{X}(\Gamma + \beta, \lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)\mathcal{X}(\Gamma, \lambda) &= \\
\mathcal{X}(\Gamma, \lambda)[\mathcal{X}(\Gamma + \beta, \lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)] &= \\
\mathcal{X}(\Gamma, \lambda)\varphi_1(\lambda), \text{ where } \varphi_1(\lambda) &= \\
\mathcal{X}(\Gamma + \beta, \lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda). &
\end{aligned}$$




FIGURE 4.3. Splitting along the edge l

Let the proposition be true for $n = r$ and

$$\mathcal{X}(\Gamma(r), \lambda) = \mathcal{X}(\Gamma, \lambda)^{r-1}\varphi_{r-1}(\lambda).$$

Then, for $n = r + 1$, we have

$$\begin{aligned}
\mathcal{X}(\Gamma(r+1), \lambda) &= \mathcal{X}(\Gamma, \lambda)\mathcal{X}(\Gamma(r), \lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)(\mathcal{X}(\Gamma, \lambda))^r = \\
&= \mathcal{X}(\Gamma, \lambda)\mathcal{X}(\Gamma, \lambda)^{r-1}\varphi_{r-1}(\lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)\mathcal{X}(\Gamma, \lambda)^r = \\
&= \mathcal{X}(\Gamma, \lambda)^r[\varphi_{r-1}(\lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)]. \quad \square
\end{aligned}$$

The following proposition is due to V. Kolmykov. For the case $\lambda = 1$, it is formulated in [Kac93, Ch.4, exs.4.12].

PROPOSITION 4.11. *If the spectrum of the Coxeter transformations for graphs Γ_1 and Γ_2 contains the eigenvalue λ , then this eigenvalue is also the eigenvalue of the graph Γ obtained by gluing as described in Proposition 4.10.*


Proof. By splitting along the edge l (4.6) we get (Fig. 4.4

$$\begin{aligned}
\mathcal{X}(\Gamma_1 + \beta + \Gamma_2, \lambda) &= \\
\mathcal{X}(\Gamma_1, \lambda)\mathcal{X}(\Gamma_2 + \beta, \lambda) - \lambda\mathcal{X}(\Gamma \setminus \alpha, \lambda)\mathcal{X}(\Gamma_2, \lambda). &
\end{aligned}$$

If $\mathcal{X}(\Gamma_1, \lambda)|_{\lambda=\lambda_0} = 0$ and $\mathcal{X}(\Gamma_2, \lambda)|_{\lambda=\lambda_0} = 0$, then also $\mathcal{X}(\Gamma, \lambda)|_{\lambda=\lambda_0} = 0$. $\square$

Let $\mathcal{X}(A_n)$ be the characteristic polynomial of the Coxeter transformation for the Dynkin diagram A_n . Then, from the recurrent formula (4.13), we have

$$\begin{aligned}
\mathcal{X}(A_1) &= -(\lambda + 1), \\
\mathcal{X}(A_2) &= \lambda^2 + \lambda + 1, \\
\mathcal{X}(A_3) &= -(\lambda^3 + \lambda^2 + \lambda + 1), \\
\mathcal{X}(A_4) &= \lambda^4 + \lambda^3 + \lambda^2 + \lambda + 1, \\
&\dots \\
\mathcal{X}(A_n) &= -(\lambda + 1)\mathcal{X}(A_{n-1}) - \lambda\mathcal{X}(A_{n-2}), \quad n > 2.
\end{aligned} \tag{4.17}$$

FIGURE 4.4. Splitting along the edge l , $\Gamma_1 \neq \Gamma_2$

REMARK 4.12. Since the characteristic polynomial can be chosen up to a factor, we prefer to consider the polynomial

$$\chi_n = (-1)^n \mathcal{X}(A_n) \quad (4.18)$$

as the characteristic polynomial of the Coxeter transformation of A_n , thus the leading coefficient is positive. Pay attention that the sign $(-1)^n$ for diagrams with n vertices should be taken into account only in recurrent calculations using formula (4.13) (cf. with calculation for $\tilde{D}_n$ and $\tilde{D}_4$ below).

REMARK 4.13. 1) J. S. Frame in [Fr51, p.784] obtained that

$$\chi(A_{m+n}) = \chi(A_m)\chi(A_n) - \lambda\chi(A_{m-1})\chi(A_{n-1}), \quad (4.19)$$

which easily follows from (4.6). Another *Frame formula* will be proved in Proposition 5.2.

2) S. M. Gussein-Zade in [Gu76] obtained the formula

$$\mathcal{X}(A_n) = -(\lambda + 1)\chi_{n-1} - \lambda\chi_{n-2}, \quad (4.20)$$

for the characteristic polynomial of the classical monodromy, see (4.17).

From (4.17) for $n > 0$, we see that χ_n is the cyclotomic polynomial whose roots are the primitive n th roots of unity:

$$\chi_n = \sum_{i=1}^n \lambda^i = \frac{\lambda^{n+1} - 1}{\lambda - 1}. \quad (4.21)$$

4.4. Formulas of the characteristic polynomials for the diagrams $T_{p,q,r}$

We give here explicit formulas of characteristic polynomials of the Coxeter transformations for the three classes of diagrams: $T_{2,3,r}$, $T_{3,3,r}$, and $T_{2,4,r}$, see §2.5, Fig. 2.3.

4.4.1. The diagrams $T_{2,3,r}$. The case $T_{2,3,r}$, where $r \geq 2$, includes diagrams D_5 , E_6 , E_7 , E_8 , $\tilde{E}_8$, E_{10} . Since $D_5 = T_{2,3,2} = E_5$ and $\tilde{E}_8 = T_{2,3,6} = E_9$, we call these diagrams the E_n -series, where $n = r + 3$. In Table 1.1 and Table 1.2 we see that characteristic polynomials of E_6 , E_7 , E_8 , $\tilde{E}_8$ constitute a series, and this series can be continued, see (4.22) and Table 4.4.

REMARK 4.14. C. McMullen observed in [McM02] that the spectral radius $\rho(\mathbf{C})$ of the Coxeter transformation for all graphs with indefinite Tits form attains its minimum when the diagram is $T_{2,3,7} = E_{10}$, see Fig. 4.5, [Hir02], [McM02], and §2.5 about *hyperbolic Dynkin diagrams*. McMullen showed that $\rho(\mathbf{C})$ is the so-called *Lehmer's number*,

$$\lambda_{Lehmer} \approx 1.176281\dots, \text{ see Table 4.4.}$$

For details and definitions, see §B.2.

Lehmer's number is a root of the polynomial

$$1 + x - x^3 - x^4 - x^5 - x^6 - x^7 + x^9 + x^{10},$$

see [Hir02], [McM02].




FIGURE 4.5. The spectral radius attains its minimum on the graph E_{10}

PROPOSITION 4.15. *The characteristic polynomials of Coxeter transformations for the diagrams $T_{2,3,n}$ are as follows:*

$$\chi(T_{2,3,n-3}) = \chi(E_n) = \lambda^n + \lambda^{n-1} - \sum_{i=3}^{n-3} \lambda^i + \lambda + 1, \quad (4.22)$$

where $\chi(E_n)$ is given up to sign $(-1)^n$, see Remark 4.12. In other words,

$$\chi(T_{2,3,n-3}) = \chi(E_n) = \frac{\lambda^{n-2}(\lambda^3 - \lambda - 1) + (\lambda^3 + \lambda^2 - 1)}{\lambda - 1}. \quad (4.23)$$

The spectral radius $\rho(T_{2,3,n-3})$ converges to the maximal root ρ_{max} of the equation

$$\lambda^3 - \lambda - 1 = 0, \quad (4.24)$$

and

$$\rho_{max} = \sqrt[3]{\frac{1}{2} + \sqrt{\frac{23}{108}}} + \sqrt[3]{\frac{1}{2} - \sqrt{\frac{23}{108}}} \approx 1.324717... \quad (4.25)$$

REMARK 4.16. Note that the spectral radius (4.25) was obtained by Y. Zhang [Zh89] and used in the study of regular components of an Auslander-Reiten quiver. The spectral radius (4.25) is also the smallest *Pisot number*, see §B.2.

Proof. Use induction and (4.13). We have

$$\begin{aligned} (-1)^{n+2}\chi(E_{n+2}) &= -(\lambda + 1)(-1)^{n+1}\chi(E_{n+1}) - \lambda(-1)^n\chi(E_n) = \\ &= (-1)^{n+2}[(\lambda + 1)\chi(E_{n+1}) - \lambda\chi(E_n)]. \end{aligned} \quad (4.26)$$

Thus,

$$\chi(E_{n+2}) = (\lambda + 1)\chi(E_{n+1}) - \lambda\chi(E_n) = \lambda(\chi(E_{n+1}) - \chi(E_n)) + \chi(E_{n+1}).$$

By the induction hypothesis, (4.22) yields

$$\lambda(\chi(E_{n+1}) - \chi(E_n)) = \lambda^{n+2} - \lambda^n - \lambda^{n-1},$$

and

$$\begin{aligned}\chi(E_{n+2}) &= \lambda^{n+2} - \lambda^n - \lambda^{n-1} + (\lambda^{n+1} + \lambda^n - \sum_{i=3}^{n-2} \lambda^i + \lambda + 1) = \\ &= \lambda^{n+2} + \lambda^{n+1} - \sum_{i=3}^{n-1} \lambda^i + \lambda + 1,\end{aligned}$$

and hence (4.22) is proved. Further,

$$\sum_{i=3}^{n-3} \lambda^i = \lambda^3 \sum_{i=0}^{n-6} \lambda^i = \lambda^3 \frac{\lambda^{n-5} - 1}{\lambda - 1} = \frac{\lambda^{n-2} - \lambda^3}{\lambda - 1},$$

and

$$\begin{aligned}\chi(E_n) &= \lambda^n + \lambda^{n-1} - \frac{\lambda^{n-2} - \lambda^3}{\lambda - 1} + \lambda + 1 = \\ &= \frac{\lambda^{n+1} - \lambda^{n-1} - \lambda^{n-2} + \lambda^3 + \lambda^2 - 1}{\lambda - 1} = \\ &= \frac{\lambda^{n-2}(\lambda^3 - \lambda - 1) + (\lambda^3 + \lambda^2 - 1)}{\lambda - 1}.\end{aligned}$$

This proves (4.23). For $\lambda > 1$, the characteristic polynomial $\chi(E_n)$ converges to

$$\frac{\lambda^{n-2}(\lambda^3 - \lambda - 1)}{\lambda - 1}.$$

By Proposition 4.21 the spectral radius ρ_{max} converges to the maximal root of equation (4.24). For the results of calculations related to the Proposition 4.15, see Table 4.4. $\square$

4.4.2. The diagrams $T_{3,3,r}$. The case $T_{3,3,r}$, where $r \geq 2$, includes diagrams E_6 and $\tilde{E}_6$, and so we call these diagrams the $E_{6,n}$ -series, where $n = r - 2$. Thus, $E_6 = T_{3,3,2} = E_{6,0}$ and $\tilde{E}_6 = T_{3,3,3} = E_{6,1}$. In Table 1.1 and Table 1.2 we see that the characteristic polynomials of $E_6, \tilde{E}_6$ constitute a series, and this series can be continued, see (4.27) and Table 4.5.

PROPOSITION 4.17. *The characteristic polynomials of Coxeter transformations for the diagrams $T_{3,3,n}$ with $n \geq 3$ are calculated as follows:*

$$\chi(T_{3,3,n}) = \chi(E_{6,n-2}) = \lambda^{n+4} + \lambda^{n+3} - 2\lambda^{n+1} - 3 \sum_{i=4}^n \lambda^i - 2\lambda^3 + \lambda + 1, \quad (4.27)$$

where $\chi(E_{6,n})$ is given up to sign $(-1)^n$, see Remark 4.12. In other words,

$$\begin{aligned}\chi(T_{3,3,n}) &= \chi(E_{6,n-2}) = \\ &= \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - 2\lambda - 1) + (\lambda^4 + 2\lambda^3 + \lambda^2 - 1)}{\lambda - 1} = \\ &= \frac{(\lambda^2 + \lambda + 1)[\lambda^{n+1}(\lambda^2 - \lambda - 1) + (\lambda^2 + \lambda - 1)]}{\lambda - 1}.\end{aligned} \quad (4.28)$$

The spectral radius $\rho(T_{3,3,n})$ converges to the maximal root ρ_{max} of the equation

$$\lambda^2 - \lambda - 1 = 0, \quad (4.29)$$

and

$$\rho_{max} = \frac{\sqrt{5} + 1}{2} \approx 1.618034... \quad (4.30)$$

TABLE 4.4. The characteristic polynomials for $T_{2,3,r}$

$T_{2,3,r}$ and its well-known name	Characteristic polynomial	Maximal eigenvalue outside the unit circle
$(2, 3, 2)$ D_5	$\lambda^5 + \lambda^4 + \lambda + 1$	—
$(2, 3, 3)$ E_6	$\lambda^6 + \lambda^5 - \lambda^3 + \lambda + 1$	—
$(2, 3, 4)$ E_7	$\lambda^7 + \lambda^6 - \lambda^4 - \lambda^3 + \lambda + 1$	—
$(2, 3, 5)$ E_8	$\lambda^8 + \lambda^7 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	—
$(2, 3, 6)$ $\tilde{E}_8$	$\lambda^9 + \lambda^8 - \lambda^6 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	—
$(2, 3, 7)$ E_{10}	$\lambda^{10} + \lambda^9 - \lambda^7 - \lambda^6 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	1.176281...
$(2, 3, 8)$ E_{11}	$\lambda^{11} + \lambda^{10} - \lambda^8 - \lambda^7 - \lambda^6 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	1.230391...
$(2, 3, 9)$ E_{12}	$\lambda^{12} + \lambda^{11} - \lambda^9 - \lambda^8 - \lambda^7 - \lambda^6 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	1.261231...
$(2, 3, 10)$ E_{13}	$\lambda^{13} + \lambda^{12} - \lambda^{10} - \lambda^9 - \lambda^8 - \lambda^7$ $- \lambda^6 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1$	1.280638...
...	...	...
$(2, 3, n)$ E_n	$\frac{\lambda^{n-2}(\lambda^3 - \lambda - 1) + (\lambda^3 + \lambda^2 - 1)}{\lambda - 1}$	1.324717... as $n \rightarrow \infty$

REMARK 4.18. For $n = 3$, the sum $\sum_{i=4}^n \lambda^i$ in (4.27) disappears and we have

$$\begin{aligned}\chi(T_{3,3,3}) &= \chi(E_{6,1}) = \chi(\tilde{E}_6) = \\ \lambda^7 + \lambda^6 - 2\lambda^4 - 2\lambda^3 + \lambda + 1 &= (\lambda^3 - 1)^2(\lambda + 1).\end{aligned}\tag{4.31}$$

Proof. As above in (4.22), we use induction and (4.13). So, by (4.13) we have

$$\begin{aligned}\chi(E_{6,n+2}) &= (\lambda + 1)\chi(E_{6,n+1}) - \lambda\chi(E_{6,n}) = \\ \lambda(\chi(E_{6,n+1}) - \chi(E_{6,n})) &+ \chi(E_{6,n+1}).\end{aligned}\tag{4.32}$$

By the induction hypothesis and (4.27) we have

$$\lambda(\chi(E_{6,n+1}) - \chi(E_{6,n})) = \lambda^{n+8} - \lambda^{n+6} - 2\lambda^{n+5} - \lambda^{n+4},\tag{4.33}$$

and

$$\begin{aligned}\chi(E_{6,n+2}) &= \\ \lambda^{n+8} - \lambda^{n+6} - 2\lambda^{n+5} - \lambda^{n+4} &+ \\ (\lambda^{n+7} + \lambda^{n+6} - 2\lambda^{n+4} - 3\sum_{i=4}^{n+3} \lambda^i - 2\lambda^3 + \lambda + 1) &= \\ \lambda^{n+8} + \lambda^{n+7} - 2\lambda^{n+5} - 3\sum_{i=4}^{n+4} \lambda^i - 2\lambda^3 + \lambda + 1;\end{aligned}\tag{4.34}$$

this proves (4.27). Further,

$$\begin{aligned}\chi(T_{3,3,n}) &= \lambda^{n+4} + \lambda^{n+3} - 2\lambda^{n+1} - 3\sum_{i=4}^n \lambda^i - 2\lambda^3 + \lambda + 1 = \\ \lambda^{n+4} + \lambda^{n+3} + \lambda^{n+1} - 3\sum_{i=3}^{n+1} \lambda^i + \lambda^3 + \lambda + 1 &= \\ \lambda^{n+1}(\lambda^3 + \lambda^2 + 1) - 3\lambda^3 \frac{\lambda^{n-1} - 1}{\lambda - 1} + \lambda^3 + \lambda + 1,\end{aligned}$$

i.e.,

$$\begin{aligned}\chi(T_{3,3,n}) &= \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 + \lambda - 1) - 3\lambda^{n+2} + 3\lambda^3 + (\lambda^4 - \lambda^3 + \lambda^2 - 1)}{\lambda - 1} = \\ \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - 2\lambda - 1) + (\lambda^4 + 2\lambda^3 + \lambda^2 - 1)}{\lambda - 1},\end{aligned}$$

and (4.28) is proved. For $\lambda > 1$, the characteristic polynomial $\chi(T_{3,3,n})$ converges to

$$\frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - 2\lambda - 1)}{\lambda - 1} = \frac{\lambda^{n+1}(\lambda^2 - \lambda - 1)(\lambda^2 + \lambda + 1)}{\lambda - 1}.$$

By Proposition 4.21 the spectral radius ρ_{max} converges to the maximal root of the equation (4.29). For the results of calculations related to the Proposition 4.17, see Table 4.5. $\square$

TABLE 4.5. The characteristic polynomials for $T_{3,3,r}$

$T_{3,3,r}$ and its well-known name	Characteristic polynomial	Maximal eigenvalue outside the unit circle
$(3, 3, 2)$ E_6	$\lambda^6 + \lambda^5 - \lambda^3 + \lambda + 1$	—
$(3, 3, 3)$ $\tilde{E}_6$	$\lambda^7 + \lambda^6 - 2\lambda^4 - 2\lambda^3 + \lambda + 1$	—
$(3, 3, 4)$ $E_{6,2}$	$\lambda^8 + \lambda^7 - 2\lambda^5 - 3\lambda^4 - 2\lambda^3 + \lambda + 1$	1.401268...
$(3, 3, 5)$ $E_{6,3}$	$\lambda^9 + \lambda^8 - 2\lambda^6 - 3\lambda^5 - 3\lambda^4 - 2\lambda^3 + \lambda + 1$	1.506136...
$(3, 3, 6)$ $E_{6,4}$	$\lambda^{10} + \lambda^9 - 2\lambda^7 - 3\lambda^6 - 3\lambda^5 - 3\lambda^4 - 2\lambda^3 + \lambda + 1$	1.556030...
$(3, 3, 7)$ $E_{6,5}$	$\lambda^{11} + \lambda^{10} - 2\lambda^8 - 3\lambda^7 - 3\lambda^6 - 3\lambda^5 - 3\lambda^4 - 2\lambda^3 + \lambda + 1$	1.582347...
$(3, 3, 8)$ $E_{6,6}$	$\lambda^{12} + \lambda^{11} - 2\lambda^9 - 3\lambda^8 - 3\lambda^7$ $- 3\lambda^6 - 3\lambda^5 - 3\lambda^4 - 2\lambda^3 + \lambda + 1$	1.597005...
...	...	...
$(3, 3, n)$ $E_{6,n-2}$	$\frac{(\lambda^2 + \lambda + 1)[\lambda^{n+1}(\lambda^2 - \lambda - 1) + (\lambda^2 + \lambda - 1)]}{\lambda - 1}$	1.618034... as $n \rightarrow \infty$

4.4.3. The diagrams $T_{2,4,r}$. The case $T_{2,4,r}$, where $r \geq 2$, includes diagrams $D_6, E_7, \tilde{E}_7$ and we call these diagrams the $E_{7,n}$ -series, where $n = r - 3$. Thus, $D_6 = T_{2,4,2} = E_{7,-1}$, $E_7 = T_{2,4,3} = E_{7,0}$, $\tilde{E}_7 = T_{2,4,4} = E_{7,1}$. In Table 1.1 and Table 1.2 we see that characteristic polynomials of $E_7, \tilde{E}_7$ constitute a series, and this series can be continued, see (4.35) and Table 4.6.

PROPOSITION 4.19. *The characteristic polynomials of Coxeter transformations for diagrams $T_{2,4,n}$, where $n \geq 3$, are calculated as follows:*

$$\chi(T_{2,4,n}) = \chi(E_{7,n-3}) = \lambda^{n+4} + \lambda^{n+3} - \lambda^{n+1} - 2 \sum_{i=4}^n \lambda^i - \lambda^3 + \lambda + 1, \quad (4.35)$$

where $\chi(E_{7,n})$ is given up to sign $(-1)^n$, see Remark 4.12. In other words

$$\chi(T_{2,4,n}) = \chi(E_{7,n-3}) = \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - \lambda - 1) + (\lambda^4 + \lambda^3 + \lambda - 1)}{\lambda - 1} = \frac{(\lambda + 1)(\lambda^{n+1}(\lambda^3 - \lambda^2 - 1) + (\lambda^3 + \lambda - 1))}{\lambda - 1}. \quad (4.36)$$

The spectral radius $\rho(T_{2,4,n})$ converges to the maximal root ρ_{max} of the equation

$$\lambda^3 - \lambda^2 - 1 = 0, \quad (4.37)$$

and

$$\rho_{max} = \frac{1}{3} + \sqrt[3]{\frac{58}{108} + \sqrt{\frac{31}{108}}} + \sqrt[3]{\frac{58}{108} - \sqrt{\frac{31}{108}}} \approx 1.465571\dots \quad (4.38)$$

REMARK 4.20. For $n = 3$, the sum $\sum_{i=4}^n \lambda^i$ disappears from (4.35) and we have

$$\begin{aligned} \chi(T_{2,4,3}) &= \chi(E_{7,0}) = \chi(E_7) = \\ \lambda^7 + \lambda^6 - \lambda^4 - \lambda^3 + \lambda + 1 &= (\lambda^4 - 1)(\lambda^3 - 1)(\lambda + 1). \end{aligned} \quad (4.39)$$

Proof. Use induction and (4.13). As above in (4.32), we have

$$\begin{aligned} \chi(E_{7,n+2}) &= \\ (\lambda + 1)\chi(E_{7,n+1}) - \lambda\chi(E_{7,n}) &= \\ \lambda(\chi(E_{7,n+1}) - \chi(E_{7,n})) + \chi(E_{7,n+1}). \end{aligned} \quad (4.40)$$

By the induction hypothesis and (4.35) we have

$$\lambda(\chi(E_{7,n+1}) - \chi(E_{7,n})) = \lambda^{n+9} - \lambda^{n+7} - \lambda^{n+6} - \lambda^{n+5}, \quad (4.41)$$

and we see that

$$\begin{aligned} \chi(E_{7,n+2}) &= \\ \lambda^{n+9} - \lambda^{n+7} - \lambda^{n+6} - \lambda^{n+5} &+ (\lambda^{n+8} + \lambda^{n+7} - \lambda^{n+5} - 2 \sum_{i=4}^{n+4} \lambda^i - \lambda^3 + \lambda + 1) = \\ \lambda^{n+9} + \lambda^{n+8} - \lambda^{n+6} - 2 \sum_{i=4}^{n+5} \lambda^i - \lambda^3 + \lambda + 1; \end{aligned} \quad (4.42)$$

this proves (4.35). Further,

$$\begin{aligned}
\chi(T_{2,4,n}) &= \lambda^{n+4} + \lambda^{n+3} - \lambda^{n+1} - 2 \sum_{i=4}^n \lambda^i - \lambda^3 + \lambda + 1 = \\
&= (\lambda^{n+4} + \lambda^{n+3} + \lambda^{n+1}) - 2 \sum_{i=3}^{n+1} \lambda^i + (\lambda^3 + \lambda + 1) = \\
&= \lambda^{n+1}(\lambda^3 + \lambda^2 + 1) - 2 \frac{\lambda^3(\lambda^{n-1} - 1)}{\lambda - 1} + (\lambda^3 + \lambda + 1) = \\
&= \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 + \lambda - 1) - 2\lambda^{n+2} + 2\lambda^3 + (\lambda^4 - \lambda^3 + \lambda^2 - 1)}{\lambda - 1} = \\
&= \frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - \lambda - 1) + (\lambda^4 + \lambda^3 + \lambda^2 - 1)}{\lambda - 1} = \\
&= \frac{\lambda^{n+1}(\lambda + 1)(\lambda^3 - \lambda^2 - 1) + (\lambda + 1)(\lambda^3 + \lambda - 1)}{\lambda - 1},
\end{aligned} \tag{4.43}$$

and (4.36) is proved. For $\lambda > 1$, the characteristic polynomial $\chi(T_{2,4,n})$ converges to

$$\frac{\lambda^{n+1}(\lambda + 1)(\lambda^3 - \lambda^2 - 1)}{\lambda - 1}.$$

By Proposition 4.21, the spectral radius ρ_{max} converges to the maximal root of the equation (4.37). For the results of calculations related to the Proposition 4.19, see Table 4.6. $\square$

The maximal eigenvalues (Tables 4.4, 4.5, 4.6) lying out the unit circle are obtained by means of the online service “Polynomial Roots Solver” of “EngineersToolbox”¹.

4.4.4. Convergence of the sequence of eigenvalues. In this section we give a substantiation (Proposition 4.21) of the fact that the maximal eigenvalues (i.e., spectral radiuses) of the characteristic polynomials for $T_{p,q,r}$ converges to a real number which is the maximal value of a known polynomial (see, Propositions 4.15, 4.17, 4.19). We will give this substantiation in a generic form. Similarly, we have a dual fact (Proposition 4.24): the minimal eigenvalues of the characteristic polynomials for $T_{p,q,r}$ converges to a real number which is the minimal value of a known polynomial.

PROPOSITION 4.21. *Let $f(\lambda)$ and $g(\lambda)$ be some polynomials in λ , and let*

$$P_n(\lambda) = \lambda^n f(\lambda) + g(\lambda) \tag{4.44}$$

be a sequence of polynomials in λ . Let $z_n \in \mathbb{R}$ be a root of $P_n(\lambda)$, lying in the interval $(1, 2)$:

$$P_n(z_n) = 0, \quad z_n \in (1, 2), \quad n = 1, 2, \dots \tag{4.45}$$

Suppose that the sequence $\{z_n\}, n = 1, 2, \dots$ is non-decreasing:

$$1 < z_1 \leq z_2 \leq \dots \leq z_n \leq \dots \tag{4.46}$$

Then the sequence z_n converges to a real number z_0 which is a root of $f(\lambda)$:

$$f(z_0) = 0. \tag{4.47}$$

¹See <http://www.engineerstoobox.com>.

TABLE 4.6. The characteristic polynomials for $T_{2,4,r}$

$T_{2,4,r}$ and its well-known name	Characteristic polynomial	Maximal eigenvalue outside the unit circle
$(2, 4, 2)$ D_6	$\lambda^6 + \lambda^5 + \lambda + 1$	—
$(2, 4, 3)$ E_7	$\lambda^7 + \lambda^6 - \lambda^4 - \lambda^3 + \lambda + 1$	—
$(2, 4, 4)$ $\tilde{E}_7$	$\lambda^8 + \lambda^7 - \lambda^5 - 2\lambda^4 - \lambda^3 + \lambda + 1$	—
$(2, 4, 5)$ $E_{7,2}$	$\lambda^9 + \lambda^8 - \lambda^6 - 2\lambda^5 - 2\lambda^4 - \lambda^3 + \lambda + 1$	1.280638...
$(2, 4, 6)$ $E_{7,3}$	$\lambda^{10} + \lambda^9 - \lambda^7 - 2\lambda^6 - 2\lambda^5 - 2\lambda^4 - \lambda^3 + \lambda + 1$	1.360000...
$(2, 4, 7)$ $E_{7,4}$	$\lambda^{11} + \lambda^{10} - \lambda^8 - 2\lambda^7 - 2\lambda^6 - 2\lambda^5 - 2\lambda^4 - \lambda^3 + \lambda + 1$	1.401268...
$(2, 4, 8)$ $E_{7,5}$	$\lambda^{12} + \lambda^{11} - \lambda^9 - 2\lambda^8 - 2\lambda^7$ $- 2\lambda^6 - 2\lambda^5 - 2\lambda^4 - \lambda^3 + \lambda + 1$	1.425005...
...	...	...
$(2, 4, n)$ E_n	$\frac{\lambda^{n+1}(\lambda^4 - \lambda^2 - \lambda - 1) + (\lambda^4 + \lambda^3 + \lambda - 1)}{\lambda - 1}$	1.465571... as $n \rightarrow \infty$

Proof. According to (4.45), the sequence (4.46) is non-decreasing and bounded from above. Therefore, there exists a real number z_0 such that

$$\lim_{k \rightarrow \infty} z_k = z_0. \quad (4.48)$$

TABLE 4.7. The diagrams $T_{p,q,r}$ and characteristic polynomials $\lambda^n f(\lambda) + g(\lambda)$

Diagram	$f(\lambda)$	$g(\lambda)$	$D(\lambda)$	$P_n(\lambda)$
$T(2, 3, r)$	$\lambda^3 - \lambda - 1$	$\lambda^3 + \lambda^2 - 1$	$\frac{1}{\lambda - 1}$	$\lambda^{n-2} f(\lambda) + g(\lambda)$
$T(3, 3, r)$	$\lambda^2 - \lambda - 1$	$\lambda^2 + \lambda - 1$	$\frac{\lambda^2 + \lambda + 1}{\lambda - 1}$	$\lambda^{n+1} f(\lambda) + g(\lambda)$
$T(2, 4, r)$	$\lambda^3 - \lambda^2 - 1$	$\lambda^3 + \lambda - 1$	$\frac{\lambda + 1}{\lambda - 1}$	$\lambda^{n+1} f(\lambda) + g(\lambda)$

Let us estimate $|f(z_0)|$ as follows:

$$|f(z_0)| < |f(z_n)| + |f(z_n) - f(z_0)|. \quad (4.49)$$

By (4.48) we obtain

$$|f(z_0)| < |f(z_n)| + \varepsilon_n, \quad \text{where } \varepsilon_n = |f(z_n) - f(z_0)| \rightarrow 0. \quad (4.50)$$

By (4.45), we have

$$f(z_n) = -\frac{g(z_n)}{z_n^n}. \quad (4.51)$$

Let

$$\delta_n = \left| \frac{g(z_n)}{z_n^n} \right| \quad \text{for each } n = 1, 2, \dots \quad (4.52)$$

Since the function $g(z)$ is uniformly bounded on the interval $(1, 2)$, it follows that $\delta_n \rightarrow 0$. By (4.50) we have

$$|f(z_0)| < \delta_n + \varepsilon_n \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (4.53)$$

Thus, $f(z_0) = 0$. $\square$

COROLLARY 4.22. *Proposition 4.21 holds also for the following non-polynomial functions P_n :*

1) For

$$P_n(\lambda) = \lambda^{n+k} f(\lambda) + g(\lambda), \quad (4.54)$$

where $k \in \mathbb{Z}$ is independent of n .

2) For

$$P_n(\lambda) = D(\lambda)(\lambda^n f(\lambda) + g(\lambda)), \quad (4.55)$$

where $D(\lambda)$ is a rational function independent of n and without roots on the interval $(1, 2)$.

COROLLARY 4.23. *The maximal values of characteristic polynomials of the Coxeter transformation for diagrams $T_{p,q,r}$ satisfy the conditions of Proposition 4.21, see Table 4.7.*

Proof. Let $\varphi_{1,r} = \varphi_1(T_{p,q,r})$ be the dominant value of the matrix DD^t (see §3.3.2) for the diagram $T_{p,q,r}$. According to Proposition 3.11 the dominant value $\varphi_{1,r}$ may only grow, i.e.,

$$\varphi_{1,r} \leq \varphi_{1,r+1}.$$

Therefore, the corresponding eigenvalue λ_1^{max} also only grows. Indeed, by (3.17) we get

$$\begin{aligned} \lambda_{1,r}^{max} &= 2\varphi_{1,r} - 1 + 2\sqrt{\varphi_{1,r}(\varphi_{1,r} - 1)} \leq \\ &2\varphi_{1,r+1} - 1 + 2\sqrt{\varphi_{1,r+1}(\varphi_{1,r+1} - 1)} = \lambda_{1,r+1}^{max}. \end{aligned} \quad (4.56)$$

In addition, from Proposition 3.11 we deduce that $\varphi_{1,r} > 1$, therefore the corresponding eigenvalue $\lambda_{1,r}$ of the Coxeter transformation is also > 1 .

It remains to show that every $\lambda_{1,r} < 2$. This is clear, because $f(\lambda) > 0$ and $g(\lambda) > 0$ for every $\lambda > 2$, see Table 4.7. $\square$

PROPOSITION 4.24. *Let $f(\lambda)$ and $g(\lambda)$ be some polynomials in λ , and let*

$$P_n(\lambda) = \lambda^n f(\lambda) + g(\lambda), \quad \text{where } n \in \mathbb{Z}_+. \quad (4.57)$$

Let $z_n \in \mathbb{R}$ be a root of $P_n(\lambda)$, lying in the interval $\left(\frac{1}{2}, 1\right)$:

$$P_n(z_n) = 0, \quad z_n \in \left(\frac{1}{2}, 1\right). \quad (4.58)$$

Suppose, that the sequence $\{z_n\}_{n \in \mathbb{Z}_+}$ is non-increasing:

$$1 > z_1 \geq z_2 \geq \dots \geq z_n \geq \dots \quad (4.59)$$

Then the sequence z_n converges to the some real number z_0 which is a root of $g(\lambda)$:

$$g(z_0) = 0. \quad (4.60)$$

Proof. As in Proposition 4.21, there is a real number z_0 such that $z_n \rightarrow z_0$ as $n \rightarrow \infty$. Since $g(z_n) \rightarrow g(z_0)$, we obtain

$$|g(z_0)| < |g(z_n)| + \varepsilon_n, \quad \text{where } \varepsilon_n = |g(z_n) - g(z_0)|.$$

By (4.58), we have

$$g(z_n) = -z_n^n f(z_n).$$

Let

$$\delta_n = |z_n^n f(z_n)| \quad \text{for each } n = 1, 2, \dots$$

Since the function $f(z_n)$ is uniformly bounded on the interval $\left(\frac{1}{2}, 1\right)$, it follows that $\delta_n \rightarrow 0$.

Thus, we have

$$|g(z_0)| < \delta_n + \varepsilon_n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus, $g(z_0) = 0$. $\square$

COROLLARY 4.25. *The minimal values of characteristic polynomials of the Coxeter transformation for diagrams $T_{p,q,r}$ satisfy the conditions of Proposition 4.21, see Table 4.7.*

Proof. The minimal value λ_1^{min} and the maximal value λ_1^{max} are reciprocal:

$$\lambda_1^{min} = \frac{1}{\lambda_1^{max}}.$$

Therefore, by (4.56), we see that the sequence of eigenvalues $\lambda_{1,r}^{min}$ is non-increasing:

$$\lambda_{1,r}^{min} \geq \lambda_{1,r+1}^{min}.$$

Since the maximal eigenvalue $\lambda_{1,r}^{max} > 1$, then $\lambda_{1,r}^{min} < 1$.

It remains to show that $\lambda_{1,r}^{min} > \frac{1}{2}$ for every $r \in \mathbb{Z}$. But this is true since $f(\lambda) < 0$ and $g(\lambda) < 0$ for every $\lambda < \frac{1}{2}$, see Table 4.7. $\square$

CHAPTER 5

R. Steinberg's theorem, B. Kostant's construction

5.1. R. Steinberg's theorem and a (p, q, r) mystery

R. Steinberg in [Stb85, p.591, (*)] observed a property of affine Coxeter transformations (i.e., transformations corresponding to extended Dynkin diagrams), which plays the main role in his derivation of the McKay correspondence. Let (p, q, r) be the same as in Table A.1, Table A.2 and relations (A.2), (A.6).

THEOREM 5.1 ([Stb85]). *The affine Coxeter transformation for the extended Dynkin diagram $\tilde{\Gamma}$ has the same eigenvalues as the product of three Coxeter transformations of types A_n , where $n = p - 1$, $q - 1$, and $r - 1$, corresponding to the branches of the Dynkin diagram Γ .*

Essentially, R. Steinberg observed that the orders of eigenvalues of the affine Coxeter transformation corresponding to the *extended* Dynkin diagram $\tilde{\Gamma}$ and given in Table 4.1 coincide with the lengths of branches of the Dynkin diagram Γ .

Now, we give the proof of R. Steinberg's theorem for the simply-laced extended Dynkin diagram by using the Subbotin-Sumin splitting formula (4.6).




FIGURE 5.1. Splitting the graph $T_{p,q,r}$ along the edge $\{\alpha\beta\}$

In §5.3.2, we generalize this theorem with some modifications to the multiply-laced case, see Theorem 5.5, Table 1.2.

Let $T_{p,q,r}$ be a connected graph with three branches of lengths p, q, r . Let χ be the characteristic polynomial of the Coxeter transformation. Split the graph along the edge $\{\alpha\beta\}$, Fig. 5.1. Then, by the Subbotin-Sumin formula (4.6) we have

$$\begin{aligned} \chi &= \mathcal{X}(A_{p+q-1})\mathcal{X}(A_{r-1}) - \lambda\mathcal{X}(A_{p-1})\mathcal{X}(A_{q-1})\mathcal{X}(A_{r-2}) = \\ &= (-1)^{p+q+r}(\chi_{p+q-1}\chi_{r-1} - \lambda\chi_{p-1}\chi_{q-1}\chi_{r-2}). \end{aligned} \tag{5.1}$$

By (4.21), we have, up to a factor $(-1)^{p+q+r}$,

$$\chi = \frac{(\lambda^{p+q} - 1)(\lambda^r - 1)}{(\lambda - 1)^2} - \frac{(\lambda^p - 1)(\lambda^q - 1)(\lambda^r - \lambda)}{(\lambda - 1)^3}. \tag{5.2}$$

For $p = q$ (e.g., E_6 , $\tilde{E}_6$, $\tilde{E}_7$, and D_n for $n \geq 4$), we have, up to a factor $(-1)^{2p+r}$,

$$\begin{aligned} \chi &= \frac{(\lambda^{2p} - 1)(\lambda^r - 1)}{(\lambda - 1)^2} - \frac{(\lambda^p - 1)^2(\lambda^r - \lambda)}{(\lambda - 1)^3} = \\ &= \frac{(\lambda^p - 1)}{(\lambda - 1)^3} ((\lambda^p + 1)(\lambda^r - 1)(\lambda - 1) - (\lambda^p - 1)(\lambda^r - \lambda)). \end{aligned} \quad (5.3)$$

For $q = 2p$ (e.g., E_7 , $\tilde{E}_8$), we have, up to a factor $(-1)^{3p+r}$,

$$\begin{aligned} \chi &= \frac{(\lambda^{3p} - 1)(\lambda^r - 1)}{(\lambda - 1)^2} - \frac{(\lambda^{2p} - 1)(\lambda^p - 1)(\lambda^r - \lambda)}{(\lambda - 1)^3} = \\ &= \frac{(\lambda^p - 1)}{(\lambda - 1)^3} ((\lambda^{2p} + \lambda^p + 1)(\lambda^r - 1)(\lambda - 1) - (\lambda^{2p} - 1)(\lambda^r - \lambda)). \end{aligned} \quad (5.4)$$

1) Case $p = q = 3$, $r = 3$ ($\tilde{E}_6$). From (5.3) we have

$$\begin{aligned} \chi &= \frac{(\lambda^3 - 1)}{(\lambda - 1)^3} [(\lambda^3 + 1)(\lambda^3 - 1)(\lambda - 1) - (\lambda^3 - 1)(\lambda^3 - \lambda)] = \\ &= \frac{(\lambda^3 - 1)^2}{(\lambda - 1)^3} [(\lambda^3 + 1)(\lambda - 1) - (\lambda^3 - \lambda)] = \\ &= \frac{(\lambda^3 - 1)^2}{(\lambda - 1)^3} (\lambda^4 - 2\lambda^3 + 2\lambda - 1) = \\ &= \frac{(\lambda^3 - 1)^2(\lambda - 1)^2(\lambda^2 - 1)}{(\lambda - 1)^3} = (\lambda - 1)^2 \chi_2^2 \chi_1. \end{aligned} \quad (5.5)$$

Polynomials χ_1 and χ_2^2 have, respectively, eigenvalues of orders 2, 3, 3 which are equal to the lengths of branches of E_6 .

2) Case $p = q = 4$, $r = 2$ ($\tilde{E}_7$). Here, from (5.3) we have

$$\begin{aligned} \chi &= \frac{(\lambda^4 - 1)}{(\lambda - 1)^3} [(\lambda^4 + 1)(\lambda^2 - 1)(\lambda - 1) - (\lambda^4 - 1)(\lambda^2 - \lambda)] = \\ &= \frac{(\lambda^4 - 1)(\lambda^2 - 1)(\lambda - 1)}{(\lambda - 1)^3} [(\lambda^4 + 1) - \lambda(\lambda^2 + 1)] = \\ &= \frac{(\lambda^4 - 1)(\lambda^3 - 1)(\lambda^2 - 1)(\lambda - 1)^2}{(\lambda - 1)^3} = (\lambda - 1)^2 \chi_3 \chi_2 \chi_1. \end{aligned} \quad (5.6)$$

Polynomials χ_1 , χ_2 and χ_3 have, respectively, eigenvalues of orders 2, 3, 4 which are equal to the lengths of branches of E_7 .

3) Case $p = 3$, $q = 6$, $r = 2$ ($\tilde{E}_8$). From (5.4), we have

$$\begin{aligned} \chi &= \frac{(\lambda^3 - 1)}{(\lambda - 1)^3} [(\lambda^6 + \lambda^3 + 1)(\lambda^2 - 1)(\lambda - 1) - (\lambda^6 - 1)(\lambda^2 - \lambda)] = \\ &= \frac{(\lambda^3 - 1)(\lambda^2 - 1)(\lambda - 1)}{(\lambda - 1)^3} [(\lambda^6 + \lambda^3 + 1) - \lambda(\lambda^4 + \lambda^2 + 1)] = \\ &= \frac{(\lambda^5 - 1)(\lambda^3 - 1)(\lambda^2 - 1)(\lambda - 1)^2}{(\lambda - 1)^3} = (\lambda - 1)^2 \chi_4 \chi_2 \chi_1. \end{aligned} \quad (5.7)$$

Polynomials χ_1 , χ_2 and χ_4 have, respectively, eigenvalues of orders 2, 3, 5 which are equal to the lengths of branches of E_8 .

4) Case $p = q = 2$, $r = 2, 3, \dots$ (D_{r+2}). From (5.3) we have

$$\begin{aligned}
 (-1)^r \mathcal{X}(D_{r+2}) &= \\
 \frac{(\lambda^2 - 1)}{(\lambda - 1)^3} [(\lambda^2 + 1)(\lambda^r - 1)(\lambda - 1) - (\lambda^2 - 1)(\lambda^r - \lambda)] &= \\
 \frac{(\lambda^2 - 1)(\lambda - 1)}{(\lambda - 1)^3} [(\lambda^2 + 1)(\lambda^r - 1) - (\lambda + 1)(\lambda^r - \lambda)] &= \\
 \frac{(\lambda^2 - 1)(\lambda - 1)^2(\lambda^{r+1} + 1)}{(\lambda - 1)^3} &= (\lambda + 1)(\lambda^{r+1} + 1),
 \end{aligned} \tag{5.8}$$

and, up to a sign,

$$\chi = (\lambda + 1)(\lambda^{r+1} + 1). \tag{5.9}$$

Pay attention that, as in (5.8), the sign $(-1)^r$ should be taken into account only in recurrent calculations (e.g., for $\tilde{D}_n$ and $\tilde{D}_4$ below), see Remark 4.12.

5) Case $\tilde{D}_{r+2}$, contains $r + 3$ points.




FIGURE 5.2. Diagrams $\tilde{D}_{r+2}$ and D_{r+2}

Here, from (4.6) we have

$$\mathcal{X}(\tilde{D}_{r+2}) = \mathcal{X}(D_r)\mathcal{X}(A_3) - \lambda\mathcal{X}(D_{r-1})\mathcal{X}(A_1)^2. \tag{5.10}$$

From (5.10) and (5.8) we have

$$\begin{aligned}
 (-1)^{r-1} \mathcal{X}(\tilde{D}_{r+2}) &= \\
 (\lambda + 1)(\lambda^{r-1} + 1)(\lambda^3 + \lambda^2 + \lambda + 1) - \lambda(\lambda + 1)(\lambda^{r-2} + 1)(\lambda + 1)^2 &= \\
 (\lambda + 1)^2 [(\lambda^{r-1} + 1)(\lambda^2 + 1) - (\lambda + 1)(\lambda^{r-1} + \lambda)] &= \\
 (\lambda + 1)^2 [(\lambda^{r+1} + \lambda^2 + \lambda^{r-1} + 1) - (\lambda^r + \lambda^{r-1} + \lambda^2 + \lambda)] &= \\
 (\lambda + 1)^2 [(\lambda^{r+1} - \lambda^r) - (\lambda - 1)] &= (\lambda + 1)^2(\lambda^r - 1)(\lambda - 1) = \\
 (\lambda - 1)^2 \chi_1^2 \chi_{r-1},
 \end{aligned}$$

and so

$$\chi = (\lambda - 1)^2 \chi_1^2 \chi_{r-1}. \tag{5.11}$$

Polynomials χ_1^2 and χ_{r-1} have, respectively, eigenvalues of orders 2, 2, r which are equal to the lengths of branches of D_{r+2} .

6) Case $\tilde{D}_4$. Here, by (5.8) we have

$$\begin{aligned}
 \mathcal{X}(\tilde{D}_4) &= \mathcal{X}(D_4)\mathcal{X}(A_1) - \lambda\mathcal{X}(A_1)^3 = -(\lambda + 1)^2(\lambda^3 + 1) + \lambda(\lambda + 1)^3 = \\
 &= -(\lambda + 1)^2(\lambda^3 + 1 - \lambda^2 - \lambda) = -(\lambda + 1)^2(\lambda^2 - 1)(\lambda - 1) = \\
 &= -(\lambda - 1)^2(\lambda + 1)^3,
 \end{aligned}$$

and so

$$\chi = (\lambda - 1)^2 \chi_1^3. \tag{5.12}$$

Polynomial χ_1^3 have eigenvalues of orders 2, 2, 2 which are equal to the lengths of branches of D_4 .

In (4.19), we saw that the simple Frame formula is a particular case of the Subbotin-Sumin formula (4.6). Now we will show another *Frame formula* from [Fr51, p.785] used by H. S. M. Coxeter in [Cox51].

PROPOSITION 5.2 ([Fr51]). *Let $T_{p,q,r}$ be a connected graph with three branches of lengths p, q, r , where the lengths include the branch point, see Fig. 2.3. Then*

$$\chi(T_{p,q,r}) = \chi(A_{p+q+r-2}) - \lambda^2 \chi(A_{p-2}) \chi(A_{q-2}) \chi(A_{r-2}). \quad (5.13)$$

Proof. By (4.6) we have

$$\chi(T_{p,q,r}) = \chi(A_{p+q-1}) \chi(A_{r-1}) - \lambda \chi(A_{p-1}) \chi(A_{q-1}) \chi(A_{r-2}). \quad (5.14)$$

This is equivalent to

$$\begin{aligned} \chi(T_{p,q,r}) = & \chi(A_{p+q-1}) \chi(A_{r-1}) - \lambda \chi(A_{p+q-2}) \chi(A_{r-2}) \\ & + \lambda \chi(A_{p+q-2}) \chi(A_{r-2}) - \lambda \chi(A_{p-1}) \chi(A_{q-1}) \chi(A_{r-2}). \end{aligned} \quad (5.15)$$

By the first Frame formula (4.19) we have

$$\chi(A_{p+q+r-2}) = \chi(A_{p+q-1}) \chi(A_{r-1}) - \lambda \chi(A_{p+q-2}) \chi(A_{r-2}), \quad (5.16)$$

and

$$\begin{aligned} \chi(T_{p,q,r}) = & \chi(A_{p+q+r-2}) + \lambda \chi(A_{p+q-2}) \chi(A_{r-2}) - \lambda \chi(A_{p-1}) \chi(A_{q-1}) \chi(A_{r-2}) = \\ & \chi(A_{p+q+r-2}) + \lambda \chi(A_{r-2}) [\chi(A_{p+q-2}) - \chi(A_{p-1}) \chi(A_{q-1})]. \end{aligned} \quad (5.17)$$

Again, by (4.19) we have

$$\chi(A_{p+q-2}) - \chi(A_{p-1}) \chi(A_{q-1}) = -\lambda \chi(A_{p-2}) \chi(A_{q-2}), \quad (5.18)$$

and from (5.17) we deduce

$$\chi(T_{p,q,r}) = \chi(A_{p+q+r-2}) - \lambda^2 \chi(A_{r-2}) \chi(A_{p-2}) \chi(A_{q-2}). \quad \square \quad (5.19)$$

5.2. Characteristic polynomials for the Dynkin diagrams

In order to calculate characteristic polynomials of the Coxeter transformations for the Dynkin diagrams, we use the Subbotin-Sumin formula (4.6), the generalized Subbotin-Sumin formula (4.11), its specialization (5.2) for arbitrary (p, q, r) -trees, and particular cases of (5.2): formula (5.3) for $p = q$, and (5.4) for $p = 2q$.

The results of calculations of this section concerning Dynkin diagrams are collected in Table 1.1.

1) Case $p = q = 3, r = 2$ (E_6). From (5.3) we have

$$\begin{aligned} \chi = & \frac{(\lambda^3 - 1)}{(\lambda - 1)^3} [(\lambda^3 + 1)(\lambda^2 - 1)(\lambda - 1) - (\lambda^3 - 1)(\lambda^2 - \lambda)] = \\ & \frac{(\lambda^3 - 1)}{(\lambda - 1)^3} (\lambda - 1)^2 [(\lambda^3 + 1)(\lambda + 1) - \lambda(\lambda^2 + \lambda + 1)] = \\ & \frac{(\lambda^3 - 1)}{(\lambda - 1)} (\lambda^4 - \lambda^2 + 1) = \\ & (\lambda^2 + \lambda + 1)(\lambda^4 - \lambda^2 + 1) = \lambda^6 + \lambda^5 - \lambda^3 + \lambda + 1. \end{aligned} \quad (5.20)$$

In another form, we have

$$\chi = \frac{(\lambda^3 - 1)}{(\lambda - 1)}(\lambda^4 - \lambda^2 + 1) = \frac{(\lambda^6 + 1)}{(\lambda^2 + 1)} \frac{(\lambda^3 - 1)}{(\lambda - 1)}. \quad (5.21)$$

2) Case $p = 2, q = 4, r = 3$ (E_7). From (5.4) we have, up to a sign,

$$\begin{aligned} \chi &= \frac{(\lambda^2 - 1)}{(\lambda - 1)^3} [(\lambda^4 + \lambda^2 + 1)(\lambda^3 - 1)(\lambda - 1) - (\lambda^4 - 1)(\lambda^3 - \lambda)] = \\ &= \frac{(\lambda^2 - 1)(\lambda - 1)^2}{(\lambda - 1)^3} [(\lambda^4 + \lambda^2 + 1)(\lambda^2 + \lambda + 1) - \\ &\quad (\lambda^3 + \lambda^2 + \lambda + 1)\lambda(\lambda + 1)] = \\ &= (\lambda + 1)[(\lambda^4 + \lambda^2 + 1)(\lambda^2 + \lambda + 1) - (\lambda^3 + \lambda^2 + \lambda + 1)(\lambda^2 + \lambda)] = \\ &= (\lambda + 1)(\lambda^6 - \lambda^3 + 1) = \lambda^7 + \lambda^6 - \lambda^4 - \lambda^3 + \lambda + 1. \end{aligned} \quad (5.22)$$

In another form, we have

$$\chi = -(\lambda + 1)(\lambda^6 - \lambda^3 + 1) = -\frac{(\lambda + 1)(\lambda^9 + 1)}{(\lambda^3 + 1)}. \quad (5.23)$$

3) Case $p = 3, q = 2, r = 5$ (E_8). From (5.2) we have

$$\begin{aligned} \chi &= \frac{(\lambda^5 - 1)^2}{(\lambda - 1)^2} - \frac{(\lambda^3 - 1)(\lambda^2 - 1)(\lambda^5 - \lambda)}{(\lambda - 1)^3} = \\ &= (\lambda^4 + \lambda^3 + \lambda^2 + \lambda + 1)^2 - \\ &\quad (\lambda^2 + \lambda + 1)(\lambda + 1)\lambda(\lambda^3 + \lambda^2 + \lambda + 1) = \\ &= \lambda^8 + \lambda^7 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1. \end{aligned} \quad (5.24)$$

Since

$$(\lambda^8 + \lambda^7 - \lambda^5 - \lambda^4 - \lambda^3 + \lambda + 1)(\lambda^2 - \lambda + 1) = \lambda^{10} - \lambda^5 + 1,$$

we have

$$\chi = \frac{\lambda^{10} - \lambda^5 + 1}{\lambda^2 - \lambda + 1} = \frac{(\lambda^{15} + 1)(\lambda + 1)}{(\lambda^5 + 1)(\lambda^3 + 1)}. \quad (5.25)$$

4) For the case D_n , see (5.8),

$$\begin{aligned} \mathcal{X}(D_n) &= (-1)^n(\lambda + 1)(\lambda^{n-1} + 1), \\ \chi &= (\lambda + 1)(\lambda^{n-1} + 1). \end{aligned} \quad (5.26)$$

5) Case F_4 . We use formula (4.11), splitting the diagram F_4 along the weighted edge into two diagrams $\Gamma_1 = A_2$ and $\Gamma_2 = A_2$. Here, $\rho = 2$.

$$\chi = \chi_2^2 - 2\lambda\chi_1^2 = (\lambda^2 + \lambda + 1)^2 - 2\lambda(\lambda + 1)^2 = \lambda^4 - \lambda^2 + 1.$$

In another form, we have

$$\chi = \lambda^4 - \lambda^2 + 1 = \frac{\lambda^6 + 1}{\lambda^2 + 1}. \quad (5.27)$$

6) Case G_2 . A direct calculation of the Coxeter transformation gives

$$\mathbf{C} = \begin{pmatrix} -1 & 3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 2 & -3 \\ 1 & -1 \end{pmatrix},$$

and

$$\chi = \lambda^2 - \lambda + 1 = \frac{\lambda^3 + 1}{\lambda + 1}. \quad (5.28)$$

7) The dual cases B_n and $C_n = B_n^\vee$. Since spectra of the Coxeter transformations of the dual graphs coincide, we can consider only the case B_n .

a) Consider B_2 . A direct calculation of the Coxeter transformation gives

$$\mathbf{C} = \begin{pmatrix} -1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ 1 & -1 \end{pmatrix},$$

and

$$\mathcal{X}(B_2) = \lambda^2 + 1. \quad (5.29)$$

b) Consider B_3 . We use formula (4.11), splitting the diagram B_3 along the weighted edge

$$\begin{aligned} \mathcal{X}(B_3) &= \mathcal{X}(A_2)\mathcal{X}(A_1) - 2\lambda\mathcal{X}(A_1) = -(\chi_2\chi_1 - 2\lambda\chi_1) = \\ &= -\chi_1(\chi_2 - 2\lambda) = -(\lambda + 1)(\lambda^2 + \lambda + 1 - 2\lambda) = \\ &= -(\lambda + 1)(\lambda^2 - \lambda + 1) = -(\lambda^3 + 1). \end{aligned} \quad (5.30)$$

c) Case B_n for $n \geq 4$.

$$\begin{aligned} \mathcal{X}(B_n) &= \\ \chi(A_{n-3})\chi(B_3) - \lambda\chi(A_{n-4})\chi(B_2) &= \chi_{n-3}\chi(B_3) - \lambda\chi_{n-4}\chi(B_2), \end{aligned}$$

i.e.,

$$\begin{aligned} (-1)^n \mathcal{X}(B_n) &= \\ \frac{\lambda^{n-2} - 1}{\lambda - 1}(\lambda^3 + 1) - \lambda \frac{\lambda^{n-3} - 1}{\lambda - 1}(\lambda^2 + 1) &= \\ \frac{(\lambda^{n-2} - 1)(\lambda^3 + 1) - (\lambda^{n-3} - 1)(\lambda^3 + \lambda)}{\lambda - 1} &= \\ \frac{\lambda^{n+1} - \lambda^n + \lambda - 1}{\lambda - 1} &= \lambda^n + 1, \end{aligned}$$

and

$$\mathcal{X}(B_n) = (-1)^n(\lambda^n + 1). \quad (5.31)$$

By (5.29), (5.30) and (5.31), we see, for the case B_n , that, up to a sign,

$$\chi = \lambda^n + 1. \quad (5.32)$$

5.3. A generalization of R. Steinberg's theorem

We will show now that R. Steinberg's theorem 5.1 can be extended to the multiply-laced case.

5.3.1. Folded Dynkin diagrams and branch points.

DEFINITION 5.3. A vertex is said to be a *branch point* of the Dynkin diagram in one of the following cases:

- (a) if it is the common endpoint of three edges (E_6, E_7, E_8, D_n);
- (b) if it is the common endpoint of two edges, one of which is non-weighted and one is weighted (B_n, C_n, F_4). Such a vertex is said to be a *non-homogeneous point*.
- (c) Let, by definition, both points of G_2 be branch points.

REMARK 5.4. Every multiply-laced Dynkin diagram (and also every extended Dynkin diagram) can be obtained by a so-called *folding* operation from a simply-laced diagrams, see Fig. 5.3, such that the branch point is transformed into the *non-homogeneous point*, see, e.g., I. Satake, [Sat60, p.109], J. Tits, [Ti66, p.39], P. Slodowy, [Sl80, Appendices I and III], or more recent works [FSS96], [Mohr04]. This fact was our motivation for considering *non-homogeneous points* from (b) as *branch points*.




FIGURE 5.3. Folding operation applied to Dynkin diagrams

Let us divide all extended Dynkin diagrams (trees) into 4 classes and let g be the number of the class:

Class $g = 0$ contains all simply-laced diagrams.

Class $g = 1$ contains multiply-laced diagrams which has only one weighted edge of type (1,2) or (2,1).

Class $g = 2$ contains multiply-laced diagrams with either two weighted edges or one weighted edge of type (1,3) or (3,1).

Class $g = 3$ contains multiply-laced diagrams with two vertices.

Thus,

$$\begin{aligned}
 g = 0 & \text{ for } \tilde{E}_6, \tilde{E}_7, \tilde{E}_8, \tilde{D}_n, \\
 g = 1 & \text{ for } \tilde{F}_{41}, \tilde{F}_{42}, \widetilde{CD}_n, \widetilde{DD}_n, \\
 g = 2 & \text{ for } \tilde{G}_{12}, \tilde{G}_{22}, \tilde{B}_n, \tilde{C}_n, \widetilde{BC}_n, \\
 g = 3 & \text{ for } \tilde{A}_{11}, \tilde{A}_{12}.
 \end{aligned} \tag{5.33}$$

5.3.2. R. Steinberg's theorem for the multiply-laced case. Now, we can generalize R. Steinberg's theorem for the multiply-laced case.

THEOREM 5.5. *The affine Coxeter transformation with the extended Dynkin diagram $\tilde{\Gamma}$ of class g (see (5.33)) has the same eigenvalues as the product of the Coxeter transformations of types A_i , where $i \in \{p-1, q-1, r-1\}$ and matches to $(3-g)$ branches of the Dynkin*

diagram Γ . In other words,

$$\begin{aligned}
 &\text{For } g = 0, \text{ the product } \chi_{p-1}\chi_{q-1}\chi_{r-1} \text{ is taken.} \\
 &\text{For } g = 1, \text{ the product } \chi_{p-1}\chi_{q-1} \text{ is taken.} \\
 &\text{For } g = 2, \text{ the product consists only of } \chi_{p-1}. \\
 &\text{For } g = 3, \text{ the product is trivial } (= 1).
 \end{aligned} \tag{5.34}$$

The remaining two eigenvalues of the affine Coxeter transformation are both equal to 1, see Table 1.2.

Proof. For $g = 0$, we have a simply-laced case considered in R. Steinberg's theorem (Theorem 5.1).

Let $g = 1$.

1) Cases $\tilde{F}_{41}$, $\tilde{F}_{42}$. Since the spectra of the characteristic polynomials of the Coxeter transformations of the dual graphs coincide, we consider $\tilde{F}_{41}$. We use formula (4.11), splitting the diagram $\tilde{F}_{41}$ along the weighted edge. We have

$$\begin{aligned}
 \mathcal{X}(\tilde{F}_{41}) &= -(\chi_3\chi_2 - 2\lambda\chi_2\chi_1) = -\chi_2(\chi_3 - 2\lambda\chi_1) = \\
 &= -\chi_2(\lambda^3 + \lambda^2 + \lambda + 1 - 2\lambda(\lambda + 1)) = \\
 &= -\chi_2(\lambda^3 - \lambda^2 - \lambda + 1) = \\
 &= -\chi_2(\lambda^2 - 1)(\lambda - 1) = -(\lambda - 1)^2\chi_2\chi_1,
 \end{aligned} \tag{5.35}$$

and, up to a sign, we have

$$\chi = (\lambda - 1)^2\chi_2\chi_1. \tag{5.36}$$

Polynomials χ_1 and χ_2 have, respectively, eigenvalues of orders 2 and 3 which are equal to the lengths of the branches A_1 and A_2 of F_4 without non-homogeneous branch point.

2) Cases $\widetilde{CD}_n$, $\widetilde{DD}_n$. These diagrams are obtained as extensions of B_n , see [Bo, Tab.II]. By (4.11) and by splitting the diagram $\widetilde{CD}_n$ along the weighted edge, we have

$$\mathcal{X}(\widetilde{CD}_n) = \mathcal{X}(D_n)\mathcal{X}(A_1) - 2\lambda\mathcal{X}(D_{n-1}).$$

By (5.26)

$$\mathcal{X}(D_n) = (-1)^n(\lambda + 1)(\lambda^{n-1} + 1).$$

Thus,

$$\begin{aligned}
 (-1)^{n-1}\mathcal{X}(\widetilde{CD}_n) &= (\lambda + 1)^2(\lambda^{n-1} + 1) - 2\lambda(\lambda + 1)(\lambda^{n-2} + 1) = \\
 &= (\lambda + 1)(\lambda^n + \lambda + \lambda^{n-1} + 1 - 2\lambda^{n-1} - 2\lambda) = \\
 &= (\lambda + 1)(\lambda^n - \lambda^{n-1} - \lambda + 1) = \\
 &= (\lambda + 1)(\lambda - 1)(\lambda^{n-1} - 1) = (\lambda - 1)^2\chi_1\chi_{n-2},
 \end{aligned} \tag{5.37}$$

and, up to a sign, we have

$$\chi = (\lambda - 1)^2\chi_1\chi_{n-2}. \tag{5.38}$$

Polynomials χ_1 and χ_{n-2} have, respectively, eigenvalues of orders 2 and 3 which are equal to the lengths of the branches A_1 and A_{n-2} of B_n without non-homogeneous branch point.

Let $g = 2$.

3) Cases $\tilde{G}_{12}$, $\tilde{G}_{22}$. Since the spectra of the Coxeter transformations of the dual graphs coincide, we consider $\tilde{G}_{12}$. By (4.11), splitting the diagram $\tilde{G}_{12}$ along the weighted edge we

have

$$\begin{aligned}\mathcal{X}(\tilde{G}_{12}) &= -(\chi_2\chi_1 - 3\lambda\chi_1) = -\chi_1(\chi_2 - 3\lambda) = \\ &= -(\lambda + 1)(\lambda^2 + \lambda + 1 - 3\lambda) = \\ &= -(\lambda - 1)^2(\lambda + 1) = -(\lambda - 1)^2\chi_1,\end{aligned}\tag{5.39}$$

and, up to a sign, we have

$$\chi = (\lambda - 1)^2\chi_1.\tag{5.40}$$

Polynomial χ_1 has the single eigenvalue of order 2, it corresponds to the length of the single branch A_1 of G_2 without non-homogeneous branch point.

4) Cases $\tilde{B}_n, \tilde{C}_n, \widetilde{BC}_n$. The characteristic polynomials of the Coxeter transformations of these diagrams coincide. Consider $\tilde{B}_n$. By (4.11), splitting the diagram $\tilde{B}_n$ along the weighted edge we have

$$\begin{aligned}\mathcal{X}(\tilde{B}_n) &= \mathcal{X}(B_n)\mathcal{X}(A_1) - 2\lambda\mathcal{X}(B_{n-1}), \text{ i.e.,} \\ (-1)^{n-1}\mathcal{X}(\tilde{B}_n) &= (\lambda^n + 1)(\lambda + 1) - 2\lambda(\lambda^{n-1} + 1) = \\ &= \lambda^{n+1} + \lambda^n + \lambda + 1 - 2\lambda^{n-1} - 2\lambda = \\ &= \lambda^{n+1} - \lambda^n - \lambda + 1 = (\lambda - 1)(\lambda^n - 1),\end{aligned}\tag{5.41}$$

and, up to a sign, we have

$$\chi = (\lambda - 1)^2\chi_{n-1}.\tag{5.42}$$

Polynomial χ_{n-1} has the single eigenvalue of order n , it corresponds to the length of the single branch A_{n-1} of B_n without non-homogeneous branch point.

Let $g = 3$.

5) Cases $\tilde{A}_{11}, \tilde{A}_{12}$. Direct calculation of the Coxeter transformation gives

$$\begin{aligned}\text{for } \tilde{A}_{11} : \mathbf{C} &= \begin{pmatrix} -1 & 4 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 3 & -4 \\ 1 & -1 \end{pmatrix}, \\ \text{for } \tilde{A}_{12} : \mathbf{C} &= \begin{pmatrix} -1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2 & -1 \end{pmatrix} = \begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix},\end{aligned}$$

and in both cases we have

$$\chi = (\lambda - 3)(\lambda + 1) + 4 = (\lambda - 1)^2. \quad \square\tag{5.43}$$

5.4. The Kostant generating function and Poincaré series

5.4.1. The generating function. Let $\text{Sym}(\mathbb{C}^2)$ be the symmetric algebra over $\mathbb{C}^2$, in other words, $\text{Sym}(\mathbb{C}^2) = \mathbb{C}[x_1, x_2]$, see (2.27). The symmetric algebra $\text{Sym}(\mathbb{C}^2)$ is a graded $\mathbb{C}$ -algebra, see (2.28):

$$\text{Sym}(\mathbb{C}^2) = \bigoplus_{n=0}^{\infty} \text{Sym}^n(\mathbb{C}^2).\tag{5.44}$$

Let π_n be the representation of $SU(2)$ in $\text{Sym}^n(\mathbb{C}^2)$ induced by its action on $\mathbb{C}^2$. The set $\{\pi_n\}$, where $n = 0, 1, \dots$ is the set of all irreducible representations of $SU(2)$, see, e.g., [Zhe73, §37]. Let G be any finite subgroup of $SU(2)$, see §(A.1). B. Kostant in [Kos84] considered the question:

how does $\pi_n|G$ decompose for any $n \in \mathbb{N}$?

The answer — the decomposition $\pi_n|G$ — is as follows:

$$\pi_n|G = \sum_{i=0}^r m_i(n) \rho_i, \quad (5.45)$$

where ρ_i are irreducible representations of G , considered in the context of *McKay correspondence*, see §(A.4). Thus, the decomposition (5.45) reduces the question to the following one:

what are the multiplicities $m_i(n)$ equal to?

B. Kostant in [Kos84] obtained the multiplicities $m_i(n)$ by means the orbit structure of the Coxeter transformation on the highest root of the corresponding Lie algebra. For further details concerning this orbit structure and the multiplicities $m_i(n)$, see §B.6.

Note, that multiplicities $m_i(n)$ in (5.45) are calculated as follows:

$$m_i(n) = \langle \pi_n|G, \rho_i \rangle, \quad (5.46)$$

(for the definition of the inner product $\langle \cdot, \cdot \rangle$, see (5.59)).

REMARK 5.6. For further considerations, we extend the relation for multiplicity (5.46) to the cases of *restricted representations* $\rho_i^\downarrow := \rho_i \downarrow_H^G$ and *induced representations* $\rho_i^\uparrow := \rho_i \uparrow_H^G$, where H is any subgroup of G (see §A.5.1):

$$m_i^\downarrow(n) = \langle \pi_n|H, \rho_i^\downarrow \rangle, \quad m_i^\uparrow(n) = \langle \pi_n|G, \rho_i^\uparrow \rangle. \quad (5.47)$$

We do not have any decomposition like (5.45) neither for restricted representations $\rho_i^\downarrow$ nor for induced representations $\rho_i^\uparrow$. Nevertheless, we will sometimes denote both multiplicities $m_i^\downarrow(n)$ and $m_i^\uparrow(n)$ in (5.47) by $m_i(n)$ as in (5.45).

REMARK 5.7. 1) A representation $\rho : G \longrightarrow GL_k(V)$ defines a k -linear action G on V by

$$gv = \rho(g)v. \quad (5.48)$$

The pair (V, ρ) is called a G -module. The case where $\rho(g) = Id_V$ is called the *trivial representation* in V . In this case

$$gv = v \text{ for all } g \in G. \quad (5.49)$$

In (5.45), the trivial representation ρ_0 corresponds to a particular vertex (see [McK80]), which extends the Dynkin diagram to the extended Dynkin diagram.

2) Let $\rho_0(H)$ (resp. $\rho_0(G)$) be the trivial representation of any subgroup $H \subset G$ (resp. of group G). The trivial representation $\rho_0(H) : H \longrightarrow GL_k(V)$ coincides with the *restricted representation* $\rho_0 \downarrow_H^G : G \longrightarrow GL_k(V)$, and the trivial representation $\rho_0(G) : G \longrightarrow GL_k(V)$ coincides with the *induced representation* $\rho_0 \uparrow_H^G : H \longrightarrow GL_k(V)$.

Since there is one-to-one correspondence between the ρ_i and the vertices of the Dynkin diagram, we can define (see [Kos84, p.211]) the vectors v_n , where $n \in \mathbb{Z}_+$, as follows:

$$v_n = \sum_{i=0}^r m_i(n) \alpha_i, \text{ where } \pi_n|G = \sum_{i=0}^r m_i(n) \rho_i, \quad (5.50)$$

where α_i are simple roots of the corresponding extended Dynkin diagram. Similarly, for the multiply-laced case, we define vectors v_n to be:

$$v_n = \sum_{i=0}^r m_i^\uparrow(n) \alpha_i \quad \text{or} \quad v_n = \sum_{i=0}^r m_i^\downarrow(n) \alpha_i, \quad (5.51)$$

where the multiplicities $m_i^\uparrow(n)$ and $m_i^\downarrow(n)$ are defined by (5.47). The vector v_n belongs to the root lattice generated by simple roots. Following B. Kostant, we define the generating function $P_G(t)$ for cases (5.50) and (5.51) as follows:

$$P_G(t) = ([P_G(t)]_0, [P_G(t)]_1, \dots, [P_G(t)]_r)^t := \sum_{n=0}^{\infty} v_n t^n, \quad (5.52)$$

the components of the vector $P_G(t)$ being the following series

$$[P_G(t)]_i = \sum_{n=0}^{\infty} \tilde{m}_i(n) t^n, \quad (5.53)$$

where $i = 0, 1, \dots, r$ and $\tilde{m}_i(n)$ designates $m_i(n)$, $m_i^\uparrow(n)$ or $m_i^\downarrow(n)$, see §1.5.2. In particular, for $i = 0$, we have

$$[P_G(t)]_0 = \sum_{n=0}^{\infty} m_0(n) t^n, \quad (5.54)$$

where $m_0(n)$ is the multiplicity of the trivial representation ρ_0 (see Remark 5.7) in $\text{Sym}^n(\mathbb{C}^2)$. By §2.7 the *algebra of invariants* R^G is a subalgebra of the *symmetric algebra* $\text{Sym}(\mathbb{C}^2)$. Thanks to (5.49), we see that R^G coincides with $\text{Sym}(\mathbb{C}^2)$, and $[P_G(t)]_0$ is the Poincaré series of the algebra of invariants $\text{Sym}(\mathbb{C}^2)^G$, i.e.,

$$[P_G(t)]_0 = P(\text{Sym}(\mathbb{C}^2)^G, t). \quad (5.55)$$

(see [Kos84, p.221, Rem.3.2]).

REMARK 5.8. According to Remark 5.7, heading 2), we have

$$[P_H(t)]_0 = P(\text{Sym}(\mathbb{C}^2)^{\rho_0 \downarrow_H^G}, t), \quad [P_G(t)]_0 = P(\text{Sym}(\mathbb{C}^2)^{\rho_0 \uparrow_H^G}, t). \quad (5.56)$$

We will need this fact in the proof of W. Ebeling's theorem for the multiply-laced case in §5.4.3, (see also Remark 5.6 and §A.5.1).

The following theorem gives a remarkable formula for calculating the Poincaré series for the binary polyhedral groups. The theorem is known in different forms. B. Kostant in [Kos84] shows it in the context of the Coxeter number h .

Theorem K-K-GV (Kostant, Knörrer, Gonsales-Sprinberg, Verdier) The Poincaré series $[P_G(t)]_0$ can be calculated as the following rational function:

$$[P_G(t)]_0 = \frac{1 + t^h}{(1 - t^a)(1 - t^b)}, \quad (5.57)$$

where

$$b = h + 2 - a, \text{ and } ab = 2|G|. \quad (5.58)$$

For a proof, see Theorem 1.4 and Theorem 1.8 from [Kos84], [Kn85, p.185], [GV83, p.428]. We call the numbers a and b the *Kostant numbers*. They can be easily calculated, see Table 5.1, compare also with Table A.2 and Table A.3. Note, that $a = 2d$, where d is the maximal coordinate of the *nil-root* vector from the kernel of the Tits form, see §2.4 and Fig. 2.2.

TABLE 5.1. The binary polyhedral groups (BPG) and the Kostant numbers a, b

Dynkin diagram	Order of group	BPG	Coxeter number	a	b
A_{n-1}	n	$\mathbb{Z}/n\mathbb{Z}$	n	2	n
D_{n+2}	$4n$	$\mathcal{D}_n$	$2n + 2$	4	$2n$
E_6	24	$\mathcal{T}$	12	6	8
E_7	48	$\mathcal{O}$	18	8	12
E_8	120	$\mathcal{J}$	30	12	20

5.4.2. The characters and the McKay operator. Let $\chi_1, \chi_2, \dots, \chi_r$ be all irreducible $\mathbb{C}$ -characters of a finite group G corresponding to irreducible representations $\rho_1, \rho_2, \dots, \rho_r$, and let χ_1 correspond to the trivial representation, i.e., $\chi_1(g) = 1$ for all $g \in G$.

All characters constitute the *character algebra* $C(G)$ of G since $C(G)$ is also a vector space over $\mathbb{C}$. An *hermitian inner product* $\langle \cdot, \cdot \rangle$ on $C(G)$ is defined as follows. For characters $\alpha, \beta \in C(G)$, let

$$\langle \alpha, \beta \rangle = \frac{1}{|G|} \sum_{g \in G} \alpha(g) \overline{\beta(g)} \quad (5.59)$$

Sometimes, we will write *inner product* $\langle \rho_i, \rho_j \rangle$ of the representations meaning actually the inner product of the corresponding characters $\langle \chi_{\rho_i}, \chi_{\rho_j} \rangle$. Let $z_{ijk} = \langle \chi_i \chi_j, \chi_k \rangle$, where $\chi_i \chi_j$ corresponds to the representation $\rho_i \otimes \rho_j$. It is known that z_{ijk} is the multiplicity of the representation ρ_k in $\rho_i \otimes \rho_j$ and $z_{ijk} = z_{jik}$. The numbers z_{ijk} are integer and are called the *structure constants*, see, e.g., [Kar92, p.765].

For every $i \in \{1, \dots, r\}$, there exists some $\hat{i} \in \{1, \dots, r\}$ such that

$$\chi_{\hat{i}}(g) = \overline{\chi_i(g)} \text{ for all } g \in G. \quad (5.60)$$

The character $\chi_i^\wedge$ corresponds to the *contragredient representation* $\rho_i^\wedge$ determined from the relation

$$\rho_i^\wedge(g) = \rho_i(g)^{\prime -1}. \quad (5.61)$$

We have

$$\langle \chi_i \chi_j, \chi_k \rangle = \langle \chi_i, \chi_j^\wedge \chi_k \rangle \quad (5.62)$$

since

$$\begin{aligned} \langle \chi_i \chi_j, \chi_k \rangle &= \frac{1}{|G|} \sum_{g \in G} \chi_i(g) \chi_j(g) \overline{\chi_k(g)} = \\ &= \frac{1}{|G|} \sum_{g \in G} \chi_i(g) \overline{(\chi_j(g) \chi_k(g))} = \langle \chi_i, \chi_j^\wedge \chi_k \rangle. \end{aligned}$$

REMARK 5.9. The group $SU(2)$ is the set of all unitary unimodular 2×2 matrices u , i.e.,

$$\begin{aligned} u^* &= u^{-1} \text{ (unitarity) }, \\ \det(u) &= 1 \text{ (unimodularity) }. \end{aligned}$$

The matrices $u \in SU(2)$ have the following form:

$$u = \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix}, \text{ and } u^* = \begin{pmatrix} a^* & -b \\ b^* & a \end{pmatrix}, \text{ where } aa^* + bb^* = 1, \quad (5.63)$$

see, e.g., [Ha89, Ch.9, §6]. The mutually inverse matrices u and u^{-1} are

$$u = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \text{ and } u^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}, \text{ where } ad - bc = 1. \quad (5.64)$$

Set

$$s = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \text{ then } s^{-1} = s^3 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (5.65)$$

For any $u \in SU(2)$, we have

$$sus^{-1} = u^{\prime -1}. \quad (5.66)$$

The element s is called the *Weyl element*.

According to (5.61) and (5.66) we see that every finite dimensional representation of the group $SL(2, \mathbb{C})$ (and hence, of $SU(2)$) is equivalent to its contragredient representation, see [Zhe73, §37, Rem.3]. Thus by (5.62), for representations ρ_i of any finite subgroup $G \subset SU(2)$, we have

$$\langle \chi_i \chi_j, \chi_k \rangle = \langle \chi_i, \chi_j \chi_k \rangle. \quad (5.67)$$

Relation (5.67) holds also for characters of restricted representations $\chi^\downarrow := \chi \downarrow_H^G$ and induced representations $\chi^\uparrow := \chi \uparrow_H^G$ (see Remark 5.6 and §A.5.1):

$$\begin{aligned} \langle \chi_i \chi_j^\downarrow, \chi_k \rangle_H &= \langle \chi_i, \chi_j^\downarrow \chi_k \rangle_H, \\ \langle \chi_i \chi_j^\uparrow, \chi_k \rangle_G &= \langle \chi_i, \chi_j^\uparrow \chi_k \rangle_G. \end{aligned} \quad (5.68)$$

Indeed, every restricted representation $\chi_j^\downarrow$ (resp. induced representation $\chi_j^\uparrow$) is decomposed into the direct sum of irreducible characters $\chi_s \in Irr(H)$ (here, $Irr(H)$ is the set of irreducible characters of H) with some integer coefficients a_s , for example:

$$\chi_j^\downarrow = \sum_{\chi_s \in Irr(H)} a_s \chi_s, \quad (5.69)$$

and since $\bar{a}_s = a_s$ for all $\chi_s \in \text{Irr}(H)$, we have

$$\begin{aligned} \langle \chi_i \chi_j^\downarrow, \chi_k \rangle_H &= \sum_{\chi_s \in \text{Irr}(H)} \langle a_s \chi_i \chi_s, \chi_k \rangle_H = \\ &= \sum_{\chi_s \in \text{Irr}(H)} a_s \langle \chi_i, \chi_s \chi_k \rangle_H = \\ &= \langle \chi_i, \sum_{\chi_s \in \text{Irr}(H)} a_s \chi_s \chi_k \rangle_H = \langle \chi_i, \chi_j^\downarrow \chi_k \rangle_H. \end{aligned} \quad (5.70)$$

The matrix of multiplicities $A := A(G)$ from (A.13) was introduced by J. McKay in [McK80]; it plays the central role in the *McKay correspondence*, see §A.4. We call this matrix — or the corresponding operator — the *McKay matrix* or the *McKay operator*.

Similarly, let $\tilde{A}$ and $\tilde{A}^\vee$ be matrices of multiplicities (A.14), (A.16). These matrices were introduced by P. Slodowy [Sl80] by analogy with the McKay matrix for the multiply-laced case, see §A.5.1. We call these matrices the *Slodowy operators*.

The following result of B. Kostant [Kos84], which holds for the McKay operator holds also for the Slodowy operators.

PROPOSITION 5.10. *If B is either the McKay operator A or the Slodowy operator $\tilde{A}$ or $\tilde{A}^\vee$, then*

$$Bv_n = v_{n-1} + v_{n+1}. \quad (5.71)$$

Proof. From now on

$$\rho_i = \begin{cases} \rho_i & \text{for } B = A, \\ \rho_i^\downarrow & \text{for } B = \tilde{A}, \\ \rho_i^\uparrow & \text{for } B = \tilde{A}^\vee, \end{cases} \quad m_i(n) = \begin{cases} m_i(n) & \text{for } B = A, \\ m_i^\downarrow(n) & \text{for } B = \tilde{A}, \\ m_i^\uparrow(n) & \text{for } B = \tilde{A}^\vee. \end{cases} \quad (5.72)$$

By (5.50), (5.51), and by definition of the McKay operator (A.13) and by definition of the Slodowy operator (A.14) and (A.16), we have

$$Bv_n = B \begin{pmatrix} m_0(n) \\ \vdots \\ m_r(n) \end{pmatrix} = \begin{pmatrix} \sum a_{0i} m_i(n) \\ \vdots \\ \sum a_{ri} m_i(n) \end{pmatrix} = \begin{pmatrix} \sum a_{0i} \langle \rho_i, \pi_n \rangle \\ \vdots \\ \sum a_{ri} \langle \rho_i, \pi_n \rangle \end{pmatrix}. \quad (5.73)$$

By (A.13), (A.14) and (A.16) we have

$$\sum_{i=1}^r a_{0i} \langle \rho_i, \pi_n \rangle = \langle \sum_{i=1}^r a_{0i} \rho_i, \pi_n \rangle = \langle \rho_{reg} \otimes \rho_0, \pi_n \rangle,$$

and from (5.73) we obtain

$$Bv_n = \begin{pmatrix} \langle \rho_{reg} \otimes \rho_0, \pi_n \rangle \\ \vdots \\ \langle \rho_{reg} \otimes \rho_r, \pi_n \rangle \end{pmatrix}. \quad (5.74)$$

Here ρ_{reg} is the regular two-dimensional representation which coincides with the representation π_1 in $\text{Sym}^2(\mathbb{C}^2)$ from §5.4.1. Thus,

$$Bv_n = \begin{pmatrix} \langle \pi_1 \otimes \rho_0, \pi_n \rangle \\ \vdots \\ \langle \pi_1 \otimes \rho_r, \pi_n \rangle \end{pmatrix}. \quad (5.75)$$

From (5.67), (5.70) we obtain

$$Bv_n = \begin{pmatrix} \langle \rho_0, \pi_1 \otimes \pi_n \rangle \\ \vdots \\ \langle \rho_r, \pi_1 \otimes \pi_n \rangle \end{pmatrix}. \quad (5.76)$$

By Clebsch-Gordan formula we have

$$\pi_1 \otimes \pi_n = \pi_{n-1} \oplus \pi_{n+1}, \quad (5.77)$$

where π_{-1} is the zero representation, see [Sp77, exs.3.2.4] or [Ha89, Ch.5, §6,§7]. From (5.76) and (5.77) we have (5.71). $\square$

For the following corollary, see [Kos84, p.222] and also [Sp87, §4.1].

COROLLARY 5.11. *Let $x = P_G(t)$ be given by (5.52). Then*

$$tBx = (1 + t^2)x - v_0, \quad (5.78)$$

where B is either the McKay operator A or the Slodowy operators $\tilde{A}$, $\tilde{A}^\vee$.

Proof. From (5.71) we obtain

$$\begin{aligned} Bx &= \sum_{n=0}^{\infty} Bv_n t^n = \sum_{n=0}^{\infty} (v_{n-1} + v_{n+1}) t^n = \sum_{n=0}^{\infty} v_{n-1} t^n + \sum_{n=0}^{\infty} v_{n+1} t^n = \\ &= t \sum_{n=1}^{\infty} v_{n-1} t^{n-1} + t^{-1} \sum_{n=0}^{\infty} v_{n+1} t^{n+1} = tx + t^{-1} \left(\sum_{n=0}^{\infty} v_n t^n - v_0 \right) = \\ &= tx + t^{-1}x - t^{-1}v_0. \quad \square \end{aligned} \quad (5.79)$$

5.4.3. The Poincaré series and W. Ebeling's theorem. W. Ebeling in [Ebl02] makes use of the Kostant relation (5.71) and deduces a new remarkable fact about the Poincaré series, a fact that shows that the Poincaré series of a binary polyhedral group (see (5.57)) is the quotient of two polynomials: the characteristic polynomial of the Coxeter transformation and the characteristic polynomial of the corresponding affine Coxeter transformation, see [Ebl02, Th.2].

We show W. Ebeling's theorem also for the multiply-laced case, see Theorem 5.12. The Poincaré series for the multiply-laced case is defined by (5.56).

THEOREM 5.12 (generalized W.Ebeling's theorem [Ebl02]). *Let G be a binary polyhedral group and $[P_G(t)]_0$ the Poincaré series (5.55) of the algebra of invariants $\text{Sym}(\mathbb{C}^2)^G$. Then*

$$[P_G(t)]_0 = \frac{\det M_0(t)}{\det M(t)}, \quad (5.80)$$

where

$$\det M(t) = \det |t^2 I - \mathbf{C}_a|, \quad \det M_0(t) = \det |t^2 I - \mathbf{C}|, \quad (5.81)$$

$\mathbf{C}$ is the Coxeter transformation and $\mathbf{C}_a$ is the corresponding affine Coxeter transformation.

Proof. By (5.78) we have

$$[(1 + t^2)I - tB]x = v_0, \quad (5.82)$$

where x is the vector $P_G(t)$ and by Cramer's rule the first coordinate $P_G(t)$ is

$$[P_G(t)]_0 = \frac{\det M_0(t)}{\det M(t)}, \quad (5.83)$$

where

$$\det M(t) = \det ((1 + t^2)I - tB), \quad (5.84)$$

TABLE 5.2. The characteristic polynomials χ , $\tilde{\chi}$ and the Poincaré series

Dynkin diagram	Coxeter transformation χ	Affine Coxeter transformation $\tilde{\chi}$	Quotient $p(\lambda) = \frac{\chi}{\tilde{\chi}}$
D_4	$(\lambda + 1)(\lambda^3 + 1)$	$(\lambda - 1)^2(\lambda + 1)^3$	$\frac{\lambda^3 + 1}{(\lambda^2 - 1)^2}$
D_{n+1}	$(\lambda + 1)(\lambda^n + 1)$	$(\lambda^{n-1} - 1)(\lambda - 1)(\lambda + 1)^2$	$\frac{\lambda^n + 1}{(\lambda^{n-1} - 1)(\lambda^2 - 1)}$
E_6	$\frac{(\lambda^6 + 1)(\lambda^3 - 1)}{(\lambda^2 + 1)(\lambda - 1)}$	$(\lambda^3 - 1)^2(\lambda + 1)$	$\frac{\lambda^6 + 1}{(\lambda^4 - 1)(\lambda^3 - 1)}$
E_7	$\frac{(\lambda + 1)(\lambda^9 + 1)}{(\lambda^3 + 1)}$	$(\lambda^4 - 1)(\lambda^3 - 1)(\lambda + 1)$	$\frac{\lambda^9 + 1}{(\lambda^4 - 1)(\lambda^6 - 1)}$
E_8	$\frac{(\lambda^{15} + 1)(\lambda + 1)}{(\lambda^5 + 1)(\lambda^3 + 1)}$	$(\lambda^5 - 1)(\lambda^3 - 1)(\lambda + 1)$	$\frac{\lambda^{15} + 1}{(\lambda^{10} - 1)(\lambda^6 - 1)}$
B_n	$\lambda^n + 1$	$(\lambda^{n-1} - 1)(\lambda^2 - 1)$	$\frac{\lambda^n + 1}{(\lambda^{n-1} - 1)(\lambda^2 - 1)}$
C_n	$\lambda^n + 1$	$(\lambda^n - 1)(\lambda - 1)$	$\frac{\lambda^n + 1}{(\lambda^n - 1)(\lambda - 1)}$
F_4	$\frac{\lambda^6 + 1}{\lambda^2 + 1}$	$(\lambda^2 - 1)(\lambda^3 - 1)$	$\frac{\lambda^6 + 1}{(\lambda^4 - 1)(\lambda^3 - 1)}$
G_2	$\frac{\lambda^3 + 1}{\lambda + 1}$	$(\lambda - 1)^2(\lambda + 1)$	$\frac{\lambda^3 + 1}{(\lambda^2 - 1)^2}$
A_n	$\frac{\lambda^{n+1} - 1}{\lambda - 1}$	$(\lambda^{n-k+1} - 1)(\lambda^k - 1)$	$\frac{\lambda^{n+1} - 1}{(\lambda - 1)(\lambda^{n-k+1} - 1)(\lambda^k - 1)}$
A_{2n-1}	$\frac{\lambda^{2n} - 1}{\lambda - 1}$	$(\lambda^n - 1)^2$ for $k = n$	$\frac{\lambda^n + 1}{(\lambda^n - 1)(\lambda - 1)}$

and $M_0(t)$ is the matrix obtained by replacing the first column of $M(t)$ by $v_0 = (1, 0, \dots, 0)^t$. The vector v_0 corresponds to the trivial representation π_0 , and by the McKay correspondence, v_0 corresponds to the particular vertex which extends the Dynkin diagram to the extended Dynkin diagram, see Remark 5.7 and (5.50). Therefore, if $\det M(t)$ corresponds to the affine Coxeter transformation, and

$$\det M(t) = \det |t^2 I - \mathbf{C}_a|, \quad (5.85)$$

then $\det M_0(t)$ corresponds to the Coxeter transformation, and

$$\det M_0(t) = \det |t^2 I - \mathbf{C}|. \quad (5.86)$$

So, it suffices to prove (5.85), i.e.,

$$\det[(1 + t^2)I - tB] = \det |t^2 I - \mathbf{C}_a|. \quad (5.87)$$

If B is the McKay operator A given by (A.13), then

$$B = 2I - K = \begin{pmatrix} 2I & 0 \\ 0 & 2I \end{pmatrix} - \begin{pmatrix} 2I & 2D \\ 2D^t & 2I \end{pmatrix} = \begin{pmatrix} 0 & -2D \\ -2D^t & 0 \end{pmatrix}, \quad (5.88)$$

where K is a symmetric Cartan matrix (3.2). If B is the Slodowy operator $\tilde{A}$ or $\tilde{A}^\vee$ given by (A.14), (A.16), then

$$B = 2I - K = \begin{pmatrix} 2I & 0 \\ 0 & 2I \end{pmatrix} - \begin{pmatrix} 2I & 2D \\ 2F & 2I \end{pmatrix} = \begin{pmatrix} 0 & -2D \\ -2F & 0 \end{pmatrix}, \quad (5.89)$$

where K is the symmetrizable but not symmetric Cartan matrix (3.4). Thus, in the generic case

$$M(t) = (1 + t^2)I - tB = \begin{pmatrix} 1 + t^2 & 2tD \\ 2tF & 1 + t^2 \end{pmatrix}. \quad (5.90)$$

Assuming $t \neq 0$ we deduce from (5.90) that

$$\begin{aligned} M(t) \begin{pmatrix} x \\ y \end{pmatrix} = 0 &\iff \begin{cases} (1 + t^2)x = -2tDy, \\ 2tFx = -(1 + t^2)y. \end{cases} \\ &\iff \begin{cases} \frac{(1 + t^2)^2}{4t^2}x = FDy, \\ \frac{(1 + t^2)^2}{4t^2}y = DFy. \end{cases} \end{aligned} \quad (5.91)$$

Here we use the Jordan form theory of the Coxeter transformations constructed in §3. According to (3.14), Proposition 3.3 and Proposition 3.9 we see that t^2 is an eigenvalue of the affine Coxeter transformation $\mathbf{C}_a$, i.e., (5.87) together with (5.85) are proved. $\square$

For the results of calculations using W. Ebeling's theorem, see Table 5.2.

REMARK 5.13. 1) The characteristic polynomials χ for the Coxeter transformation and $\tilde{\chi}$ for the affine Coxeter transformation in Table 5.2 are taken from Tables 1.1 and 1.2. Pay attention to the fact that the affine Dynkin diagram for B_n is $\widetilde{C\bar{D}}_n$, ([Bo, Tab.2]), and the affine Dynkin diagram for C_n is $\tilde{C}_n$, ([Bo, Tab.3]), see Fig. 2.2.

2) The characteristic polynomial χ for the affine Coxeter transformation of A_n depends on the *index of the conjugacy class* k of the Coxeter transformation, see (4.5). In the case of A_n (for every $k = 1, 2, \dots, n$) the quotient $p(\lambda) = \frac{\chi}{\tilde{\chi}}$ contains three factors in the denominator, and its form is different from (5.57), see Table 5.2.

For the case A_{2n-1} and $k = n$, we have

$$p(\lambda) = \frac{\lambda^{2n} - 1}{(\lambda - 1)(\lambda^{2n-k} - 1)(\lambda^k - 1)} = \frac{\lambda^{2n} - 1}{(\lambda - 1)(\lambda^n - 1)(\lambda^n - 1)} = \frac{\lambda^n + 1}{(\lambda^n - 1)(\lambda - 1)} \quad (5.92)$$

and $p(\lambda)$ again is of the form (5.57), see Table 5.2.

3) The quotients $p(\lambda)$ coincide for the following pairs:

$$\begin{array}{ll} D_4 \text{ and } G_2, & E_6 \text{ and } F_4, \\ D_{n+1} \text{ and } B_n (n \geq 4), & A_{2n-1} \text{ and } C_n. \end{array} \quad (5.93)$$

Note that the second elements of the pairs are obtained by *folding* operation from the first ones, see Remark 5.4.

CHAPTER 6

Regular representations of quivers

6.1. The regular representations and the Dlab-Ringel criterion

The regular representations are the most complicated in the category of all representation of the given quiver. For every Dynkin diagram the category of regular representations is empty, there are only finite number non-regular representations (P.Gapriel's theorem [Gab72]). For this reason, the Dynkin diagrams are called *finite type quivers* in the representation theory of quivers. The regular representations have been completely described only for the extended Dynkin diagrams, which for this reason were dubbed *tame quivers* in the representation theory of quivers, ([Naz73], [DR76]), see §1.6.1.

DEFINITION 6.1. 1) The vector $z \in \mathcal{E}_\Gamma$ is said to be Δ' -regular or *regular in the orientation* Δ' if

$$\mathbf{C}_{\Delta'}^k z > 0 \text{ for all } k \in \mathbb{Z}. \quad (6.1)$$

2) The representation V is said to be a *regular representation in the orientation* Δ' of the quiver Γ if its dimension $\dim V$ is a Δ' -regular vector, i.e.,

$$\mathbf{C}_{\Delta'}^k(\dim V) > 0 \text{ for all } k \in \mathbb{Z}. \quad (6.2)$$

Let the Coxeter transformation $\mathbf{C}_{\Delta'}$ correspond to any orientation Δ' and let the Coxeter transformation $\mathbf{C}_\Delta$ correspond to a bicolored orientation Δ . Then $\mathbf{C}_{\Delta'}$ and $\mathbf{C}_\Delta$ are conjugate if the quiver is a tree. Let T be an element in the Weyl group which relates $\mathbf{C}_{\Delta'}$ and $\mathbf{C}_\Delta$:

$$\mathbf{C}_{\Delta'} = T^{-1} \mathbf{C}_\Delta T. \quad (6.3)$$

6.2. Necessary conditions of regularity of the extended Dynkin diagrams

Let $(\alpha_1, \tilde{\alpha}_1, \alpha_1^{\varphi_2}, \alpha_2^{\varphi_2}, \dots)$ be coordinates of the vector Tz in the Jordan basis of eigenvectors and adjoint vectors (3.29) – (3.31):

$$Tz = \alpha_1 z^1 + \tilde{\alpha}_1 \tilde{z}^1 + \alpha_1^{\varphi_2} z_1^{\varphi_2} + \alpha_2^{\varphi_2} z_2^{\varphi_2} + \dots \quad (6.4)$$

According to (3.32), (3.33), (3.34), we have

$$\mathbf{C}_\Delta^k Tz = \alpha_1 z^1 + \tilde{\alpha}_1 (kz^1 + \tilde{z}^1) + \alpha_1^{\varphi_2} (\lambda_1^{\varphi_2})^k z_1^{\varphi_2} + \alpha_2^{\varphi_2} (\lambda_2^{\varphi_2})^k z_2^{\varphi_2} + \dots \quad (6.5)$$

and

$$\begin{aligned} \mathbf{C}_{\Delta'}^k z &= T^{-1} \mathbf{C}_\Delta^k Tz = \\ &= \alpha_1 z^1 + \tilde{\alpha}_1 (kz^1 + T^{-1} \tilde{z}^1) + \\ &+ \alpha_1^{\varphi_2} (\lambda_1^{\varphi_2})^k T^{-1} z_1^{\varphi_2} + \alpha_2^{\varphi_2} (\lambda_2^{\varphi_2})^k T^{-1} z_2^{\varphi_2} + \dots \end{aligned} \quad (6.6)$$

Since $z^1 > 0$, $k \in \mathbb{Z}$, and $|\lambda_{1,2}^{\varphi_i}| = 1$, we have the following necessary condition of Δ' -regularity:

$$\tilde{\alpha}_1 = 0, \quad (6.7)$$

where $\tilde{\alpha}_1$ is the coordinate of the vector Tz corresponding to the adjoint basis vector $\tilde{z}^1$.

a) Simply-laced case. The adjoint vector $\tilde{z}^1$ is orthogonal to vectors $z_i^{\varphi_j}$ for $i = 1, 2$; the numbers φ_j are eigenvalues of DD^t since the corresponding components $\mathbb{X}$ and $\mathbb{Y}$ are orthogonal, see Proposition 3.8. Further, $\tilde{z}^1$ is also orthogonal to z^1 : indeed,

$$\begin{aligned} \langle z^1, \tilde{z}^1 \rangle &= \langle \mathbb{X}^1, \mathbb{X}^1 \rangle - \langle D^t \mathbb{X}^1, D^t \mathbb{X}^1 \rangle = \\ &= \langle \mathbb{X}^1, \mathbb{X}^1 \rangle - \langle \mathbb{X}^1, DD^t \mathbb{X}^1 \rangle = \langle \mathbb{X}^1, \mathbb{X}^1 \rangle - \langle \mathbb{X}^1, \mathbb{X}^1 \rangle = 0. \end{aligned}$$

Thus, from (6.4) we have

$$\langle Tz, \tilde{z}^1 \rangle = \tilde{\alpha}_1 \langle \tilde{z}^1, \tilde{z}^1 \rangle \quad (6.8)$$

and (6.7) is equivalent to the following relation:

$$\langle Tz, \tilde{z}^1 \rangle = 0. \quad (6.9)$$

b) Multiply-laced case. The normal basis of DF (resp. FD) is not orthogonal, however, the dual graph will help us.

DEFINITION 6.2. Let u be an eigenvector of the Coxeter transformation given by (3.29), (3.30). The vector $\tilde{u}$ is said to be *conjugate* to u , if it is obtained from u by changing the sign of $\mathbb{Y}$ -component and replacing the eigenvalue λ by $\frac{1}{\lambda}$.

(It is easily to see that the vectors z^1 and $4\tilde{z}^1$ are conjugate.)

PROPOSITION 6.3. 1) Let φ be an eigenvalue for DF and $\varphi^\vee \neq \varphi$ an eigenvalue for $D^\vee F^\vee$. The eigenvectors of DF and $D^\vee F^\vee$ corresponding to these eigenvalues are orthogonal.

2) Let z_φ be an eigenvector with eigenvalue $\varphi \neq 0$ for the extended Dynkin diagram Γ and $z_\varphi^\vee$ an eigenvector with eigenvalue φ for the dual diagram $\Gamma^\vee$. Let $\tilde{z}_\varphi^\vee$ be conjugate to $z_\varphi^\vee$. Then z_φ and $\tilde{z}_\varphi^\vee$ are orthogonal.

Proof. 1) Let $DFx = \varphi x$ and $D^\vee F^\vee x = \varphi^\vee x^\vee$. Since $\varphi \neq \varphi^\vee$, one of these eigenvalues is $\neq 0$. Let, for example, $\varphi \neq 0$. Then

$$\langle x, x^\vee \rangle = \frac{1}{\varphi} \langle DFx, x^\vee \rangle = \frac{1}{\varphi} \langle x, (DF)^t x^\vee \rangle. \quad (6.10)$$

From (6.10) and (3.8) we have

$$\langle x, x^\vee \rangle = \frac{1}{\varphi} \langle x, D^\vee F^\vee x^\vee \rangle = \frac{\varphi^\vee}{\varphi} \langle x, x^\vee \rangle.$$

Since $\varphi \neq \varphi^\vee$, we have $\langle x, x^\vee \rangle = 0$. $\square$

2) Let us express vectors z_φ and $\tilde{z}_\varphi^\vee$ as follows:

$$z_\varphi = \begin{pmatrix} x \\ -\frac{2}{\lambda+1}Fx \end{pmatrix}, \quad \tilde{z}_\varphi^\vee = \begin{pmatrix} x^\vee \\ \frac{2\lambda}{\lambda+1}F^\vee x^\vee \end{pmatrix}. \quad (6.11)$$

According to (3.7) and (3.8) we have

$$\begin{aligned} \langle z_\varphi, \tilde{z}_\varphi^\vee \rangle &= \langle x, x^\vee \rangle - \frac{4\lambda}{(\lambda+1)^2} \langle Fx, F^\vee x^\vee \rangle = \\ &= \langle x, x^\vee \rangle - \frac{1}{\varphi} \langle DFx, x^\vee \rangle = \langle x, x^\vee \rangle - \langle x, x^\vee \rangle = 0. \quad \square \end{aligned}$$

Thus, by Proposition 6.3 heading 1), the vector $\tilde{z}^{1\vee}$ is orthogonal to the z_i^φ for $i = 1, 2$ and by Proposition 6.3 heading 2), the vector $\tilde{z}^{1\vee}$ is orthogonal to z^1 . Therefore from (6.4) we deduce that

$$\langle Tz, \tilde{z}^{1\vee} \rangle = \tilde{\alpha}_1 \langle \tilde{z}^{1\vee}, \tilde{z}^{1\vee} \rangle \quad (6.12)$$

and (6.7) is equivalent to the following relation:

$$\langle Tz, \tilde{z}^{1\vee} \rangle = 0. \quad (6.13)$$

Relation (6.8) for the simply-laced case and relation (6.12) for the multiply-laced case motivate the following

DEFINITION 6.4. The linear form $\rho_{\Delta'}(z)$ given as follows

$$\rho_{\Delta'}(z) = \begin{cases} \langle Tz, \tilde{z}^1 \rangle & \text{for a simply-laced case,} \\ \langle Tz, \tilde{z}^{1\vee} \rangle & \text{for a multiply-laced case} \end{cases} \quad (6.14)$$

is said to be the Δ' -defect of the vector z .

From (6.9) and (6.13) we derive the following theorem:

THEOREM 6.5 ([SuSt75],[SuSt78]). *If z is a regular vector for the extended Dynkin diagram Γ in the orientation Δ' , then*

$$\rho_{\Delta'}(z) = 0. \quad (6.15)$$

Below, in Proposition 6.13, we will show that the condition (6.15) is also sufficient, if z is a root in the root system related to the given extended Dynkin diagram.

6.3. The Dlab-Ringel definition of the defect

The vector z^1 from the kernel of the Tits form is the fixed point for the Weyl group, so $T^{-1}z^1 = z^1$, and from (6.4) we deduce

$$z = \alpha_1 z^1 + \tilde{\alpha}_1 T^{-1} \tilde{z}^1 + \alpha_1^{\varphi_2} T^{-1} z_1^{\varphi_2} + \alpha_2^{\varphi_2} T^{-1} z_2^{\varphi_2} + \dots \quad (6.16)$$

and from (6.6) for $k = h$ being the Coxeter number, i.e., $(\lambda_{1,2}^{\varphi_2})^k = 1$, we obtain

$$\mathbf{C}_{\Delta'}^k z = \alpha_1 z^1 + \tilde{\alpha}_1 (kz^1 + T^{-1} \tilde{z}^1) + \alpha_1^{\varphi_2} T^{-1} z_1^{\varphi_2} + \alpha_2^{\varphi_2} T^{-1} z_2^{\varphi_2} + \dots \quad (6.17)$$

From (6.16) and (6.17) we get the following formula due to V. Dlab and C. M. Ringel, see [DR76]:

$$\mathbf{C}_{\Delta'}^h z = z + h \tilde{\alpha}_1 z^1, \quad (6.18)$$

where h is the Coxeter number and $\tilde{\alpha}_1 = \tilde{\alpha}_1(z)$ is the linear form proportional to $\rho_{\Delta'}(z)$. By (6.8) for the simply-laced case and by (6.12) for the multiply-laced case the form $\tilde{\alpha}_1(z)$ can be calculated as follows:

$$\tilde{\alpha}_1(z) = \begin{cases} \frac{\langle Tz, \tilde{z}^1 \rangle}{\langle \tilde{z}^1, \tilde{z}^1 \rangle} & \text{for a simply-laced case,} \\ \frac{\langle Tz, \tilde{z}^{1\vee} \rangle}{\langle \tilde{z}^{1\vee}, \tilde{z}^{1\vee} \rangle} & \text{for a multiply-laced case.} \end{cases} \quad (6.19)$$

Dlab and Ringel introduced in [DR76] the defect $\delta_{\Delta'}$ as a vector from $\mathcal{E}_{\Gamma}^*$ obtained as a solution of the equation

$$\mathbf{C}_{\Delta'}^* \delta_{\Delta'} = \delta_{\Delta'}. \quad (6.20)$$

PROPOSITION 6.6 ([St85]). *The Dlab-Ringel defect $\delta_{\Delta'}$ given by (6.20) coincides (up to coefficient) with the Δ' -defect $\rho_{\Delta'}$ given by Definition 6.4.*

Proof. 1) Let us show that $\delta_{\Delta'}$ is obtained from δ_{Δ} in the same way as $\rho_{\Delta'}$ is obtained from ρ_{Δ} , so it suffices to prove the proposition only for bicolored orientations Δ . Indeed,

$$\mathbf{C}_{\Delta'} = T^{-1}\mathbf{C}_{\Delta}T \quad \text{implies that} \quad \mathbf{C}_{\Delta'}^* = T^*\mathbf{C}_{\Delta}^*T^{-1*}.$$

Since $\mathbf{C}_{\Delta'}^*\delta_{\Delta'} = \delta_{\Delta'}$, we obtain

$$\mathbf{C}_{\Delta}^*T^{-1*}\delta_{\Delta'} = T^{-1*}\delta_{\Delta'},$$

and

$$T^{-1*}\delta_{\Delta'} = \delta_{\Delta}, \quad \text{i.e.,} \quad \delta_{\Delta'} = T^*\delta_{\Delta}.$$

Further,

$$\langle \delta_{\Delta}, Tz \rangle = \langle T^*\delta_{\Delta}, z \rangle = \langle \delta_{\Delta'}, z \rangle.$$

Thus, $\delta_{\Delta'}$ is obtained from δ_{Δ} in the same way as $\rho_{\Delta'}$ is obtained from ρ_{Δ} .

2) Now, let us prove the proposition for the bicolored orientation Δ , i.e., let us prove that ρ_{Δ} is proportional to δ_{Δ} . According to (3.1), (3.2) and (3.4) the relation

$$\mathbf{C}_{\Delta}^*z = z \quad \text{is equivalent to} \quad w_1^*z = w_2^*z,$$

which, in turn, is equivalent to the following:

$$\begin{pmatrix} -I & 0 \\ -2D^t & I \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} I & -2F \\ 0 & -I \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

By (3.7) we have

$$\begin{cases} x = F^ty = D^{\vee}y, \\ y = D^tx = F^{\vee}x. \end{cases}$$

Thus,

$$z = \begin{pmatrix} x \\ F^{\vee}x \end{pmatrix}, \quad \text{where } x = F^tD^tx = D^{\vee}F^{\vee}x,$$

i.e., x corresponds to $\lambda = 1$, an eigenvalue of $\mathbf{C}$, and to $\varphi = 1$, an eigenvalue of $(DF)^{\vee}$. By (6.11) we have $z = \tilde{z}^{1\vee}$, i.e., δ_{Δ} is proportional to ρ_{Δ} , see (6.14). $\square$

6.4. Necessary conditions of regularity for diagrams with indefinite Tits form

Now, consider the case where $\mathcal{B}$ is indefinite, i.e., Γ is any tree, which is neither Dynkin diagram nor extended Dynkin diagram.

Let $(\alpha_1^m, \alpha_2^m, \alpha_1^{\varphi_2}, \alpha_2^{\varphi_2}, \dots)$ be coordinates of the vector Tz in the Jordan basis of eigenvectors and adjoint vectors (3.29) – (3.31), where α_1^m and α_2^m are coordinates corresponding to eigenvectors z_1^m and z_2^m . The vectors z_1^m and z_2^m correspond to the maximal eigenvalue $\varphi^m = \varphi^{\max}$ of DD^t and D^tD , respectively. Let us decompose the vector Tz as follows:

$$Tz = \alpha_1^m z_1^m + \alpha_2^m z_2^m + \alpha_1^{\varphi_2} z_1^{\varphi_2} + \alpha_2^{\varphi_2} z_2^{\varphi_2} + \dots \quad (6.21)$$

According to (3.32), (3.33), (3.34) we have

$$\begin{aligned} \mathbf{C}_{\Delta}^k Tz = & (\lambda_1^{\varphi_m})^k \alpha_1^m z_1^m + (\lambda_2^{\varphi_m})^k \alpha_2^m z_2^m + (\lambda_1^{\varphi_2})^k \alpha_1^{\varphi_2} z_1^{\varphi_2} + (\lambda_1^{\varphi_2})^k \alpha_2^{\varphi_2} z_2^{\varphi_2} + \dots \end{aligned} \quad (6.22)$$

and

$$\begin{aligned} \mathbf{C}_{\Delta'}^k z = T^{-1}\mathbf{C}_{\Delta}^k Tz = & (\lambda_1^{\varphi_m})^k \alpha_1^m T^{-1}z_1^m + (\lambda_2^{\varphi_m})^k \alpha_2^m T^{-1}z_2^m + \\ & (\lambda_1^{\varphi_2})^k \alpha_1^{\varphi_2} T^{-1}z_1^{\varphi_2} + (\lambda_1^{\varphi_2})^k \alpha_2^{\varphi_2} T^{-1}z_2^{\varphi_2} + \dots \end{aligned} \quad (6.23)$$

It will be shown in Theorem 6.11 that the transforming element T can be modified so that its decomposition does not contain any given reflection σ_{α} . Since the coordinates of

the eigenvectors $z_{1,2}^m$ are all positive, see Corollary 3.7 and (3.32), we see that each vector $T^{-1}z_{1,2}^m$ has at least one positive coordinate. Besides,

$$|\lambda_1^{\varphi_m}| > |\lambda_{1,2}^{\varphi_j}| > |\lambda_2^{\varphi_m}| \quad (6.24)$$

because

$$\begin{aligned} \lambda_1^{\varphi_m} &= \frac{1}{\lambda_2^{\varphi_m}} \quad \text{and} \\ \lambda_1^{\varphi_m} &= 2\varphi^m - 1 + 2\sqrt{\varphi^m(\varphi^m - 1)} > 2\varphi^i - 1 \pm 2\sqrt{\varphi^i(\varphi^i - 1)}. \end{aligned}$$

Thus, since $T^{-1}z_{1,2}^m$ have at least one positive coordinate, we deduce from (6.23) and (6.24) that

$$\alpha_1^{\varphi_m} \geq 0, \quad \alpha_2^{\varphi_m} \geq 0. \quad (6.25)$$

As above for the case $\mathcal{B} \geq 0$, let us calculate $\alpha_{1,2}^{\varphi_m}$. The vector $\tilde{z}_1^m$ conjugate to the vector z_1^m is orthogonal to the vectors $z_{1,2}^2, z_{1,2}^3, \dots$. Let us show that $\tilde{z}_1^m$ is also orthogonal to z_1^m . Indeed, the vectors z_1^m and $\tilde{z}_1^m$ can be expressed as follows:

$$z_1^m = \begin{pmatrix} x^m \\ -\frac{2}{\lambda_1^m + 1} D^t x^m \end{pmatrix}, \quad \tilde{z}_1^m = \begin{pmatrix} x^m \\ \frac{2\lambda_1^m}{\lambda_1^m + 1} D^t x^m \end{pmatrix}, \quad (6.26)$$

where for brevity we designate $\lambda_1^{\varphi_{max}}$ by λ_1^m , $z_1^{\varphi_{max}}$ by z_1^m , and $x^{\varphi_{max}}$ by x^m . Then,

$$\begin{aligned} < z_1^m, \tilde{z}_1^m > = \\ < x^m, x^m > - \frac{4\lambda_1^m}{(\lambda_1^m + 1)^2} < D^t x^m, D^t x^m > = \\ < x^m, x^m > - \frac{1}{\varphi_m} < x^m, DD^t x^m > = \\ < x^m, x^m > - < x^m, x^m > = 0. \end{aligned} \quad (6.27)$$

The conjugate vector $\tilde{z}_1^m$ is not orthogonal only to z_2^m in decomposition (6.21), and similarly $\tilde{z}_2^m$ is not orthogonal only to z_1^m . From (6.21) we get

$$< Tz, \tilde{z}_1^m > = \alpha_2^m < z_2^m, \tilde{z}_1^m >, \quad < Tz, \tilde{z}_2^m > = \alpha_1^m < z_1^m, \tilde{z}_2^m >. \quad (6.28)$$

Let us find $< z_2^m, \tilde{z}_1^m >$ and $< z_1^m, \tilde{z}_2^m >$. We have:

$$\begin{aligned} < z_2^m, \tilde{z}_1^m > &= < x^m, x^m > - \frac{4}{(\lambda_2^m + 1)^2} < x^m, DD^t x^m > = \\ < x^m, x^m > &(1 - \frac{4}{(\lambda_2^m + 1)^2} \frac{(\lambda_2^m + 1)^2}{4(\lambda_2^m)^2}) = < x^m, x^m > (1 - (\lambda_1^m)^2). \end{aligned} \quad (6.29)$$

Similarly,

$$< z_1^m, \tilde{z}_2^m > = < x^m, x^m > (1 - (\lambda_2^m)^2). \quad (6.30)$$

Thus, from (6.28), (6.29) and (6.30) we get

$$\begin{aligned} < Tz, \tilde{z}_1^m > &= \alpha_2^m < x^m, x^m > (1 - (\lambda_1^m)^2), \\ < Tz, \tilde{z}_2^m > &= \alpha_1^m < x^m, x^m > (1 - (\lambda_2^m)^2). \end{aligned} \quad (6.31)$$

THEOREM 6.7 ([SuSt75], [SuSt78]). *If z is a regular vector for the graph Γ with indefinite Tits form $\mathcal{B}$ in the orientation Δ' , then*

$$< Tz, \tilde{z}_1^m > \leq 0, \quad < Tz, \tilde{z}_2^m > \geq 0. \quad (6.32)$$

Proof. Since $\lambda_1^m > 1$ and $\lambda_2^m < 1$, the theorem follows from (6.31) and (6.25). $\square$

We denote the linear form $\langle Tz, \tilde{z}_1^m \rangle$ (resp. $\langle Tz, \tilde{z}_2^m \rangle$) by $\rho_{\Delta'}^1$ (resp. $\rho_{\Delta'}^2$). Then, conditions (6.32) have the following form:

$$\rho_{\Delta'}^1(z) \leq 0, \quad \rho_{\Delta'}^2(z) \geq 0. \quad (6.33)$$

A similar results were obtained by Y. Zhang in [Zh89, Prop.1.5], and by J. A. de la Peña, M. Takane in [PT90, Th.2.3].

For an application of (6.32) to the star graph, see §6.6.4.

6.5. Transforming elements and sufficient conditions of regularity

In this section we consider only extended Dynkin diagrams. We will show in Theorem 6.14 that *the necessary condition of regularity of the vector z (6.15) coincides with the sufficient condition (Proposition 6.8) only if vector z is a positive root in the corresponding root system*. For an arbitrary vector z this is not true. Since dimension $\dim V$ of the indecomposable representation V is a positive root (the Gabriel theorem for the Dynkin diagrams [Gab72], [BGP73], the Dlab-Ringel theorem for the extended Dynkin diagram [DR76], [Naz73], the Kac theorem for any diagrams [Kac80], [Kac82]), then *for the indecomposable representation V the necessary condition of regularity of the vector $\dim V$ coincides with the sufficient condition*. For an arbitrary decomposable representation this is not true.

We start from the bicolored orientation Δ .

6.5.1. Sufficient conditions of regularity for the bicolored orientation.

PROPOSITION 6.8 ([St82]). *Let Γ be an extended Dynkin diagram, i.e., $\mathcal{B} \geq 0$. Let z be a root in the root system associated with Γ . If the Δ -defect of the vector z is zero:*

$$\rho_{\Delta}(z) = 0, \quad (6.34)$$

then z is regular in the bicolored orientation Δ .

Proof. Let

$$H = \{z \mid z > 0, z \text{ is a root}, \rho_{\Delta}(z) = 0\}. \quad (6.35)$$

It suffices to prove that

$$w_1 H \subset H, \quad w_2 H \subset H. \quad (6.36)$$

Indeed, if (6.36) holds, then

$$\mathbf{C}_{\Delta}^k H \subset H \quad \text{for all } k \in \mathbb{N},$$

hence $\mathbf{C}_{\Delta}^k z > 0$ if z is a positive root satisfying the condition (6.34).

So, let us prove, for example, that $w_1 H \subset H$. Note that z and $w_1 z$ are roots simultaneously. Thus, either $w_1 z > 0$ or $w_1 z < 0$. Suppose, $w_1 z < 0$. Together with $z > 0$, by (3.4) we have $y = 0$. Hence

$$\rho_{\Delta}(z) = \langle z, \tilde{z}^{1\vee} \rangle = \langle x, \tilde{x}^{1\vee} \rangle.$$

The coordinates of $\tilde{x}^{1\vee}$ are positive, the coordinates of x are non-negative. If $\rho_{\Delta}(z) = 0$, then $x = 0$; this contradicts to the condition $z > 0$. Therefore, $w_1 z > 0$. It remains to show that

$$\rho_{\Delta}(w_1 z) = 0.$$

Again, by (3.4), (3.7) and (3.30) we have

$$\begin{aligned} \langle w_1 z, \tilde{z}^{1\vee} \rangle &= \left\langle \begin{pmatrix} -x - 2Dy \\ y \end{pmatrix}, \begin{pmatrix} x^{1\vee} \\ F^{\vee} x^{1\vee} \end{pmatrix} \right\rangle = \\ &= -\langle x, x^{1\vee} \rangle - 2\langle y, D^t x^{1\vee} \rangle + \langle y, F^{\vee} x^{1\vee} \rangle = \\ &= -\langle x, x^{1\vee} \rangle - \langle y, F^{\vee} x^{1\vee} \rangle = -\langle z, \tilde{z}^{1\vee} \rangle = 0. \end{aligned} \quad (6.37)$$

Thus, $w_1H \subset H$. Similarly, $w_2H \subset H$. $\square$

To prove the sufficient condition of regularity for arbitrary orientation, we need some properties of transforming elements T .

6.5.2. A theorem on transforming elements.

PROPOSITION 6.9 ([St82]). *Let Δ', Δ'' be two arbitrary orientations of the graph Γ that differ by the direction of k edges. Consider the chain of orientations, in which every two adjacent orientations differ by the direction of one edge:*

$$\Delta' = \Delta_0, \Delta_1, \Delta_2, \dots, \Delta_{k-1}, \Delta_k = \Delta''. \quad (6.38)$$

Then, in the Weyl group, there exist elements P_i and S_i , where $i = 1, 2, \dots, k$, such that

$$\begin{aligned} \mathbf{C}_{\Delta_0} &= P_1 S_1, \\ \mathbf{C}_{\Delta_1} &= S_1 P_1 = P_2 S_2, \\ &\vdots \\ \mathbf{C}_{\Delta_{k-1}} &= S_{k-1} P_{k-1} = P_k S_k, \\ \mathbf{C}_{\Delta_k} &= S_k P_k. \end{aligned} \quad (6.39)$$

In addition, for each reflection and each $i = 1, 2, \dots, k$, this reflection does not occur in the decomposition of either P_i or S_i^{-1} .

Proof. It suffices to consider the case $k = 1$. Let us consider the graph

$$\Gamma_1 \cup \Gamma_2 = \Gamma \setminus l.$$

The graph Γ can be depicted as follows:

$$\begin{array}{ccc} \Delta' & \dots & \longleftarrow \dots \\ & \Gamma_1 & \Gamma_2 \\ \Delta'' & \dots & \longrightarrow \dots \\ & \Gamma_1 & \Gamma_2 \end{array} \quad (6.40)$$

The orientations Δ'' and Δ' induce the same orientations on the graphs Γ_1 and Γ_2 , and therefore they induce the same Coxeter transformations on subgraphs Γ_1 and Γ_2 . Denote the corresponding Coxeter transformations by $\mathbf{C}_{\Gamma_1}$ and $\mathbf{C}_{\Gamma_2}$. Then

$$\mathbf{C}_{\Delta'} = \mathbf{C}_{\Gamma_2} \mathbf{C}_{\Gamma_1}, \quad \mathbf{C}_{\Delta''} = \mathbf{C}_{\Gamma_1} \mathbf{C}_{\Gamma_2}.$$

Here,

$$P_1 = \mathbf{C}_{\Gamma_2}, \quad S_1 = \mathbf{C}_{\Gamma_1}. \quad \square$$

REMARK 6.10. Observe, that Proposition 6.9 gives, in particular, a simple proof of the fact that all the Coxeter transformations form one conjugacy class, cf. [Bo, Ch.5, §6].

THEOREM 6.11 ([St82]). 1) *Under the condition of Proposition 6.9,*

$$T^{-1} \mathbf{C}_{\Delta'} T = \mathbf{C}_{\Delta''}$$

for the following $k + 1$ transforming elements $T := T_i$:

$$\begin{aligned}
T_1 &= P_1 P_2 P_3 \dots P_{k-2} P_{k-1} P_k, \\
T_2 &= P_1 P_2 P_3 \dots P_{k-2} P_{k-1} S_k^{-1}, \\
T_3 &= P_1 P_2 P_3 \dots P_{k-2} S_{k-1}^{-1} S_k^{-1}, \\
&\dots \\
T_{k-1} &= P_1 P_2 S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}, \\
T_k &= P_1 S_2^{-1} S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}, \\
T_{k+1} &= S_1^{-1} S_2^{-1} S_3^{-1} \dots S_{k-2}^{-1} S_{k-1}^{-1} S_k^{-1}.
\end{aligned} \tag{6.41}$$

In addition, for each reflection σ_α , there exists a T_i whose decomposition does not contain this reflection.

2) The following relation holds:

$$T_p T_q^{-1} = \mathbf{C}_{\Delta'}^{q-p}. \tag{6.42}$$

Proof. 1) There are altogether 2^k transforming elements of the form

$$T = X_1 X_2 \dots X_{k-1} X_k, \text{ where } X_i \in \{P_i, S_i^{-1}\}. \tag{6.43}$$

By Proposition 6.9, for each reflection σ_α and for each i , we can select $X_i = P_i$ or $X_i = S_i^{-1}$ such that this reflection does not occur in T . Taking the product of all these elements X_i we obtain the transforming element T whose decomposition does not contain this reflection. It remains to show that every transforming element from the list (6.43) containing 2^k elements is of the form (6.41). By (6.39) we have

$$S_q^{-1} P_{q+1} = P_q S_{q+1}^{-1}.$$

Thus, all symbols S_i^{-1} can be shifted to the right and all symbols P_i can be shifted to the left.

2) It suffices to show (6.42) for $p < q$. By (6.41) we have

$$\begin{aligned}
T_p T_q^{-1} &= (P_1 P_2 \dots P_{k-p+1} S_{k-p+2}^{-1} \dots S_k^{-1}) \times (S_k \dots S_{k-q+2} P_1 P_{k-q+1}^{-1} \dots P_1^{-1}) = \\
&P_1 P_2 \dots P_{k-p+1} (S_{k-p+1} S_{k-p} \dots S_{k-q+2}) P_{k-q+1}^{-1} \dots P_1^{-1}.
\end{aligned} \tag{6.44}$$

Here, $k - p + 1 \geq k - q + 2$. In order to simplify (6.44), we use formulas (6.39):

$$\begin{aligned}
P_{k-p} P_{k-p+1} S_{k-p+1} S_{k-p} &= P_{k-p} (S_{k-p} P_{k-p}) S_{k-p} = (P_{k-p} S_{k-p})^2, \\
P_{k-p-1} (P_{k-p} S_{k-p})^2 S_{k-p-1} &= \\
P_{k-p-1} (S_{k-p-1} P_{k-p-1})^2 S_{k-p-1} &= (P_{k-p-1} S_{k-p-1})^3,
\end{aligned} \tag{6.45}$$

$$P_{k-q+2} P_{k-q+3} \dots P_{k-p+1} S_{k-p+1} \dots S_{k-q+2} S_{k-q+2} = (P_{k-q+2} S_{k-q+2})^{q-p}.$$

Thus, from (6.44) and (6.45) we deduce:

$$T_p T_q^{-1} = P_1 P_2 \dots P_{k-q+1} (P_{k-q+2} S_{k-q+2})^{q-p} P_{k-q+1}^{-1} \dots P_1^{-1}. \tag{6.46}$$

Again, by (6.39) we have

$$\begin{aligned}
P_{k-q+1}(P_{k-q+2}S_{k-q+2})^{q-p}P_{k-q+1}^{-1} &= \\
P_{k-q+1}(S_{k-q+1}P_{k-q+1})^{q-p}P_{k-q+1}^{-1} &= (P_{k-q+1}S_{k-q+1})^{q-p}, \\
P_{k-q}(P_{k-q+1}S_{k-q+1})^{q-p}P_{k-q}^{-1} &= \\
P_{k-q}(S_{k-q}P_{k-q})^{q-p}P_{k-q}^{-1} &= (P_{k-q}S_{k-q})^{q-p},
\end{aligned} \tag{6.47}$$

$$P_1(P_2S_2)^{q-p}P_1^{-1} = P_1(S_1P_1)^{q-p}P_1^{-1} = (P_1S_1)^{q-p}.$$

Therefore,

$$T_pT_q^{-1} = (P_1S_1)^{q-p} = \mathbf{C}_{\Delta_0}^{q-p} = \mathbf{C}_{\Delta'}^{q-p}. \quad \square \tag{6.48}$$

REMARK 6.12. Theorem 6.11 allows us to select transforming element T in such a way that its decomposition does not contain any given reflection σ_i , and therefore T does not change any given coordinate i . This fact was already used once in §6.4 for the proof of the necessary conditions of regularity for diagrams with indefinite Tits form. Now, we will use Theorem 6.11 to carry the sufficient condition of regularity from a bicolored orientation Δ in Proposition 6.8 to an arbitrary orientation Δ' , see Definition 6.1.

6.5.3. Sufficient conditions of regularity for an arbitrary orientation.

PROPOSITION 6.13 ([St82]). *Let Γ be an extended Dynkin diagram. Let $\Delta = \Delta_0$, $\Delta' = \Delta_k$. If z is a positive root with zero Δ' -defect, then z is the Δ' -regular vector.*

Proof. If $z > 0$, then there exists a positive coordinate $z_\alpha > 0$. Take transforming element T_i whose decomposition does not contain the reflection σ_α . Then $(T_i z)_\alpha > 0$. Since $T_i z$ is the root, we have $T_i z > 0$. The equality $\rho_{\Delta'}(z) = 0$ means that $\rho_\Delta(T_i z) = 0$, see Definition 6.4. Since $T_i z > 0$ and $T_i z$ is the root, we see by Proposition 6.8 that $T_i z$ is Δ -regular and

$$\mathbf{C}_\Delta^m T_i z > 0 \text{ for all } m \in \mathbb{N}. \tag{6.49}$$

Suppose that

$$u = T_i^{-1} \mathbf{C}_\Delta^m T_i z < 0 \text{ for some } m = m_0.$$

Then by Theorem 6.11 there exists T_j such that $T_j u < 0$, i.e.,

$$T_j T_i^{-1} \mathbf{C}_\Delta^m T_i z < 0 \text{ for } m = m_0. \tag{6.50}$$

Again by Theorem 6.11 we have $T_j T_i^{-1} = \mathbf{C}_\Delta^{i-j}$ and by (6.50) the following relation holds:

$$\mathbf{C}_\Delta^{i-j} \mathbf{C}_\Delta^m T_i z < 0 \text{ for } m = m_0 \tag{6.51}$$

that contradicts to (6.49). Therefore,

$$u = T_i^{-1} \mathbf{C}_\Delta^m T_i z > 0 \text{ for all } m \in \mathbb{N}.$$

Thus, z is a Δ' -regular vector. $\square$

From Theorem 6.5 together with Proposition 6.13 we get the following

THEOREM 6.14. *The indecomposable representation V of the graph Γ (which is an extended Dynkin diagram) with orientation Δ' is regular in the orientation Δ' if and only if*

$$\rho_{\Delta'}(\dim V) = 0.$$

Proof. Indeed, the dimensions of the indecomposable representations of the extended Dynkin diagrams are roots, [DR76], [Kac80]. However, there are indecomposable representations whose dimensions are not usual (real) roots but *imaginary roots*, see §2.4. They are vectors from the kernel of the Tits form, and are proportional to vectors z^1 which are fixed points of the Weyl group. In particular, $Tz^1 = z^1$, so $\langle z^1, \tilde{z}^{1\vee} \rangle = 0$ directly implies $\langle Tz^1, \tilde{z}^{1\vee} \rangle = 0$. $\square$

6.5.4. Invariance of the defect. We will show that the Δ' -defect $\rho_{\Delta'}$ is invariant under the Coxeter transformation $\mathbf{C}_{\Delta'}$ and $\rho_{\Delta'}$ does not depend on the choice of the transforming element T_i in the Weyl group, see Definition 6.4 and Theorem 6.11. In other words, the following proposition holds.

PROPOSITION 6.15. 1) *The Coxeter transformation $\mathbf{C}_{\Delta'}$ preserves the linear form $\rho_{\Delta'}$:*

$$\rho_{\Delta'}(\mathbf{C}_{\Delta'} z) = \rho_{\Delta'}(z) \text{ for any vector } z. \quad (6.52)$$

2) *If T_i and T_j are transforming elements defined by (6.41), i.e.,*

$$T_i^{-1} \mathbf{C}_{\Delta} T_i = \mathbf{C}_{\Delta'} \text{ and } T_j^{-1} \mathbf{C}_{\Delta} T_j = \mathbf{C}_{\Delta'}, \quad (6.53)$$

then we have

$$\langle T_i z, \tilde{z}^{1\vee} \rangle = \langle T_j z, \tilde{z}^{1\vee} \rangle \text{ for any vector } z. \quad (6.54)$$

Proof. First, observe that (6.54) does not hold for an arbitrary matrix T , since the matrix kT , where $k \in \mathbb{R}$, also satisfies (6.53), but does not satisfy (6.54).

In (6.37) we showed that

$$\langle w_i z, \tilde{z}^{1\vee} \rangle = - \langle z, \tilde{z}^{1\vee} \rangle. \quad (6.55)$$

By (6.55) we have

$$\langle z, \tilde{z}^{1\vee} \rangle = \langle \mathbf{C}_{\Delta} z, \tilde{z}^{1\vee} \rangle \text{ for any vector } z, \quad (6.56)$$

i.e., (6.52) holds for any bicolored orientation Δ . Since z in (6.56) is an arbitrary vector, we have

$$\langle Tz, \tilde{z}^{1\vee} \rangle = \langle \mathbf{C}_{\Delta} Tz, \tilde{z}^{1\vee} \rangle = \langle T \mathbf{C}_{\Delta'} z, \tilde{z}^{1\vee} \rangle, \quad (6.57)$$

or

$$\rho_{\Delta'}(z) = \rho_{\Delta'}(\mathbf{C}_{\Delta'} z), \quad (6.58)$$

i.e., (6.52) holds for an arbitrary orientation Δ' .

Further, from (6.56) we have

$$\rho_{\Delta}(z) = \rho_{\Delta}(\mathbf{C}_{\Delta} z) = \rho_{\Delta}(\mathbf{C}_{\Delta}^{i-j} z). \quad (6.59)$$

By (6.42)

$$\rho_{\Delta}(z) = \rho_{\Delta}(T_j T_i^{-1} z). \quad (6.60)$$

Substituting $T_i z$ instead of z we have

$$\rho_{\Delta}(T_i z) = \rho_{\Delta}(T_j z),$$

and (6.54) is proved. $\square$

6.6. Examples of necessary conditions of regularity

DEFINITION 6.16. Let Γ be a graph, $x_0 \in \Gamma$ be a point of maximal branching degree and let all arrows of Γ be directed to the point x_0 . The corresponding orientation is said to be the *central orientation* and denoted by Δ_0 .

We will consider the necessary conditions of regularity for some diagrams in bicolored, central, and other orientations. If an orientation Δ' is obtained from another orientation Δ'' by reversing of all arrows of a graph, then the corresponding conditions of the regularity coincide:

$$\rho_{\Delta'} = \rho_{\Delta''} \quad \text{if } \mathcal{B} > 0 \quad (6.61)$$

or

$$\rho_{\Delta'}^1 = \rho_{\Delta''}^1, \quad \rho_{\Delta'}^2 = \rho_{\Delta''}^2 \quad \text{if } \mathcal{B} \text{ is indefinite.} \quad (6.62)$$

Let us introduce an equivalence relation $\mathcal{R}$ on the set of orientations of the graph Γ . Two orientations Δ' and Δ'' will be called *equivalent*, if one orientation can be obtained from the other one by reversing all arrows or by an automorphism of the diagram; for equivalent orientations, we write:

$$\Delta' \equiv \Delta'' \pmod{\mathcal{R}}.$$

Equivalent orientations have the identical conditions of regularity.




FIGURE 6.1. For $\tilde{D}_4$, the bicolored and central orientations coincide

6.6.1. The three equivalence classes of orientations of $\tilde{D}_4$. In the case of $\tilde{D}_4$, the bicolored orientation Δ coincides with the central orientation Δ_0 . The orientations Δ , Δ' , and Δ'' cover all possible equivalence classes.

a) The bicolored orientation Δ , see Fig. 6.1.

Here, $T = I$ and by Theorem 6.5 and Proposition 6.8 we have

$$z^1 = \begin{pmatrix} 2 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} x_0 \\ y_1 \\ y_2 \\ y_3 \\ y_4 \end{matrix}, \quad \tilde{z}^1 = \begin{pmatrix} 2 \\ -1 \\ -1 \\ -1 \\ -1 \end{pmatrix}, \quad \rho_{\Delta}(z) = y_1 + y_2 + y_3 + y_4 - 2x_0. \quad (6.63)$$

Originally, the linear form $\rho_{\Delta}(z)$ in (6.63) was obtained by I. M. Gelfand and V. A. Ponomarev in the work devoted to classifications of quadruples of linear subspaces of arbitrary dimension [GP72].

b) The orientation Δ' , see Fig. 6.1. Here we have

$$\mathbf{C}_{\Delta} = \sigma_{y_4} \sigma_{y_3} \sigma_{y_2} \sigma_{y_1} \sigma_{x_0}, \quad \mathbf{C}_{\Delta'} = \sigma_{y_4} \sigma_{y_3} \sigma_{y_1} \sigma_{x_0} \sigma_{y_2}, \quad T = \sigma_{y_2}. \quad (6.64)$$

Then, by Theorem 6.5 and Proposition 6.13 we get the following condition of the Δ' -regularity:

$$Tz = \begin{pmatrix} x_0 \\ y_1 \\ x_0 - y_2 \\ y_3 \\ y_4 \end{pmatrix}, \quad \begin{aligned} y_1 + (x_0 - y_2) + y_3 + y_4 - 2x_0 &= 0, \text{ or} \\ y_1 + y_3 + y_4 &= y_2 + x_0. \end{aligned} \quad (6.65)$$

c) The orientation Δ'' , see Fig. 6.1. Here, $T = \sigma_{y_2}\sigma_{y_3}$ (or $\sigma_{y_1}\sigma_{y_4}$), We have the following condition of the Δ'' -regularity:

$$\begin{aligned} y_1 + (x_0 - y_2) + (x_0 - y_3) + y_4 - 2x_0 &= 0, \text{ or} \\ y_1 + y_4 &= y_2 + y_3. \end{aligned} \quad (6.66)$$




FIGURE 6.2. For $\tilde{E}_6$, the bicolored and central orientations

6.6.2. The bicolored and central orientations of $\tilde{E}_6$. We consider only the bicolored orientation Δ and the central orientation Δ_0 , Fig. 6.2. The Coxeter transformations and transforming element T are:

$$\begin{aligned} C_\Delta &= \sigma_{y_3}\sigma_{y_2}\sigma_{y_1}\sigma_{x_3}\sigma_{x_2}\sigma_{x_1}\sigma_{x_0}, \\ C_{\Delta_0} &= \sigma_{x_3}\sigma_{x_2}\sigma_{x_1}\sigma_{y_3}\sigma_{y_2}\sigma_{y_1}\sigma_{x_0}, \\ T &= \sigma_{x_3}\sigma_{x_2}\sigma_{x_1}. \end{aligned} \quad (6.67)$$

We have

$$z^1 = \begin{pmatrix} 3 \\ 1 \\ 1 \\ 1 \\ 2 \\ 2 \\ 2 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \\ x_3 \\ y_1 \\ y_2 \\ y_3 \end{pmatrix}, \quad \tilde{z}^1 = \begin{pmatrix} 3 \\ 1 \\ 1 \\ 1 \\ -2 \\ -2 \\ -2 \end{pmatrix}, \quad Tz = \begin{pmatrix} x_0 \\ y_1 - x_1 \\ y_2 - x_2 \\ y_3 - x_3 \\ y_1 \\ y_2 \\ y_3 \end{pmatrix}, \quad (6.68)$$

and by Theorem 6.5 and Proposition 6.8 a condition of Δ -regularity is

$$\rho_\Delta(z) = 3x_0 + x_1 + x_2 + x_3 + 3x_0 - 2y_1 - 2y_2 - 2y_3. \quad (6.69)$$

The choice of element T is ambiguous, but it was shown in Proposition 6.15 that the regularity condition does not depend on this choice.

So, a condition of Δ_0 -regularity is

$$\begin{aligned} \rho_{\Delta_0}(z) &= \\ 3x_0 + (y_1 - x_1) + (y_2 - x_2) + (y_3 - x_3) - 2(y_1 + y_2 + y_3) &= 0, \end{aligned}$$

or

$$\rho_{\Delta_0}(z) = x_1 + x_2 + x_3 + y_1 + y_2 + y_3 - 3x_0. \quad (6.70)$$




FIGURE 6.3. The two orientations of $\tilde{G}_{21}$ and $\tilde{G}_{22} = \tilde{G}_{21}^V$

6.6.3. The multiply-laced case. The two orientations of $\tilde{G}_{21}$ and $\tilde{G}_{22} = \tilde{G}_{21}^V$. We have

$$z^1 = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} \begin{matrix} x_1 \\ y_1 \\ y_2 \end{matrix}, \quad \tilde{z}^1 = \begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix}, \quad z^{1V} = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} x_1 \\ y_1 \\ y_2 \end{matrix}, \quad \tilde{z}^{1V} = \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}. \quad (6.71)$$

Take vectors $\tilde{z}^1$ and $\tilde{z}^{1V}$ from (6.71) and substitute them in (6.13). We get the following regularity condition ρ_{Δ} (resp. ρ_{Δ}^V):

$$\begin{aligned} \rho_{\Delta}(z) &= y_1 + y_2 - 2x_1 \quad \text{for } \tilde{G}_{21}, \\ \rho_{\Delta}^V(z) &= y_1 + 3y_2 - 2x_1 \quad \text{for } \tilde{G}_{22}. \end{aligned} \quad (6.72)$$

Now consider the orientation Δ' , Fig. 6.3. The Coxeter transformations can be expressed as follows:

$$\begin{aligned} \mathbf{C}_{\Delta} &= \sigma_{y_2} \sigma_{y_1} \sigma_{x_1}, & \mathbf{C}_{\Delta}^V &= \sigma_{y_2}^V \sigma_{y_1}^V \sigma_{x_1}^V, \\ \mathbf{C}_{\Delta'} &= \sigma_{y_1} \sigma_{x_1} \sigma_{y_2}, & \mathbf{C}_{\Delta'}^V &= \sigma_{y_1}^V \sigma_{x_1}^V \sigma_{y_2}^V, \end{aligned} \quad (6.73)$$

Transforming elements T and T^V are:


$$T = \sigma_{y_2}, \quad T^V = \sigma_{y_2}^V. \quad (6.74)$$

and

$$Tz = \begin{pmatrix} x_1 \\ y_1 \\ 3x_1 - y_2 \end{pmatrix}, \quad T^V z = \begin{pmatrix} x_1 \\ y_1 \\ x_1 - y_2 \end{pmatrix}. \quad (6.75)$$

Finally, we have

$$\begin{aligned} \rho_{\Delta'}(z) &= y_1 - y_2 + x_1 \quad \text{for } \tilde{G}_{21}, \\ \rho_{\Delta'}^V(z) &= y_1 - 3y_2 + x_1 \quad \text{for } \tilde{G}_{22}. \end{aligned} \quad (6.76)$$

FIGURE 6.4. A bicolored orientation of the star $*_{n+1}$

6.6.4. The case of indefinite $\mathcal{B}$. The oriented star $*_{n+1}$. Consider the oriented star $*_{n+1}$ with a bicolored orientation.

According to Remark 3.6 the matrix DD^t is a scalar. By (3.21) $DD^t = \frac{n}{4}$ and the maximal eigenvalue $\varphi^m = \frac{n}{4}$. By (3.20) we have

$$\lambda_{1,2}^m = \frac{n-2 \pm \sqrt{n(n-4)}}{2}. \quad (6.77)$$

Let $x_m = 1$. Then

$$\frac{-2}{\lambda_{1,2}^m + 1} D^t x^m = \frac{-2}{\lambda_{1,2}^m + 1} [1, 1, \dots, 1]^t.$$

Thus, we have the following eigenvectors z_1^m, z_2^m :

$$z_1^m = \begin{pmatrix} \lambda_1^m + 1 \\ 1 \\ 1 \\ \dots \\ 1 \end{pmatrix}, \quad z_2^m = \begin{pmatrix} \lambda_2^m + 1 \\ 1 \\ 1 \\ \dots \\ 1 \end{pmatrix},$$

and their conjugate vectors $\tilde{z}_1^m, \tilde{z}_2^m$ (see Definition 6.2 and Proposition 6.3):

$$\tilde{z}_1^m = \begin{pmatrix} \lambda_2^m + 1 \\ -1 \\ -1 \\ \dots \\ -1 \end{pmatrix}, \quad \tilde{z}_2^m = \begin{pmatrix} \lambda_1^m + 1 \\ -1 \\ -1 \\ \dots \\ -1 \end{pmatrix}.$$

According to (6.32) we obtain the following condition of Δ -regularity:

$$(\lambda_2^m + 1)x_0 - \sum y_i \leq 0, \quad (\lambda_1^m + 1)x_0 - \sum y_i \geq 0. \quad (6.78)$$

From (6.77) and (6.78) we deduce

$$\begin{aligned}
& \frac{n - \sqrt{n(n-4)}}{2} x_0 \leq \sum y_i \leq \frac{n + \sqrt{n(n-4)}}{2} x_0 \text{ or} \\
& \frac{n - \sqrt{n(n-4)}}{2} \sum y_i \leq x_0 \leq \frac{n + \sqrt{n(n-4)}}{2} \sum y_i, \text{ or} \\
& |x_0 - \frac{1}{2} \sum y_i| \leq \frac{\sqrt{n(n-4)}}{2n} \sum y_i \text{ or} \\
& x_0^2 - x_0 \sum y_i + \frac{1}{4} (\sum y_i)^2 \leq \frac{1}{4} (\sum y_i)^2 - \frac{1}{n} (\sum y_i)^2 \text{ or} \\
& x_0^2 - x_0 \sum y_i \leq -\frac{1}{n} (\sum y_i)^2 \text{ or} \\
& x_0^2 - x_0 \sum y_i + \sum y_i^2 \leq \sum y_i^2 - \frac{1}{n} (\sum y_i)^2
\end{aligned} \tag{6.79}$$

Since the left hand side of the latter inequality is the Tits form $\mathcal{B}$, we obtain the following condition of Δ -regularity:

$$\mathcal{B}(z) \leq \frac{1}{n} \sum_{0 < i < j} (y_i - y_j)^2. \tag{6.80}$$

APPENDIX A

The McKay correspondence and the Slodowy generalization

We have seen during the past few years a major assault on the problem of determining all the finite simple groups. ... If I am right, I foresee new proofs of classification which will owe little or nothing to the current proofs. They will be much shorter and will help us to understand the simple groups in a context much wider than finite group theory.

J. McKay, [McK80, p.183], 1980

A.1. Finite subgroups of $SU(2)$ and $SO(3, \mathbb{R})$

Let us consider the special unitary group $SU(2)$, the subgroup of unitary transformations in $GL(2, \mathbb{C})$ with determinant 1 and its quotient group $PSU(2) = SU(2)/\{\pm 1\}$ acting on the complex projective line $\mathbb{CP}^1$, see §B.4. The projective line $\mathbb{CP}^1$ can be identified with the sphere $S^2 \subset \mathbb{R}^3$, see §B.4 and (B.16).

The transformations of the sphere S^2 induced by elements of $SO(3, \mathbb{R})$ (orientation preserving rotations of $\mathbb{R}^3$) correspond under these identifications to transformations of $\mathbb{CP}^1$ in the group $PSU(2)$:

$$PSU(2) \cong SO(3, \mathbb{R}), \tag{A.1}$$

see [Sp77, Prop.4.4.3], [PV94, §0.13].

It is well known that the finite subgroups of $SO(3, \mathbb{R})$ are precisely the rotation groups of the following polyhedra: the regular n -angled pyramid, the n -angled dihedron (a regular plane n -gon with two faces), the tetrahedron, the cube (its the rotation group coincides with the rotation group of the octahedron), the icosahedron (its the rotation group coincides with the rotation group of the dodecahedron). By (A.1) we have a classification of the finite subgroups of $PSU(2)$, see Table A.1.

To get a classification of all finite subgroups of $SU(2)$ we make use of the double covering

$$\pi : SU(2) \longrightarrow SO(3, \mathbb{R}).$$

If G is a finite subgroup of $SO(3, \mathbb{R})$, we see that the preimage $\pi^{-1}(G)$ is a finite subgroup of $SU(2)$ and $|\pi^{-1}(G)| = 2|G|$. The finite subgroups of $SO(3, \mathbb{R})$ are called *polyhedral groups*, see Table A.1. The finite subgroups of $SU(2)$ are naturally called *binary polyhedral groups*, see Table A.2.

REMARK A.1. 1) The cyclic group is an exceptional case. The preimage of the cyclic group $G = \mathbb{Z}/n\mathbb{Z}$ is the even cyclic group $\mathbb{Z}/2n\mathbb{Z}$, which is a binary cyclic group. The cyclic group $\mathbb{Z}/(2n-1)\mathbb{Z}$ is not a preimage with respect to π , so $\mathbb{Z}/(2n-1)\mathbb{Z}$ is not a binary polyhedral group. Anyway, the cyclic groups $\mathbb{Z}/n\mathbb{Z}$ complete the list of finite subgroups of $SU(2)$.

TABLE A.1. The polyhedral groups in $\mathbb{R}^3$

Polyhedron	Orders of symmetries	Rotation group	Group order
Pyramid	—	cyclic	n
Dihedron	$n \ 2 \ 2$	dihedral	$2n$
Tetrahedron	$3 \ 2 \ 3$	$\mathcal{A}_4$	12
Cube	$4 \ 2 \ 3$	$\mathcal{S}_4$	24
Octahedron	$3 \ 2 \ 4$	$\mathcal{S}_4$	24
Dodecahedron	$5 \ 2 \ 3$	$\mathcal{A}_5$	60
Icosahedron	$3 \ 2 \ 5$	$\mathcal{A}_5$	60

Here, $\mathcal{S}_m$ (resp. $\mathcal{A}_m$) denotes the symmetric, (resp. alternating) group of all (resp. of all even) permutations of m letters.

2) Every finite subgroup of $SL(2, \mathbb{C})$ is conjugate to a subgroup in $SU(2)$, so we also have a classification of finite subgroups of $SL(2, \mathbb{C})$, [Sp77].

For every polyhedral group G , the axis of rotations under an element $\gamma \in G$ passes through either the mid-point of a face, or the mid-point of an edge, or a vertex. We denote the orders of symmetry of these axes by p , q , and r , respectively, see [Sl83]. These numbers are listed in Table A.1. The triples p, q, r listed in Table A.1 are exactly the solutions of the diophantine inequality

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1, \quad (\text{A.2})$$

see [Sp77].

A.2. Generators and relations in polyhedral groups

The quaternion group introduced by W. R. Hamilton is defined as follows: It is generated by three generators i , j , and k subject to the relations

$$i^2 = j^2 = k^2 = ijk = -1; \quad (\text{A.3})$$

for references, see [Cox40], [CoxM84]. A natural generalization of the quaternion group is the group generated by three generators R , S , and T subject to the relations

$$R^p = S^q = T^r = RST = -1. \quad (\text{A.4})$$

Denote by $\langle p, q, r \rangle$ the group defined by (A.4).

W. Threlfall, (see [CoxM84]) has observed that

$$\langle 2, 2, n \rangle, \langle 2, 3, 3 \rangle, \langle 2, 3, 4 \rangle, \langle 2, 3, 5 \rangle$$

are the *binary polyhedral groups*, of order

$$\frac{4}{\frac{1}{p} + \frac{1}{q} + \frac{1}{r} - 1}$$

TABLE A.2. The finite subgroups of $SU(2)$

$\langle l, m, n \rangle$	Order	Denotation	Well-known name
$-$	n	$\mathbb{Z}/n\mathbb{Z}$	cyclic group
$\langle 2, 2, n \rangle$	$4n$	$\mathcal{D}_n$	binary dihedral group
$\langle 2, 3, 3 \rangle$	24	$\mathcal{T}$	binary tetrahedral group
$\langle 2, 3, 4 \rangle$	48	$\mathcal{O}$	binary octahedral group
$\langle 2, 3, 5 \rangle$	120	$\mathcal{I}$	binary icosahedral group

H. S. M. Coxeter proved in [Cox40, p.370] that having added one more generator Z one can replace (A.4) by the relations:

$$R^p = S^q = T^r = RST = Z. \quad (\text{A.5})$$

For the groups $\langle 2, 2, n \rangle$ and $\langle 2, 3, n \rangle$, where $n = 3, 4, 5$, relation (A.5) implies

$$Z^2 = 1.$$

The polyhedral groups from Table A.1 are described by generations and relations as follows:

$$R^p = S^q = T^r = RST = 1, \quad (\text{A.6})$$

see [Cox40], [CoxM84].

A.3. Kleinian singularities and the Du Val resolution

Consider the quotient variety $\mathbb{C}^2/G$, where G is a binary polyhedral group $G \subset SU(2)$ from Table A.2. According to (§B.5.4 and (B.40)) $X = \mathbb{C}^2/G$ is the orbit space given by the prime spectrum on the algebra of invariants R^G :

$$X := \text{Spec}(R^G), \quad (\text{A.7})$$

where $R = \mathbb{C}[z_1, z_2]$ (see eq.(2.27)) which coincides with the symmetric algebra $\text{Sym}((\mathbb{C}^2)^*)$ (see §2.7.1; for more details, see [Sp77], [PV94], [Sl80].)

F. Klein [Kl1884] observed that the algebra of invariants $\mathbb{C}[z_1, z_2]^G$ for every binary polyhedral group $G \subset SU(2)$ from Table A.2 can be considered by one approach:

THEOREM A.2 (F. Klein, [Kl1884]). *The algebra of invariants $\mathbb{C}[z_1, z_2]^G$ is generated by 3 variables x, y, z , subject to one essential relation*

$$R(x, y, z) = 0, \quad (\text{A.8})$$

where $R(x, y, z)$ is defined in column 2, Table A.3. In other words, the algebra of invariants $\mathbb{C}[z_1, z_2]^G$ coincides with the coordinate algebra (see §B.5.3) of a curve defined by the eq. (A.8), i.e.,

$$\mathbb{C}[z_1, z_2]^G \simeq \mathbb{C}[x, y, z]/(R(x, y, z)). \quad (\text{A.9})$$

TABLE A.3. The relations $R(x, y, z)$ describing the algebra of invariants $\mathbb{C}[z_1, z_2]^G$

Finite subgroup of $SU(2)$	Relation $R(x, y, z)$	Dynkin diagram
$\mathbb{Z}/n\mathbb{Z}$	$x^n + yz$	A_{n-1}
$\mathcal{D}_n$	$x^{n+1} + xy^2 + z^2$	D_{n+2}
$\mathcal{T}$	$x^4 + y^3 + z^2$	E_6
$\mathcal{O}$	$x^3y + y^3 + z^2$	E_7
$\mathcal{J}$	$x^5 + y^3 + z^2$	E_8

The quotient X from (A.7) has no singularity except at the origin $O \in \mathbb{C}^3$. The quotient variety X is called a *Kleinian singularity* also known as a *Du Val singularity*, a *simple surface singularity* or a *rational double point*.

See Remark A.3, [SI83, §5]. The quotient variety X can be embedded as a surface $X \subset \mathbb{C}^3$ with an isolated singularity at the origin.

REMARK A.3. According to Theorem 2.3 (Sheppard-Todd-Chevalley-Serre) the algebra of invariants $k[V]^G$ is isomorphic to a polynomial algebra in some number of variables if the image of G in $GL(V)$ is generated by reflections. Every binary polyhedral group G is generated by reflections, see, e.g., [CoxM84], therefore the algebra of invariants $k[V]^G$ for the binary polyhedral group is a polynomial algebra.

EXAMPLE A.4. Consider the cyclic group $G = \mathbb{Z}/r\mathbb{Z}$ of the order r . The group G acts on $\mathbb{C}[z_1, z_2]$ as follows:

$$(z_1, z_2) \mapsto (\varepsilon z_1, \varepsilon^{r-1} z_2), \quad (\text{A.10})$$

where $\varepsilon = e^{2\pi i/r}$, and polynomials

$$x = z_1 z_2, \quad y = -z_1^r, \quad z = z_2^r \quad (\text{A.11})$$

are invariant polynomials in $\mathbb{C}[x, y, z]$ which satisfy the following relation

$$x^r + yz = 0, \quad (\text{A.12})$$

see Table A.3. We have

$$k[V]^G = \mathbb{C}[z_1 z_2, z_1^r, z_2^r] \simeq \mathbb{C}[x, y, z]/(x^r + yz).$$

see, e.g., [Sp77, pp.95-97], [PV94, p.143].

Du Val obtained the following description of the minimal resolution $\pi : \tilde{X} \rightarrow X$ of a Kleinian singularity $X = \mathbb{C}^2/G$, see [DuVal34], [Sl80, §6.1, §6.2] [Sl83, §5]¹. The *exceptional divisor* (the preimage of the singular point O) is a finite union of complex projective lines:

$$\pi^{-1}(O) = L_1 \cup \cdots \cup L_n, \quad L_i \simeq \mathbb{CP}^1 \text{ for } i = 1, \dots, n.$$

The intersection $L_i \cap L_j$ is empty or consists of exactly one point for $i \neq j$.

To each complex projective line L_i (which can be identified with the sphere $S^2 \subset \mathbb{R}^3$, see §B.4) we assign a vertex i , and two vertices are connected by an edge if the corresponding projective lines intersect. The corresponding diagrams are Dynkin diagrams (this phenomenon was observed by Du Val in [DuVal34]), see Table A.3.

In the case of the binary dihedral group $\mathcal{D}_2$ the real resolution of the real variety

$$\mathbb{C}^3/R(x, y, z) \cap \mathbb{R}^3$$

gives a quite faithful picture of the complex situation, the minimal resolution $\pi^{-1} : \tilde{X} \rightarrow X$ for $X = \mathcal{D}_2$ depicted on Fig. A.1. Here $\pi^{-1}(O)$ consists of four circles, the corresponding diagram is the Dynkin diagram D_4 .




FIGURE A.1. The minimal resolution $\pi^{-1} : \tilde{X} \rightarrow X$ for $X = \mathcal{D}_2$

A.4. The McKay correspondence

Let G be a finite subgroup of $SU(2)$. Let $\{\rho_0, \rho_1, \dots, \rho_n\}$ be the set of irreducible finite dimensional complex representations of G , of which ρ_0 is the trivial one. Let $\rho : \rightarrow SU(2)$ be a faithful representation. Then, for each group G , we can define a matrix $A(G) = (a_{ij})$, by decomposing the tensor products:

$$\rho \otimes \rho_j = \bigoplus_{k=0}^r a_{jk} \rho_k, \quad j, k = 0, 1, \dots, r, \quad (\text{A.13})$$

where a_{jk} is the multiplicity of ρ_k in $\rho \otimes \rho_j$. McKay [McK80] observed that

¹For more details, see also [Gb02], [Rie02], [Hob02], [Cr01].

The matrix $2I - A(G)$ is the Cartan matrix of the extended Dynkin diagram $\tilde{\Delta}(G)$ associated to G . There is one-to-one correspondence between finite subgroups of $SU(2)$ and simply-laced extended Dynkin diagrams.

This remarkable observation, called the *McKay correspondence*, was based first on an explicit verification [McK80].

For the multiply-laced case, the McKay correspondence was extended by D. Happel, U. Preiser, and C. M. Ringel in [HPR80], and by P. Slodowy in [Sl80, App.III]. We consider P. Slodowy's approach in §A.5.

The systematic proof of the McKay correspondence based on the study of affine Coxeter transformations was given by R. Steinberg in [Stb85].

Other explanations of the McKay correspondence have been given by G. Gonzalez-Sprinberg and J.-L. Verdier in [GV83], by H. Knörrer in [Kn85]. A nice review is given by J. van Hoboken in [Hob02].

B. Kostant used the *McKay matrix* $A(G)$ (or *McKay operator*) in [Kos84] and showed that the multiplicities $m_i(n)$ in the decomposition

$$\pi_n|G = \sum_{i=0}^r m_i(n) \rho_i,$$

(see §1.5.2) come in an amazing way from the orbit structure of the Coxeter transformation on the highest root of the corresponding Lie algebra $\mathfrak{g}$, see §B.6. More explicitly, the multiplicities $m_i(n)$ come from the orbit structure of the affine Coxeter transformation on the set of roots of $\mathfrak{g}$. To calculate these multiplicities, Kostant employed generating functions and Poincaré series. We applied Kostant's technique in §5.4 in order to show a relation between Poincaré series and a ratio of characteristic polynomials of the Coxeter transformations.

A.5. The Slodowy generalization of the McKay correspondence

We consider here the Slodowy generalization [Sl80] of the McKay correspondence to the multiply-laced case and illustrate Slodowy's approach with the diagrams $\tilde{F}_{41}$ and $\tilde{F}_{42}$, see Fig. 2.2.

Slodowy's approach is based on the consideration of *restricted representations* and *induced representations* instead of an original representation. Let $\rho : G \rightarrow GL(V)$ be a representation of a group G . We denote the *restricted representation* of ρ to a subgroup $H \subset G$ by $\rho \downarrow_H^G$, or, briefly, $\rho^\downarrow$ for fixed G and H . Let $\tau : H \rightarrow GL(V)$ be a representation of a subgroup H . We denote by $\tau \uparrow_H^G$ the representation *induced* by τ to a representation of the group G containing H ; we briefly write $\tau^\uparrow$ for fixed G and H . For a detailed definition on restricted and induced representations, see, for example, [Kar92] or [Bak04].

Let us consider pairs of groups $H \triangleleft G$, where H and G are binary polyhedral groups from Tables A.2 and A.3.

A.5.1. The Slodowy correspondence. Let us fix a pair $H \triangleleft G$ from Table A.4. We formulate now the essence of the Slodowy correspondence [Sl80, App.III].

1) Let ρ_i , where $i = 1, \dots, n$, be irreducible representations of G ; let $\rho_i^\downarrow$ be the corresponding restricted representations of the subgroup H . Let ρ be a faithful representation of H , which may be considered as the restriction of the fixed faithful representation ρ_f of G . Then the following decomposition formula makes sense

$$\rho \otimes \rho_i^\downarrow = \bigoplus a_{ji} \rho_j^\downarrow \tag{A.14}$$

TABLE A.4. The pairs $H \triangleleft G$ of binary polyhedral groups

Subgroup H	Dynkin diagram $\Gamma(H)$	Group R	Dynkin diagram $\Gamma(G)$	Index $[G : H]$
$\mathcal{D}_2$	D_4	$\mathcal{T}$	E_6	3
$\mathcal{T}$	E_6	$\mathcal{O}$	E_7	2
$\mathcal{D}_{r-1}$	D_{n+1}	$\mathcal{D}_{2(r-1)}$	D_{2n}	2
$\mathbb{Z}/2r\mathbb{Z}$	A_{n-1}	$\mathcal{D}_r$	D_{r+2}	2

and uniquely determines an $n \times n$ matrix $\tilde{A} = (a_{ij})$ such that

$$K = 2I - \tilde{A} \quad (\text{A.15})$$

(see [S180, p.163]), where K is the Cartan matrix of the corresponding folded extended Dynkin diagram given in Table A.5

2) Let τ_i , where $i = 1, \dots, n$, be irreducible representations of the subgroup H , let $\tau_i^\uparrow$ be the induced representations of the group G . Then the following decomposition formula makes sense

$$\rho \otimes \tau_i^\uparrow = \bigoplus a_{ij} \tau_j^\uparrow, \quad (\text{A.16})$$

i.e., the decomposition of the induced representation is described by the matrix $A^\vee = A^t$ which satisfies the relation

$$K^\vee = 2I - \tilde{A}^\vee \quad (\text{A.17})$$

(see [S180, p.164]), where $K^\vee$ is the Cartan matrix of the dual folded extended Dynkin diagram given in Table A.5.

We call matrices $\tilde{A}$ and $\tilde{A}^\vee$ the *Slodowy matrices*, they are analogs of the McKay matrix. The *Slodowy correspondence* is an analogue to the McKay correspondence for the multiply-laced case, so one can speak about the *McKay-Slodowy correspondence*.

A.5.2. The binary tetrahedral group and the binary octahedral group. Now we will illustrate the Slodowy correspondence for the binary tetrahedral and octahedral groups, i.e., H is the binary tetrahedral group $\mathcal{T}$ and G is the binary octahedral group $\mathcal{O}$, $\mathcal{T} \triangleleft \mathcal{O}$. These groups have orders $|\mathcal{T}| = 24$ and $|\mathcal{O}| = 48$, see Table A.2.

TABLE A.5. The pairs $H \triangleleft G$ and folded extended Dynkin diagrams

Groups $H \triangleleft G$	Dynkin diagram $\tilde{\Gamma}(H)$ and $\tilde{\Gamma}(G)$	Folded extended Dynkin diagram
$\mathcal{D}_2 \triangleleft \mathcal{T}$	D_4 and E_6	$\tilde{G}_{21}$ and $\tilde{G}_{22}$
$\mathcal{T} \triangleleft \mathcal{O}$	E_6 and E_7	$\tilde{F}_{41}$ and $\tilde{G}_{42}$
$\mathcal{D}_{r-1} \triangleleft \mathcal{D}_{2(r-1)}$	D_{n+1} and D_{2n}	$\widetilde{DD}_n$ and $\widetilde{CD}_n$
$\mathbb{Z}/2r\mathbb{Z} \triangleleft \mathcal{D}_r$	A_{n-1} and D_{r+2}	$\tilde{B}_n$ and $\tilde{C}_n$

We will use the Springer formula for elements of the group $\mathcal{O}$ from [Sp77, §4.4.11]. Let

$$a = \begin{pmatrix} \varepsilon & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix}, \quad b = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad c = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{-1} & \varepsilon^{-1} \\ -\varepsilon & \varepsilon \end{pmatrix}, \quad (\text{A.18})$$

where $\varepsilon = e^{\pi i/4}$. Then each of the 48 different elements $x \in \mathcal{O}$ may be expressed as follows:

$$x = a^h b^j c^l, \quad 0 \leq h < 8, \quad 0 \leq j < 2, \quad 0 \leq l < 3. \quad (\text{A.19})$$

The elements $x \in \mathcal{O}$ and their traces are collected in Table A.6. Observe that every element $u \in SL(2, \mathbb{C})$ from Table A.6 is of the form

$$u = \begin{pmatrix} \alpha & \beta \\ -\beta^* & \alpha^* \end{pmatrix}, \quad (\text{A.20})$$

see (5.63). We can now distinguish the elements of $\mathcal{O}$ by their traces and by means of the 1-dimensional representation ρ_1 such that

$$\rho_1(a) = -1, \quad \rho_1(b) = 1, \quad \rho_1(c) = 1. \quad (\text{A.21})$$

PROPOSITION A.5. *There are 8 conjugacy classes in the binary octahedral group $\mathcal{O}$. Rows of Table A.7 constitute these conjugacy classes.*

Proof. In order to prove the proposition we need a number of conjugacy relations.

1) It is easy to verify that

$$b \begin{pmatrix} x & y \\ u & v \end{pmatrix} b^{-1} = \begin{pmatrix} v & u \\ y & x \end{pmatrix} \quad \text{for all } x, y, u, v \in \mathbb{C}, \quad (\text{A.22})$$

TABLE A.6. The elements of the binary octahedral group

Elements $x = a^p b^j c^l$ $0 \leq p < 8$	Matrix form u	Trace $tr(x)$
a^p	$\begin{pmatrix} \varepsilon^p & 0 \\ 0 & \varepsilon^{-p} \end{pmatrix}$	$2 \cos \frac{\pi p}{4}$
$a^p b$	$\begin{pmatrix} 0 & i\varepsilon^p \\ i\varepsilon^{-p} & 0 \end{pmatrix}$	0
$a^p c$	$\frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{p-1} & \varepsilon^{p-1} \\ -\varepsilon^{-(p-1)} & \varepsilon^{-(p-1)} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{p-1} & \varepsilon^{p-1} \\ \varepsilon^{-p+5} & \varepsilon^{-p+1} \end{pmatrix}$	$\sqrt{2} \cos \frac{\pi(p-1)}{4}$
$a^p bc$	$\frac{1}{\sqrt{2}} \begin{pmatrix} -i\varepsilon^{p+1} & i\varepsilon^{p+1} \\ i\varepsilon^{-(p+1)} & i\varepsilon^{-(p+1)} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{p-1} & \varepsilon^{p+3} \\ \varepsilon^{-p+1} & \varepsilon^{-p+1} \end{pmatrix}$	$\sqrt{2} \cos \frac{\pi(p-1)}{4}$
$a^p c^2$	$\frac{1}{\sqrt{2}} \begin{pmatrix} -\varepsilon^{p+1} & \varepsilon^{p-1} \\ -\varepsilon^{-(p-1)} & -\varepsilon^{-(p+1)} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{p+5} & \varepsilon^{p-1} \\ \varepsilon^{-p+5} & \varepsilon^{-p+3} \end{pmatrix}$	$-\sqrt{2} \cos \frac{\pi(p+1)}{4}$
$a^p bc^2$	$\frac{1}{\sqrt{2}} \begin{pmatrix} -i\varepsilon^{p+1} & -i\varepsilon^{p-1} \\ -i\varepsilon^{-(p-1)} & i\varepsilon^{-(p+1)} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{p-1} & \varepsilon^{p-3} \\ \varepsilon^{-p-1} & \varepsilon^{-p+1} \end{pmatrix}$	$\sqrt{2} \cos \frac{\pi(p-1)}{4}$

and

$$a \begin{pmatrix} x & y \\ u & v \end{pmatrix} a^{-1} = \begin{pmatrix} x & \varepsilon^2 y \\ \varepsilon^2 u & v \end{pmatrix} \text{ for all } x, y, u, v \in \mathbb{C}. \quad (\text{A.23})$$

TABLE A.7. The conjugacy classes in the binary octahedral group

Trace $tr(x)$	Representation ρ_1	Conjugacy class $Cl(g)$	Class order	Representative g
2	1	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	1	1
-2	1	$b^2 = c^3 = a^4 = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	1	-1
0	-1	$ab, a^3b, a^5b, a^7b,$ $a^3bc, a^7bc, ac^2, a^5c^2,$ $a^3c, a^7c, a^3bc^2, a^7bc^2$	12	ab
0	1	$a^2, a^6, b, a^2b, a^4b, a^6b$	6	b
-1	1	$a^4c, a^6c, a^4bc, a^6bc,$ $c^2, a^6c^2, a^4bc^2, a^6bc^2$	8	c^2
1	1	$c, a^2c, bc, a^2bc,$ $a^4c^2, a^2c^2, bc^2, a^2bc^2$	8	c
$\sqrt{2}$	-1	$a, a^7, ac, abc, a^3c^2, abc^2$	6	a
$-\sqrt{2}$	-1	$a^3, a^5, a^5c, a^5bc, a^7c^2, a^5bc^2$	6	a^3

From Table A.6 and (A.22) and since $a^p = a^{p+8}$, we see, for all $p \in \mathbb{Z}$, that

$$b(a^p)b^{-1} = a^{-p}, \quad (\text{A.24})$$

$$b(a^p c)b^{-1} = a^{2-p}bc, \quad (\text{A.25})$$

¹See Remark A.6.

$$b(a^p b c^2) b^{-1} = a^{2-p} b c^2, \quad (\text{A.26})$$

$$b(a^p c^2) b^{-1} = a^{6-p} c^2. \quad (\text{A.27})$$

From Table A.6 and (A.23), we see, for all $p \in \mathbb{Z}$, that

$$a(a^p c^2) a^{-1} = a^{p-2} b c. \quad (\text{A.28})$$

By (A.24)

$$a \sim a^7, \quad a^2 \sim a^6, \quad a^3 \sim a^5. \quad (\text{A.29})$$

By (A.25)

$$\begin{aligned} c &\sim a^2 b c, \quad a c \sim a b c, \quad a^2 c \sim b c, \quad a^3 c \sim a^7 b c, \\ a^4 c &\sim a^6 b c, \quad a^5 c \sim a^5 b c, \quad a^6 c \sim a^4 b c, \quad a^7 c \sim a^3 b c. \end{aligned} \quad (\text{A.30})$$

By (A.26)

$$b c^2 \sim a^2 b c^2, \quad a^3 b c^2 \sim a^7 b c^2, \quad a^4 b c^2 \sim a^6 b c^2. \quad (\text{A.31})$$

By (A.27)

$$c^2 \sim a^6 c^2, \quad a c^2 \sim a^5 c^2, \quad a^2 c^2 \sim a^4 c^2. \quad (\text{A.32})$$

By (A.28)

$$\begin{aligned} c^2 &\sim a^6 b c, \quad a c^2 \sim a^7 b c, \quad a^2 c^2 \sim b c, \quad a^3 c^2 \sim a b c, \\ a^4 c^2 &\sim a^2 b c, \quad a^5 c^2 \sim a^3 b c, \quad a^6 c^2 \sim a^4 b c, \quad a^7 c^2 \sim a^5 b c. \end{aligned} \quad (\text{A.33})$$

From (A.24) we have

$$b a^{-1} = a b, \quad b a^{-p} = a^p b. \quad (\text{A.34})$$

By (A.24) $a^p b \sim a a^p b a^{-1} = a a^p a b$, i.e.,

$$a^p b \sim a^{p+2} b, \quad (\text{A.35})$$

i.e.,

$$\begin{aligned} b &\sim a^2 b \sim a^4 b \sim a^6 b, \\ a b &\sim a^3 b \sim a^5 b \sim a^7 b. \end{aligned} \quad (\text{A.36})$$

2) We need some conjugacy relations performed by c . By (5.64) the inverse elements in $SL(2, \mathbb{C})$ are

$$u^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \text{ for every } u = \begin{pmatrix} a & b \\ c & d \end{pmatrix}. \quad (\text{A.37})$$

So, we have

$$c = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon^{-1} & \varepsilon^{-1} \\ -\varepsilon & \varepsilon \end{pmatrix} \text{ and } c^{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} \varepsilon & -\varepsilon^{-1} \\ \varepsilon & \varepsilon^{-1} \end{pmatrix}. \quad (\text{A.38})$$

Thus

$$c a^p c^{-1} = \frac{1}{2} \begin{pmatrix} \varepsilon^p + \varepsilon^{-p} & -\varepsilon^{p-2} + \varepsilon^{-p-2} \\ -\varepsilon^{p+2} + \varepsilon^{-p+2} & \varepsilon^p + \varepsilon^{-p} \end{pmatrix}. \quad (\text{A.39})$$

Since

$$\varepsilon^2 = i, \quad \varepsilon^2 = -i, \quad \varepsilon^4 = \varepsilon^{-4} = -1, \quad (\text{A.40})$$

we obtain from Table A.6 and (A.39):

$$c a^2 c^{-1} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = a^2 b. \quad (\text{A.41})$$

Also, by Table A.6 and (A.39) we have

$$ca^{-1}c^{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} = ac, \text{ and } ac^2 = ca^{-1}. \quad (\text{A.42})$$

By (A.41) and (A.42) we have

$$\begin{aligned} cac^{-1} &= ca^2c^{-1}ca^{-1}c^{-1} = (a^2b)(ac) = a^2a^{-1}bc = abc, \text{ i.e.,} \\ a &\sim abc. \end{aligned} \quad (\text{A.43})$$

Further,

$$cbc^{-1} = \begin{pmatrix} \varepsilon^2 & 0 \\ 0 & \varepsilon^{-2} \end{pmatrix} = a^2. \quad (\text{A.44})$$

By (A.43) and (A.44) we have

$$\begin{aligned} cabc^{-1} &= cac^{-1}cbc^{-1} = (abc)(a^2) \sim a^2(abc) = a^3bc, \\ ab &\sim a^3bc. \end{aligned} \quad (\text{A.45})$$

By (A.44)

$$cbc^{-1} = a^2, \quad cb^2c^{-1} = a^4, \quad (\text{A.46})$$

so, by (A.41) and (A.34) we have $ca^6c^{-1} = a^2ba^2ba^2b = a^2a^2b^{-1}ba^2b = a^6b$, i.e.,

$$b \sim a^2, \quad b^2 \sim a^4, \quad (\text{A.47})$$

and

$$a^2 \sim a^2b, \quad a^6 \sim a^6b. \quad (\text{A.48})$$

In addition, by (A.34) and (A.42) we have

$$a^pbc^2 = a^{p+1}a^{-1}bc^2 = a^{p+1}bac^2 = a^{p+1}bca^{-1} \sim a^{-1}a^{p+1}bca^{-1}a = a^pbc, \quad (\text{A.49})$$

i.e.,

$$a^pbc^2 \sim a^pbc, \quad (\text{A.50})$$

and

$$\begin{aligned} bc &\sim bc^2, \quad abc \sim abc^2, \quad a^2bc \sim a^2bc^2, \quad a^3bc \sim a^3bc^2, \\ a^4bc &\sim a^4bc^2, \quad a^5bc \sim a^5bc^2, \quad a^6bc \sim a^6bc^2, \quad a^7bc \sim a^7bc^2. \end{aligned} \quad (\text{A.51})$$

REMARK A.6. We denote by $Cl(g)$ the conjugacy class containing the element $g \in \mathcal{O}$. The union $Cl(b) \cup \{1, -1\}$ constitutes the 8-element subgroup

$$\{1, a^2, a^4, a^6, b, a^2b, a^4b, a^6b\} = \{1, a^2, -1, -a^2, b, a^2b, -b, -a^2b\}. \quad (\text{A.52})$$

Setting $i = b$, $j = a^2$, $k = a^2b$ we see that

$$\begin{aligned} i^2 &= j^2 = k^2 = -1, \\ ji &= -ij = k, \quad ik = -ki = j, \quad kj = -jk = i, \end{aligned} \quad (\text{A.53})$$

i.e., group (A.52) is the quaternion group Q_8 , see (A.3).

According to (A.34), (A.41) and (A.44) we deduce that Q_8 is a normal subgroup in $\mathcal{O}$:

$$Q_8 \triangleleft \mathcal{O}. \quad (\text{A.54})$$

3) Now we can prove conjugacy of elements in every row of Table A.7.

Conjugacy class $Cl(ab)$. By (A.36) and (A.45)

$$ab \sim a^3b \sim a^5b \sim a^7b \sim a^3bc.$$

By (A.33), (A.30) and (A.51)

$$a^3bc \sim a^5c^2, \quad a^3bc \sim a^7c, \quad a^3bc \sim a^3bc^2.$$

By (A.31), (A.51), (A.30) and (A.33)

$$a^3bc^2 \sim a^7bc^2 \sim a^7bc \sim a^3c, \quad a^7bc \sim ac^2.$$

Therefore, the elements

$$ab, a^3b, a^5b, a^7b, a^3bc, a^7bc, ac^2, a^5c^2, a^3c, a^7c, a^3bc^2, a^7bc^2$$

constitute a conjugacy class.

Conjugacy class $Cl(b)$. By (A.36), (A.47) and (A.29)

$$b \sim a^2b \sim a^4b \sim a^6b, \quad b \sim a^2, \quad a^2 \sim a^6,$$

i.e., the elements

$$a^2, a^6, b, a^2b, a^4b, a^6b$$

constitute a conjugacy class.

Conjugacy class $Cl(c^2)$. By (A.33), (A.32), (A.30), (A.33) and (A.51)

$$c^2 \sim a^6bc \sim a^4c, \quad c^2 \sim a^6c^2 \sim a^4bc \sim a^4bc^2, \quad a^6bc \sim a^6bc^2, \quad a^6c \sim a^4bc.$$

Thus, we have a conjugacy class

$$c^2, a^6c^2, a^4bc^2, a^6bc^2, a^4c, a^6c, a^4bc, a^6bc.$$

Conjugacy class $Cl(c)$. By (A.30), (A.33), (A.32) and (A.31)

$$c \sim a^2bc \sim a^4c^2 \sim a^2c^2, \quad a^2bc \sim a^2bc^2 \sim bc^2 \sim bc \sim a^2c,$$

i.e., the elements

$$c, a^2c, bc, a^2bc, a^4c^2, a^2c^2, bc^2, a^2bc^2$$

constitute a conjugacy class.

Conjugacy class $Cl(a)$. By (A.29), (A.42), (A.30), (A.33) and (A.51)

$$a \sim a^7 \sim ac \sim abc \sim a^3c^2, \quad abc \sim abc^2,$$

and we have a conjugacy class

$$a, a^7, ac, abc, a^3c^2, abc^2.$$

Conjugacy class $Cl(a^3)$. Since $a^7c^2 \sim c(a^7c^2)c^{-1} = ca^{-1}c$, we have by (A.42):

$$a^7c^2 \sim ca^{-1}c = ac^2c = ac^3.$$

Since $c^3 = a^4 = -1$, we deduce that

$$a^7c^2 \sim a^5. \tag{A.55}$$

By (A.29), (A.55), (A.33), (A.30) and (A.51)

$$a^3 \sim a^5 \sim a^7c^2 \sim a^5bc \sim a^5c, \quad a^5bc \sim a^5bc^2.$$

The elements

$$a^3, a^5, a^5c, a^5bc, a^7c^2, a^5bc^2.$$

constitute a conjugacy class. The proposition A.5 is proved. $\square$

Now consider conjugacy classes in the binary tetrahedral group $\mathcal{T}$. According to the Springer formula [Sp77, §4.4.10], the elements of the group $\mathcal{T}$ are given by (A.19) with even numbers h . In other words, each of the 24 different elements $x \in \mathcal{T}$ may be given as follows:

$$x = a^{2h}b^jc^l, \quad 0 \leq h < 4, \quad 0 \leq j < 2, \quad 0 \leq l < 3. \quad (\text{A.56})$$

There are 24 elements of type (A.56). In the octahedral group $\mathcal{O}$, the elements (A.56) constitute 5 conjugacy classes:

$$\{1\}, \{-1\}, Cl(b), Cl(c) \text{ and } Cl(c^2),$$

see Table A.7. We will see now, that in the tetrahedral group $\mathcal{T}$, the elements (A.56) constitute 7 conjugacy classes:

$$\{1\}, \{-1\}, Cl(b), Cl(c), Cl(a^4c) \text{ and } Cl(a^4c^2).$$

The elements $x \in \mathcal{T}$ and their traces are collected in Table A.8. We can now distinguish the elements of $\mathcal{T}$ by their traces and by means of two 1-dimensional representations τ_1 and τ_2 such that

$$\begin{aligned} \tau_1(a) &= 1, & \tau_1(b) &= 1, & \tau_1(c) &= \omega_3, \\ \tau_2(a) &= 1, & \tau_2(b) &= 1, & \tau_2(c) &= \omega_3^2, \end{aligned} \quad (\text{A.57})$$

where $\omega_3 = e^{2\pi i/3}$.

PROPOSITION A.7. *There are 7 conjugacy classes in the binary tetrahedral group $\mathcal{T}$. The rows of Table A.8 constitute these conjugacy classes.*

Proof. We will prove that the classes $Cl(b)$, $Cl(a^4c)$, $Cl(c^2)$, $Cl(c)$, and $Cl(a^4c^2)$ constitute all conjugacy classes.

Conjugacy class $Cl(b)$. Observe that relation (A.36) uses conjugating element a which does not lie in $\mathcal{T}$. We conjugate by a^{2p} :

$$b \sim a^2ba^{-2} = a^4b, \quad a^2b \sim a^6a^2ba^{-6} = a^6b. \quad (\text{A.58})$$

Then by (A.58), (A.29), (A.41) and (A.47) we have

$$b \sim a^4b, \quad a^2b \sim a^6b, \quad a^2 \sim a^6, \quad a^2 \sim a^2b, \quad b \sim a^2, \quad (\text{A.59})$$

i.e., the elements b , a^2 , a^6 , a^2b , a^4b , and a^6b constitute a conjugacy class.

Conjugacy class $Cl(a^4c)$. By (A.30) conjugating by b we get

$$a^4bc \sim a^6c, \quad a^6bc \sim a^4c. \quad (\text{A.60})$$

Further,

$$a^2ca^{-2} = \begin{pmatrix} \varepsilon^{-1} & \varepsilon^3 \\ -\varepsilon^{-3} & \varepsilon \end{pmatrix} = \begin{pmatrix} \varepsilon^{-1} & \varepsilon^3 \\ \varepsilon & \varepsilon \end{pmatrix}, \quad (\text{A.61})$$

and by Table A.6 we have

$$a^2ca^{-2} = bc, \quad c \sim bc, \quad (\text{A.62})$$

where conjugating element is a^2 . From (A.62) we obtain

$$a^6bc = a^6(a^2ca^{-2}) = ca^{-2} \sim a^6(ca^{-2})a^{-6} = a^6c. \quad (\text{A.63})$$

So,

$$a^6bc \sim a^6c, \quad (\text{A.64})$$

and together with (A.60) we see that the elements a^4c , a^4bc , a^6c , and a^6bc are conjugate.

Conjugacy class $Cl(c^2)$. By (A.32) and (A.31)

$$c^2 \sim a^6c^2, \quad a^4bc^2 \sim a^6bc^2. \quad (\text{A.65})$$

TABLE A.8. The conjugacy classes in the binary tetrahedral group

Trace $tr(x)$	Repr. τ_1	Repr. τ_2	Conjugacy class $Cl(g)$	Class order	Representative g
2	1	1	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	1	1
-2	1	1	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	1	-1
0	1	1	$a^2, a^6, b, a^2b, a^4b, a^6b$	6	b
-1	ω_3	ω_3^2	$a^4c, a^6c, a^4bc, a^6bc,$	4	$a^4c = -c$
-1	ω_3^2	ω_3	$c^2, a^6c^2, a^4bc^2, a^6bc^2$	4	c^2
1	ω_3	ω_3^2	c, a^2c, bc, a^2bc	4	c
1	ω_3^2	ω_3	$a^4c^2, a^2c^2, bc^2, a^2bc^2$	4	$a^4c^2 = -c^2$

Further, by (A.62), (A.44) and (A.34)

$$a^2c^2a^{-2} = (a^2ca^{-2})^2 = (bc)^2 = b(cb)c = b(a^2c)c = a^{-2}bc^2 = a^6bc^2, \quad (\text{A.66})$$

i.e.,

$$c^2 \sim a^6bc^2, \quad (\text{A.67})$$

and, with (A.65), we deduce that the elements c^2 , a^6c^2 , a^4bc^2 , and a^6bc^2 are conjugate.

Conjugacy class $Cl(c)$. By (A.62), (A.30)

$$c \sim bc, \quad c \sim a^2bc, \quad bc \sim a^2c, \quad (A.68)$$

i.e., c , bc , a^2bc , and a^2c are conjugate.

Conjugacy class $Cl(a^4c^2)$. By (A.30), (A.31)

$$a^4c^2 \sim a^2c^2, \quad bc^2 \sim a^2bc^2. \quad (A.69)$$

Further, by (A.46) we have

$$bc^2 \sim c(bc^2)c^{-1} = (cb)c = (a^2c)c = a^2c. \quad (A.70)$$

Thus, the elements a^4c^2 , a^2c^2 , bc^2 , a^2c are conjugate. The proposition A.7 is proved. $\square$

TABLE A.9. The characters of the binary octahedral group

Character ψ_i	Conjugacy class $Cl(g)$ and its order $ Cl(g) $ under it								Note on represent. ρ_i
	$Cl(1)$ 1	$Cl(-1)$ 1	$Cl(ab)$ 12	$Cl(b)$ 6	$Cl(c^2)$ 8	$Cl(c)$ 8	$Cl(a)$ 6	$Cl(a^3)$ 6	
ψ_0	1	1	1	1	1	1	1	1	trivial
ψ_1	1	1	-1	1	1	1	-1	-1	$\rho_1(a) = -a$
ψ_2	2	2	0	2	-1	-1	0	0	$\gamma_2\pi_2$
ψ_3	2	-2	0	0	-1	1	$\sqrt{2}$	$-\sqrt{2}$	faithful
ψ_4	2	-2	0	0	-1	1	$-\sqrt{2}$	$\sqrt{2}$	faithful
ψ_5	3	3	-1	-1	0	0	1	1	$\gamma_5\pi_{56}$
ψ_6	3	3	1	-1	0	0	-1	-1	$\gamma_6\pi_{56}$
ψ_7	4	-4	0	0	1	-1	0	0	$\rho_2 \otimes \rho_3$

A.5.3. Representations of the binary octahedral and tetrahedral groups.

PROPOSITION A.8. *The group $\mathcal{O}$ has the following 8 irreducible representations.*

1) *Two 1-dimensional representations:*

$$\begin{aligned} \rho_0(a) &= \rho_0(b) = \rho_0(c) = 1; \\ \rho_1(a) &= -1, \quad \rho_1(b) = \rho_1(c) = 1, \end{aligned} \quad (A.71)$$

2) *Two faithful 2-dimensional representations:*

$$\begin{aligned} \rho_3(a) &= a, \quad \rho_3(b) = b, \quad \rho_3(c) = c; \\ \rho_4(a) &= -a, \quad \rho_4(b) = b, \quad \rho_4(c) = c. \end{aligned} \quad (A.72)$$

3) *The 2-dimensional representation ρ_2 constructed by means of an epimorphism to the symmetric group S_3*

$$\mathcal{O} \longrightarrow \mathcal{O}/Q_8 \simeq S_3, \quad (A.73)$$

where Q_8 is the quaternion group.

4) Two 3-dimensional representations ρ_5 and ρ_6 constructed by means of an epimorphism to the symmetric group S_4

$$\mathcal{O} \longrightarrow \mathcal{O}/\{1, -1\} \simeq S_4. \quad (\text{A.74})$$

The representations ρ_5 and ρ_6 are related as follows

$$\rho_6(a) = -\rho_5(a), \quad \rho_6(b) = \rho_5(a), \quad \rho_6(c) = \rho_5(c). \quad (\text{A.75})$$

5) The 4-dimensional representation ρ_7 constructed as the tensor product $\rho_2 \otimes \rho_3$ (it coincides with $\rho_2 \otimes \rho_4$).

The characters of representations ρ_i for $i = 0, \dots, 7$ are collected in Table A.9.

Proof. 1) and 2) are clear from constructions (A.71) and (A.72).

3) We construct the third 2-dimensional representation ρ_2 by using the homomorphism

$$\pi_2 : \mathcal{O} \longrightarrow \mathcal{O}/Q_8, \quad (\text{A.76})$$

see (A.54). The quotient group $\mathcal{O}/Q_8$ is isomorphic to the symmetric group S_3 consisting of 6 elements. The cosets of $\mathcal{O}/Q_8$ are

$$\begin{aligned} \{Q_8, cQ_8, c^2Q_8, abQ_8, acQ_8, ac^2Q_8\} = \\ \{\{1\}, \{c\}, \{c^2\}, \{ab\}, \{ac\}, \{ac^2\}\}. \end{aligned} \quad (\text{A.77})$$

By (A.43), (A.42) and (A.27)

$$\begin{aligned} (ab)(ab) &= aa^{-1}b^2 = -1 \in Q_8, \text{ i.e., } \{ab\}^2 = \{1\}, \\ (ac)(ac) &= a(ca)c = a(abc^2)c = -a^2b \in Q_8, \text{ i.e., } \{ac\}^2 = \{1\}, \\ (ac^2)(ac^2) &= ca^{-1}(ac^2) = c^3 = -1 \in Q_8, \text{ i.e., } \{ac^2\}^2 = \{1\}, \\ (ab)(ac) &= bc \in cQ_8 \text{ i.e., } \{ab\}\{ac\} = \{c\}, \\ (ac)(ab) &= a(ca)b = a(abc^2)b = -a^2b(c^2b^{-1}) = \\ &= -a^2b(b^{-1}a^6c^2) = -c^2 \in c^2Q_8, \text{ i.e., } \{ab\}\{ac\} = \{c^2\}, \\ (c)(ab) &= (ca)b = (abc^2)b = -ab(c^2b^{-1}) = \\ &= -ab(b^{-1}a^6c^2) = a^3c^2 \in Q_8ac^2 = ac^2Q_8, \text{ i.e., } \{c\}\{ab\} = \{ac^2\}, \end{aligned} \quad (\text{A.78})$$

and so on. Thus,

$$\begin{aligned} \mathcal{O}/Q_8 &\simeq S_3, \\ \{ab\} &\simeq (12), \quad \{ac\} \simeq (13), \quad \{ac^2\} \simeq (23), \\ \{c\} &\simeq (123), \quad \{c^2\} \simeq (132). \end{aligned} \quad (\text{A.79})$$

The symmetric group S_3 has a 2-dimensional representation γ_2 such that

$$\begin{aligned} \text{tr } \gamma_2(12) &= \text{tr } \gamma_2(13) = \text{tr } \gamma_2(23) = 0, \quad \text{i.e.,} \\ \text{tr } \gamma_2\{ab\} &= \text{tr } \gamma_2\{ac\} = \text{tr } \gamma_2\{ac^2\} = 0, \\ \text{tr } \gamma_2(123) &= \text{tr } \gamma_2(132) = -1, \text{ i.e., } \text{tr } \gamma_2\{c\} = \text{tr } \gamma_2\{c^2\} = -1, \\ \text{tr } \gamma_2(1) &= 2, \end{aligned}$$

see, e.g., [CR62, §32]. We consider now the representation ρ_2 as the composition of epimorphism π_2 and the representation γ_2 , i.e.,

$$\rho_2 = \gamma_2 \pi_2. \quad (\text{A.80})$$

So, for all $u \in Cl(ab) \cup Cl(a) \cup Cl(a^3)$, we see that

$$\pi_2(u) \in \{ab\} \cup \{ac\} \cup \{ac^2\} \text{ and } \text{tr } \rho_2(u) = 0.$$

For all $u \in Cl(c) \cup Cl(c^2)$, we see that

$$\pi_2(u) \in \{c\} \cup \{c^2\} \text{ and } tr \rho_2(u) = -1.$$

Finally, for all $u \in Cl(1) \cup Cl(-1) \cup Cl(b)$ we see that

$$\pi_2(u) \in \{1\} \text{ and } tr \rho_2(u) = 2.$$

Thus we obtain the row of characters ψ_3 .

TABLE A.10. The 3-dimensional characters of S_4

Character	Conjugacy class $C_i \subset S_4$ and its order $ C_i $ under it				
	C_1	C_2	C_3	C_4	C_5
	1	6	8	6	3
γ_5	3	-1	0	1	-1
γ_6	3	1	0	-1	-1

4) We construct representations ρ_5 and ρ_6 by means of the epimorphism

$$\pi_{56} : \mathcal{O} \longrightarrow \mathcal{O}/\{1, -1\}. \quad (\text{A.81})$$

The epimorphism π_{56} is well-defined because the subgroup $\{1, -1\}$ is normal:

$$\{1, -1\} \triangleleft \mathcal{O}. \quad (\text{A.82})$$

The quotient group $\mathcal{O}/\{1, -1\}$ is the 24-element octahedral group coinciding with the symmetric group S_4 . By [CR62, §32] S_4 has two 3-dimensional representations, γ_5 and γ_6 , with characters as in Table A.10.

In Table A.10 we give conjugacy classes C_i of the group S_4 together with the number of elements of these classes:

$$\begin{aligned}
C_1 &= \{1\}, \\
C_2 &= \{ab, a^3b, a^3c, a^3bc, ac^2, a^3bc^2\}, \\
C_3 &= \{c, c^2, bc, a^2c, a^2bc, bc^2, a^2c^2, a^2bc^2\}, \\
C_4 &= \{a, a^3, ac, abc, a^3c^2, abc^2\}, \\
C_5 &= \{a^2, b, a^2b\}.
\end{aligned} \quad (\text{A.83})$$

We have

$$\rho_5 = \gamma_5 \pi_{56}, \quad \rho_6 = \gamma_6 \pi_{56}. \quad (\text{A.84})$$

For $u \in Cl(1) \cup Cl(-1)$, we see that

$$\pi_{56}(u) = 1 \text{ and } tr \rho_5(u) = 3.$$

For all $u \in Cl(ab)$, we see that

$$\pi_{56}(u) \in C_2 \text{ and } tr \rho_5(u) = -1.$$

For $u \in Cl(b)$, we see that

$$\pi_{56}(u) \in C_5 \text{ and } tr \rho_5(u) = -1.$$

For $u \in Cl(c) \cup Cl(c^2)$, we see that

$$\pi_{56}(u) \in C_3 \text{ and } tr \rho_5(u) = 0.$$

Finally, for every $u \in Cl(a) \cup Cl(a^2)$ we see that

$$\pi_{56}(u) \in C_4 \text{ and } tr \rho_5(u) = 1.$$

Thus we obtain the row of characters ψ_5 .

Note, that ρ_6 can be obtained from ρ_5 by the following relations:

$$\rho_6(a) = -\rho_5(a), \quad \rho_6(b) = \rho_5(b), \quad \rho_6(c) = \rho_5(c). \quad (\text{A.85})$$

5) Finally, the 4-dimensional representation ρ_7 is constructed as either of the tensor products $\rho_2 \otimes \rho_3$ or $\rho_2 \otimes \rho_4$. Observe that $\psi_2\psi_3$ and $\psi_2\psi_4$ have the same characters, see Table A.11.

TABLE A.11. The character of the 4-dimensional representation ρ_7

	$Cl(1)$	$Cl(-1)$	$Cl(ab)$	$Cl(b)$	$Cl(c^2)$	$Cl(c)$	$Cl(a)$	$Cl(a^3)$
$\psi_7 = \psi_2\psi_3 = \psi_2\psi_4$	4	-4	0	0	1	-1	0	0

The irreducibility of $\rho_2 \otimes \rho_3$ follows from the fact that

$$\langle \psi_2\psi_3, \psi_2\psi_3 \rangle = \frac{16 + 16 + 8 + 8}{48} = 1. \quad \square$$

Select ρ_3 as a faithful representation of $\mathcal{O}$ from the McKay correspondence. All irreducible representations ρ_i of $\mathcal{O}$ (see Proposition A.8 and Table A.9) can be placed in vertices of the extended Dynkin diagram $\tilde{E}_7$, see (A.86):

$$\begin{array}{ccccccccccc}
 \rho_0 & \text{---} & \rho_3 & \text{---} & \rho_5 & \text{---} & \rho_7 & \text{---} & \rho_6 & \text{---} & \rho_4 & \text{---} & \rho_1 \\
 & & & & & & \downarrow & & & & & & \\
 & & & & & & \rho_2 & & & & & &
 \end{array} \quad (\text{A.86})$$

Then, according to the McKay correspondence we have the following decompositions of the tensor products $\rho_3 \otimes \rho_i$:

$$\begin{aligned}
 \rho_3 \otimes \rho_0 &= \rho_3, \\
 \rho_3 \otimes \rho_1 &= \rho_4, \\
 \rho_3 \otimes \rho_2 &= \rho_7, \\
 \rho_3 \otimes \rho_3 &= \rho_0 + \rho_5, \\
 \rho_3 \otimes \rho_4 &= \rho_1 + \rho_6, \\
 \rho_3 \otimes \rho_5 &= \rho_3 + \rho_7, \\
 \rho_3 \otimes \rho_6 &= \rho_4 + \rho_7, \\
 \rho_3 \otimes \rho_7 &= \rho_2 + \rho_5 + \rho_6.
 \end{aligned} \quad (\text{A.87})$$

In Table A.12 we assume that $\omega_3 = e^{2\pi i/3}$.

PROPOSITION A.9. *The group $\mathcal{T}$ has the following 7 irreducible representations:*

TABLE A.12. The characters of the binary tetrahedral group

Character χ_i	Conjugacy class $Cl(g)$ and its order $ Cl(g) $ under it							Note on τ_i
	$Cl(1)$	$Cl(-1)$	$Cl(b)$	$Cl(c)$	$Cl(c^2)$	$Cl(-c)$	$Cl(-c^2)$	
	1	1	6	4	4	4	4	
χ_0	1	1	1	1	1	1	1	trivial
χ_1	1	1	1	ω_3	ω_3^2	ω_3	ω_3^2	$\tau_1(c) = \omega_3$
χ_2	1	1	1	ω_3^2	ω_3	ω_3^2	ω_3	$\tau_2(c) = \omega_3^2$
χ_3	2	-2	0	1	-1	-1	1	faithful
χ_4	2	-2	0	ω_3	$-\omega_3^2$	$-\omega_3$	ω_3^2	$\tau_3 \otimes \tau_1$
χ_5	2	-2	0	ω_3^2	$-\omega_3$	$-\omega_3^2$	ω_3	$\tau_3 \otimes \tau_2$
χ_6	3	3	-1	0	0	0	0	$\gamma_6 \pi_6$

1) *Three 1-dimensional representations*

$$\begin{aligned}
\tau_0(a) &= \tau_0(b) = \tau_0(c) = 1, \\
\tau_1(a) &= \tau_1(b) = 1, \quad \tau_1(c) = \omega_3, \\
\tau_2(a) &= \tau_2(b) = 1, \quad \tau_2(c) = \omega_3^2,
\end{aligned} \tag{A.88}$$

Representations τ_1 and τ_2 can be constructed by using an epimorphism onto the alternating group A_4 :

$$\mathcal{T} \longrightarrow \mathcal{T}/\{1, -1\} = A_4. \tag{A.89}$$

2) *The faithful 2-dimensional representation*

$$\tau_3(a) = a, \quad \tau_3(b) = b, \quad \tau_0(c) = c, \tag{A.90}$$

3) *Two 2-dimensional representation τ_4 and τ_5 constructed as tensor products*

$$\tau_4 = \tau_3 \otimes \tau_1, \quad \tau_5 = \tau_3 \otimes \tau_2. \tag{A.91}$$

4) *The 3-dimensional representation τ_6 constructed by using an epimorphism (A.89) onto the alternating group A_4 .*

The characters of representations τ_i for $i = 0, \dots, 6$ are collected in Table A.12.

Proof. 1), 2) and 3) are easily checked.

4) Consider the 3-dimensional representation of the alternating group A_4 with the character given in Table A.13, see, e.g., [CR62, §32]:

In Table A.13 we have $C_1 = \{1\}$, C_2 contains only elements of order 2, and C_3, C_4 contain only elements of order 3, see (A.92).

$$\begin{aligned}
C_1 &= \{1\}, \\
C_2 &= \{a^2, b, a^2b\}, \\
C_3 &= \{c, a^2c, bc, a^2bc\}, \\
C_4 &= \{c^2, a^2c^2, bc^2, a^2bc^2\}.
\end{aligned} \tag{A.92}$$

TABLE A.13. The 3-dimensional character of A_4

Character	Conjugacy class $C_i \subset A_4$ and its order $ C_i $ under it			
	C_1	C_2	C_3	C_4
	1	3	4	4
γ_6	3	-1	0	0

If π_6 is an epimorphism (A.89), then

$$\tau_6 = \gamma_6 \pi_6, \quad (\text{A.93})$$

and

$$\begin{aligned}
\tau_6 : Cl(1) \cup Cl(-1) &\longrightarrow C_1, & tr \tau_6(u) &= 3, \\
\tau_6 : Cl(b) &\longrightarrow C_2, & tr \tau_6(u) &= -1, \\
\tau_6 : Cl(c) \cup Cl(-c) &\longrightarrow C_3, & tr \tau_6(u) &= 0, \\
\tau_6 : Cl(c^2) \cup Cl(-c^2) &\longrightarrow C_4, & tr \tau_6(u) &= 0.
\end{aligned} \quad (\text{A.94})$$

Thus we get the last row in Table A.12. $\square$

Select τ_3 as a faithful representation of $\mathcal{T}$ from the McKay correspondence. All irreducible representations τ_i of $\mathcal{T}$ (see Proposition A.9 and Table A.12) can be placed in vertices of the extended Dynkin diagram $\tilde{E}_6$, see (A.95):

$$\begin{array}{ccccccc}
\tau_1 & \text{---} & \tau_4 & \text{---} & \tau_6 & \text{---} & \tau_5 & \text{---} & \tau_2 \\
& & & & | & & & & \\
& & & & \tau_3 & & & & \\
& & & & | & & & & \\
& & & & \tau_0 & & & &
\end{array} \quad (\text{A.95})$$

Then, according to the McKay correspondence, we have the following decompositions of the tensor products $\tau_3 \otimes \tau_i$:

$$\begin{aligned}
\tau_3 \otimes \tau_0 &= \tau_3, \\
\tau_3 \otimes \tau_1 &= \tau_4, \\
\tau_3 \otimes \tau_2 &= \tau_5, \\
\tau_3 \otimes \tau_3 &= \tau_0 + \tau_6, \\
\tau_3 \otimes \tau_4 &= \tau_1 + \tau_6, \\
\tau_3 \otimes \tau_5 &= \tau_2 + \tau_6, \\
\tau_3 \otimes \tau_6 &= \tau_3 + \tau_4 + \tau_5.
\end{aligned} \quad (\text{A.96})$$

A.5.4. Induced and restricted representations. Let us denote the characters of induced and restricted representations of $\mathcal{T}$ and $\mathcal{O}$ as follows:

$$\begin{aligned}
 &\text{irreducible representations } \tau \text{ of } \mathcal{T} \iff \text{characters are denoted by } \chi, \\
 &\text{irreducible representations } \rho \text{ of } \mathcal{O} \iff \text{characters are denoted by } \psi, \\
 &\text{induced representations } \tau \uparrow_{\mathcal{T}}^{\mathcal{O}} \text{ of } \mathcal{O} \iff \text{characters are denoted by } \chi^{\uparrow}, \\
 &\text{restricted representations } \rho \downarrow_{\mathcal{T}}^{\mathcal{O}} \text{ of } \mathcal{T} \iff \text{characters are denoted by } \psi^{\downarrow}.
 \end{aligned} \tag{A.97}$$

TABLE A.14. The restricted characters of the binary octahedral group

Character	Conjugacy class $Cl(g)$ and order of class $ Cl(g) $						
$\psi_i^{\downarrow}$	$Cl_{\mathcal{O}}(1)$	$Cl_{\mathcal{O}}(-1)$	$Cl_{\mathcal{O}}(b)$	$Cl_{\mathcal{O}}(c^2)$		$Cl_{\mathcal{O}}(c)$	
	1	1	6	8		8	
	$Cl_{\mathcal{T}}(1)$	$Cl_{\mathcal{T}}(-1)$	$Cl_{\mathcal{T}}(b)$	$Cl_{\mathcal{T}}(-c)$	$Cl_{\mathcal{T}}(c^2)$	$Cl_{\mathcal{T}}(c)$	$Cl_{\mathcal{T}}(-c^2)$
	1	1	6	4	4	4	4
$\psi_1^{\downarrow} = \psi_0^{\downarrow}$	1	1	1	1	1	1	1
$\psi_2^{\downarrow}$	2	2	2	-1	-1	-1	-1
$\psi_4^{\downarrow} = \psi_3^{\downarrow}$	2	-2	0	-1	-1	1	1
$\psi_6^{\downarrow} = \psi_5^{\downarrow}$	3	3	-1	0	0	0	0
$\psi_7^{\downarrow}$	4	-4	0	1	1	-1	-1


Consider the restriction of the binary octahedral group $\mathcal{O}$ onto the binary tetrahedral subgroup $\mathcal{T}$. Then the conjugacy classes $Cl(a)$, $Cl(a^3)$ and $Cl(ab)$ disappear, and the remaining 5 classes split into 7 conjugacy classes, see Table A.14. We denote the conjugacy classes of $\mathcal{O}$ by $Cl_{\mathcal{O}}$ and conjugacy classes of $\mathcal{T}$ by $Cl_{\mathcal{T}}$.

Now consider the restricted representations $\psi_i^{\downarrow}$ from $\mathcal{O}$ onto $\mathcal{T}$. By Table A.9 ψ_0 (resp. ψ_3 or ψ_5) differs from ψ_1 (resp. ψ_4 or ψ_6) only on $Cl(a)$, $Cl(a^3)$, and $Cl(ab)$, so we have the following coinciding pairs of restricted representations:

$$\psi_0^{\downarrow} = \psi_1^{\downarrow}, \quad \psi_3^{\downarrow} = \psi_4^{\downarrow}, \quad \psi_5^{\downarrow} = \psi_6^{\downarrow}. \tag{A.98}$$

The values of characters $\psi_i^{\downarrow}$ for $i = 0, 2, 3, 5, 7$ are easily obtained from the corresponding characters ψ_i .

Observe that $\rho_3^{\downarrow} = \tau_3$ is a faithful representation of $\mathcal{T}$ with character $\psi_3^{\downarrow} = \chi_3$. All irreducible representations $\rho_i^{\downarrow}$ for $i = 0, 2, 3, 5, 7$ can be placed in vertices of the extended

FIGURE A.2. The induced and restricted representations of $\mathcal{T} \triangleleft \mathcal{O}$

Dynkin diagram $\tilde{F}_{42}$, see Fig. A.2. From Table A.14 we have

$$\begin{aligned}
 \tau_3 \otimes \rho_0^\downarrow &= \rho_3^\downarrow \otimes \rho_0^\downarrow = \rho_3^\downarrow, \\
 \tau_3 \otimes \rho_2^\downarrow &= \rho_3^\downarrow \otimes \rho_2^\downarrow = \rho_7^\downarrow, \\
 \tau_3 \otimes \rho_3^\downarrow &= \rho_3^\downarrow \otimes \rho_3^\downarrow = \rho_0^\downarrow + \rho_5^\downarrow, \\
 \tau_3 \otimes \rho_5^\downarrow &= \rho_3^\downarrow \otimes \rho_5^\downarrow = \rho_3^\downarrow + \rho_7^\downarrow, \\
 \tau_3 \otimes \rho_7^\downarrow &= \rho_3^\downarrow \otimes \rho_7^\downarrow = \rho_2^\downarrow + 2\rho_5^\downarrow.
 \end{aligned} \tag{A.99}$$

Thus, the decompositions (A.99) constitute the following matrix

$$\tilde{A} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 2 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \rho_2^\downarrow \\ \rho_7^\downarrow \\ \rho_5^\downarrow \\ \rho_3^\downarrow \\ \rho_0^\downarrow \end{pmatrix} \tag{A.100}$$

where the row associated with $\rho_i^\downarrow$ for $i = 2, 7, 5, 3, 0$ consists of the decomposition coefficients (A.99) of $\tau_3 \otimes \rho_i^\downarrow$. The matrix $\tilde{A}$ satisfies the relation

$$\tilde{A} = 2I - K, \tag{A.101}$$

where K is the Cartan matrix for the extended Dynkin diagram $\tilde{F}_{42}$. The matrix K in (A.101) does not correspond to the bicolored order of vertices (see §3.1.3), but can be obtained from the bicolored one by permutations of rows and columns. We call the matrix $\tilde{A}$ the *Slodowy matrix*; it is an analog of the McKay matrix for the multiply-laced case.

Now we move on to the dual case and consider induced representations. To obtain induced representations $\chi \uparrow_T^{\mathcal{O}}$, we use the *Frobenius reciprocity formula* connecting restricted and induced representations, see, e.g., [JL2001, 21.16]

$$\langle \psi, \chi \uparrow_H^G \rangle_G = \langle \psi \downarrow_H^G, \chi \rangle_H. \quad (\text{A.102})$$

By (A.102) we have the following expression for the characters of induced representations

$$\chi \uparrow_H^G = \sum_{\psi_i \in \text{Irr}(G)} \langle \psi_i, \chi \uparrow_H^G \rangle_G \psi_i = \sum_{\psi_i \in \text{Irr}(G)} \langle \psi \downarrow_H^G, \chi \rangle_H \psi_i. \quad (\text{A.103})$$

Thus, to calculate the characters $\chi^\uparrow = \chi \uparrow_T^{\mathcal{O}}$, we only need to calculate the inner products

$$\langle \psi^\downarrow, \chi \rangle_T = \langle \psi \downarrow_T^{\mathcal{O}}, \chi \rangle_T. \quad (\text{A.104})$$

TABLE A.15. The inner products $\langle \psi^\downarrow, \chi \rangle$

	$\psi_1^\downarrow = \psi_0^\downarrow$	$\psi_2^\downarrow$	$\psi_3^\downarrow = \psi_4^\downarrow$	$\psi_5^\downarrow = \psi_6^\downarrow$	$\psi_7^\downarrow$
χ_0	1	0	0	0	0
χ_1	0	1	0	0	0
χ_2	0	1	0	0	0
χ_3	0	0	1	0	0
χ_4	0	0	0	0	1
χ_5	0	0	0	0	1
χ_6	0	0	0	1	0

One can obtain the inner products (A.104) from Tables A.12 and A.14. The results are given in Table A.15. Further, from Table A.15 and (A.103) we deduce

$$\begin{aligned}
\chi_0^\uparrow &= \psi_0 + \psi_1, & \tau_0^\uparrow &= \rho_0 + \rho_1, \\
\chi_1^\uparrow &= \chi_2^\uparrow = \psi_2, & \tau_1^\uparrow &= \tau_2^\uparrow = \rho_2, \\
\chi_3^\uparrow &= \psi_3 + \psi_4, & \tau_3^\uparrow &= \rho_3 + \rho_4, \\
\chi_4^\uparrow &= \chi_5^\uparrow = \psi_7, & \tau_4^\uparrow &= \tau_5^\uparrow = \rho_7, \\
\chi_6^\uparrow &= \psi_5 + \psi_6, & \tau_6^\uparrow &= \rho_5 + \rho_6.
\end{aligned} \quad (\text{A.105})$$

Let us find the tensor products $\rho_f \otimes \tau_i^\uparrow = \rho_3 \otimes \tau_i^\uparrow$, where ρ_f means the faithful representation of $\mathcal{O}$. By (A.87) and (A.105) we have

$$\begin{aligned}
\rho_3 \otimes \tau_0^\uparrow &= \rho_3 \otimes (\rho_0 + \rho_1) = \rho_3 \otimes \rho_0 + \rho_3 \otimes \rho_1 = \rho_3 + \rho_4 = \tau_3^\uparrow, \\
\rho_3 \otimes \tau_1^\uparrow &= \rho_3 \otimes \tau_2^\uparrow = \rho_3 \otimes \rho_2 = \rho_7 = \tau_4^\uparrow = \tau_5^\uparrow, \\
\rho_3 \otimes \tau_3^\uparrow &= \rho_3 \otimes (\rho_3 + \rho_4) = \\
&\quad \rho_3 \otimes \rho_3 + \rho_3 \otimes \rho_4 = (\rho_0 + \rho_5) + (\rho_1 + \rho_6) = \tau_0^\uparrow + \tau_6^\uparrow, \\
\rho_3 \otimes \tau_4^\uparrow &= \rho_3 \otimes \tau_5^\uparrow = \rho_3 \otimes \rho_7 = \rho_2 + \rho_5 + \rho_6 = \tau_1^\uparrow + \tau_6^\uparrow = \tau_2^\uparrow + \tau_6^\uparrow, \\
\rho_3 \otimes \tau_6^\uparrow &= \rho_3 \otimes (\rho_5 + \rho_6) = \rho_3 \otimes \rho_5 + \rho_3 \otimes \rho_6 = \\
&\quad (\rho_3 + \rho_7) + (\rho_4 + \rho_7) = \tau_3^\uparrow + 2\tau_4^\uparrow = \tau_3^\uparrow + 2\tau_5^\uparrow.
\end{aligned} \tag{A.106}$$

Here ρ_3 is the faithful representation of $\mathcal{O}$ with character ψ_3 . All irreducible representations $\tau_i^\uparrow$ for $i = 0, 2, 3, 5, 6$ can be placed in vertices of the extended Dynkin diagram $\tilde{F}_{41}$, see Fig. A.2. For other details, see [Sl80, App.III, p. 164].

The decompositions (A.106) constitute the following matrix

$$\tilde{A}^\vee = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix} \begin{matrix} \tau_2^\uparrow \\ \tau_5^\uparrow \\ \tau_6^\uparrow \\ \tau_3^\uparrow \\ \tau_0^\uparrow \end{matrix} \tag{A.107}$$

where the row associated with $\tau_i^\uparrow$ for $i = 2, 5, 6, 3, 0$ consists of the decomposition coefficients (A.106) of $\rho_3 \otimes \tau_i^\uparrow$. As in (A.101), the matrix $\tilde{A}^\vee$ satisfies the relation

$$\tilde{A}^\vee = 2I - K^\vee, \tag{A.108}$$

where $K^\vee$ is the Cartan matrix for the extended Dynkin diagram $\tilde{F}_{41}$. We see that the matrices $\tilde{A}$ and $\tilde{A}^\vee$ are mutually transposed:

$$\tilde{A}^t = \tilde{A}^\vee. \tag{A.109}$$

As in (A.101), we call the matrix $\tilde{A}^\vee$ the *Slodowy matrix*.

APPENDIX B

Miscellaneous

B.1. The triangle groups and Hurwitz groups

The group generated by X, Y, Z that satisfy the relations (A.6)

$$X^p = Y^q = Z^r = XYZ = 1$$

is said to be a *triangle group*. As it was mentioned in §A.2, the finite polyhedral groups from Table A.1 are triangle groups. Set

$$\mu(p, q, r) := \frac{1}{p} + \frac{1}{q} + \frac{1}{r}.$$

The triangle group is finite if and only if $\mu(p, q, r) > 1$. There are only three triangle groups

$$(2, 4, 4), \quad (2, 3, 6), \quad (3, 3, 3),$$

for which $\mu(p, q, r) = 1$. These groups are infinite and soluble. For references, see [Con90], [Mu01]. For all other triangle groups, we have

$$\mu(p, q, r) < 1.$$

These groups are infinite and insoluble. The value $1 - \mu(p, q, r)$ attains the minimum value $\frac{1}{42}$ at $(2, 3, 7)$. Thus, $(2, 3, 7)$ is, in a sense, the minimal infinite insoluble triangle group. The importance of the triangle group $(2, 3, 7)$ is revealed by the following theorem due to Hurwitz.

THEOREM B.1 (Hurwitz, [Hur1893]). *If X is a compact Riemann surface of genus $g > 1$, then $|AutX| \leq 84(g - 1)$, and moreover, the upper bound of this order is attained if and only if $|AutX|$ is a homomorphic image of the triangle group $(2, 3, 7)$.*

For further references, see [Con90], [Con03].

A *Hurwitz group* is any finite nontrivial quotient of the triangle group $(2, 3, 7)$. In other words, the finite group G is the Hurwitz group if it has generators $X, Y, Z \in G$ such that

$$X^2 = Y^3 = Z^7 = XYZ = 1.$$

M. Conder writes that the significance of the Hurwitz groups “...is perhaps best explained by referring to some aspects of the theory of Fuchsian groups, hyperbolic geometry, Riemann surfaces...”, see [Con90, p.359] and a bibliography cited there.

M. Conder [Con80] using the method of coset graphs developed by G. Higman have shown that the alternating group A_n is a Hurwitz group for all $n \geq 168$.

Recently, A. Lucchini, M. C. Tamburini and J. S. Wilson showed that most finite simple classical groups of sufficiently large rank are Hurwitz groups, see [LuT99], [LuTW00]. For example, the groups $SL_n(q)$ are Hurwitz, for all $n > 286$ [LuTW00], and the groups $Sp_{2n}(q)$, $SU_{2n}(q)$ are Hurwitz, for all $n > 371$ [LuT99]. (Mentioned groups act in the n -dimensional vector space over the field F_q of the prime characteristic q).

The sporadic groups have been treated in a series of papers by Woldar and others, see [Wi01] for a survey and references. R. A. Wilson shows in [Wi01], that the Monster is also a Hurwitz group.

B.2. Algebraic integers

If λ is a root of the polynomial equation

$$a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0, \quad (\text{B.1})$$

where a_i for $i = 0, 1, \dots, n$ are integers and λ satisfies no similar equation of degree $< n$, then λ is said to be an *algebraic number* of degree n . If λ is an algebraic number and $a_n = 1$, then λ is called an *algebraic integer*.

A polynomial $p(x)$ in which the coefficient of the highest order term is equal to 1 is called the *monic polynomial*. The polynomial (B.1) with integer coefficients and $a_n = 1$ is the *monic integer polynomial*.

The algebraic integers of degree 1 are the ordinary integers (elements of $\mathbb{Z}$). If α is an algebraic number of degree n satisfying the polynomial equation

$$(x - \alpha)(x - \beta)(x - \gamma) \cdots = 0,$$

then there are $n-1$ other algebraic numbers $\beta, \gamma, \dots$ called the *conjugates* of α . Furthermore, if α satisfies any other algebraic equation, then its conjugates also satisfy the same equation.

DEFINITION B.2. An algebraic integer $\lambda > 1$ is said to be a *Pisot number* if its conjugates (other than λ itself) satisfy $|\lambda'| < 1$.

The smallest Pisot number,

$$\lambda_{\text{Pisot}} \approx 1.324717\dots, \quad (\text{B.2})$$

is a root of $\lambda^3 - \lambda - 1 = 0$ (for details and references, see [McM02]). This number appears in Proposition 4.15 as a limit of the spectral radius $\rho(T_{2,3,n})$ as $n \rightarrow \infty$. This number was also obtained by Y. Zhang [Zh89] and used in the study of regular components of an Auslander-Reiten quiver, so we call the number (B.2) the *Zhang number*.

DEFINITION B.3. Let $p(x)$ be a monic integer polynomial, and define its *Mahler measure* to be

$$\|p(x)\| = \prod_{\beta} |\beta|, \quad (\text{B.3})$$

where β runs over all (complex) roots of $p(x)$ outside the unit circle.

It is well known that $\|p(x)\| = 1$ if and only if all roots of $p(x)$ are roots of unity. In 1933, Lehmer [Leh33] asks whether for each $\Delta \geq 1$, there exists an algebraic integer such that

$$1 < \|\alpha\| < 1 + \Delta \quad (\text{B.4})$$

Lehmer found polynomials with smallest Mahler measure for small degrees and stated in [Leh33, p.18] that the polynomial with minimal root α (in the sense of B.4) he could find is the polynomial of degree 10:

$$1 + x - x^3 - x^4 - x^5 - x^6 - x^7 + x^9 + x^{10}, \quad (\text{B.5})$$

see [Hir02], [McM02]; cf. Remark 4.14. Outside the unit circle, the polynomial (B.5) has only one root

$$\lambda_{\text{Lehmer}} \approx 1.176281\dots \quad (\text{B.6})$$

The number (B.6) is called *Lehmer's number*; see Proposition 4.15, Remark 4.14 and Table 4.4.

DEFINITION B.4. An algebraic integer $\lambda > 1$ is said to be a *Salem number* if its conjugates λ' (other than λ itself) satisfy $|\lambda'| \leq 1$ and include $\frac{1}{\lambda}$.

It is known that every Pisot number is a limit of Salem numbers; for details and references, see [McM02, p.177]. Conjecturally, Lehmer's number (B.6) is the smallest Salem number, [Leh33], [GH01].

The positive root of the quadratic equation $\lambda^2 - \lambda - 1 = 0$ is a well-known constant

$$\lambda_{\text{Golden}} \approx 1.618034\dots, \quad (\text{B.7})$$

called the *Golden mean* or *Divine proportion*. This number appears in Proposition 4.17 as a limit of the spectral radius $\rho(T_{3,3,n})$ as $n \rightarrow \infty$.

The smallest Mahler measure among polynomials of degree at most 6 is

$$M_6 = \|x^6 - x^4 + x^3 - x^2 + 1\| \approx 1.401268\dots, \quad (\text{B.8})$$

see [Mos98, p.1700]. This number appears in Proposition 4.17 as a root of

$$\chi(T_{3,3,4}) = x^8 + x^7 - 2x^5 - 3x^4 - 2x^3 + x + 1.$$

This is the polynomial of minimal degree among polynomials $\chi(T_{3,3,n})$, where $n = 4, 5, 6, \dots$ with indefinite Tits form, see Proposition 4.17 and Table 4.5.

B.3. The Perron-Frobenius Theorem

We say that a matrix is *positive* (resp. *nonnegative*) if all its entries are positive (resp. nonnegative). We use the notation $A > 0$ (resp. $A \geq 0$) for positive (resp. nonnegative) matrix. A square $n \times n$ matrix A is called *reducible* if the indices $1, 2, \dots, n$ can be divided into the disjoint union of two nonempty sets $\{i_1, i_2, \dots, i_p\}$ and $\{j_1, j_2, \dots, j_q\}$ (with $p + q = n$) such that

$$a_{i_\alpha j_\beta} = 0, \text{ for } \alpha = 1, \dots, p \text{ and } \beta = 1, \dots, q.$$

In other words, A is *reducible* if there exists a permutation matrix P , such that

$$PAP^t = \begin{pmatrix} B & 0 \\ C & D \end{pmatrix},$$

where B and D are square matrices. A square matrix which is not reducible is said to be *irreducible*.

THEOREM B.5 (Perron-Frobenius). *Let A be $n \times n$ nonnegative irreducible matrix. Then the following holds:*

1) *One of its eigenvalues λ is positive and greater than or equal to (in absolute value) all other eigenvalues λ_i :*

$$|\lambda_i| \leq \lambda, \quad i = 1, 2, \dots, n.$$

2) *There is a positive eigenvector z corresponding to that eigenvalue:*

$$Az = \lambda z, \text{ where } z = (z_1, \dots, z_n)^t \text{ and } z_i > 0 \text{ for } i = 1, 2, \dots, n.$$

Such an eigenvalue λ is called the dominant eigenvalue of A .

3) *The eigenvalue λ is a simple root of the characteristic equation of A .*

The following important corollary from the Perron-Frobenius theorem holds for the eigenvalue λ :

$$\lambda = \max_{z \geq 0} \min_i \frac{(Az)_i}{z_i} \quad (z_i \neq 0),$$

$$\lambda = \min_{z \geq 0} \max_i \frac{(Az)_i}{z_i} \quad (z_i \neq 0).$$

For details, see [MM64], [Ga90].

B.4. The complex projective line and stereographic projection

The n -dimensional *complex projective space* $\mathbb{C}P^n$ is the set of all complex lines in $\mathbb{C}^{n+1}$ passing through the origin. A point $z \in \mathbb{C}P^n$ is defined up to the *complex* factor w . In other words, two points

$$z_1 = (z_1^0, z_1^1, z_1^2, \dots, z_1^n), \quad z_2 = (z_2^0, z_2^1, z_2^2, \dots, z_2^n) \in \mathbb{C}^{n+1}$$

lie on the same line, if

$$\begin{aligned} z_2 &= wz_1 \text{ for some complex multiplier } w \in \mathbb{C}, \text{ i.e.} \\ z_2^i &= wz_1^i \text{ for } i = 0, 1, \dots, n. \end{aligned} \tag{B.9}$$

The points (B.9) constitute an equivalence class denoted by $[z^0 : z^1 : \dots : z^n]$.

Clearly, the *complex projective line* $\mathbb{C}P^1$ is the set of all lines in $\mathbb{C}^2$. By (B.9), the points of $\mathbb{C}P^1$ are classes of complex pairs $z = [z^0, z^1]$ up to a factor $w \in \mathbb{C}$.

The correspondence

$$[z^0 : z^1] \mapsto z^0/z^1 \tag{B.10}$$

sets the following bijection maps:

$$\begin{aligned} \mathbb{C}P^1 \setminus \{[1 : 0]\} &\Longleftrightarrow \mathbb{C}, \text{ and} \\ \mathbb{C}P^1 &\Longleftrightarrow \mathbb{C} \cup \infty. \end{aligned} \tag{B.11}$$

Let z_1, z_2 be two vectors from $\mathbb{C}^2$. Define a map

$$F : \mathbb{C}^2 \longrightarrow \mathbb{R}^3 \tag{B.12}$$

by setting

$$F(z_1, z_2) := \left(\frac{z_1 \bar{z}_2 + \bar{z}_1 z_2}{z_1 \bar{z}_1 + \bar{z}_2 z_2}, \frac{z_1 \bar{z}_2 - \bar{z}_1 z_2}{i(z_1 \bar{z}_1 + \bar{z}_2 z_2)}, \frac{z_1 \bar{z}_1 - \bar{z}_2 z_2}{z_1 \bar{z}_1 + \bar{z}_2 z_2} \right), \tag{B.13}$$

(see, for example, [Alv02]).

Let $z_2 = wz_1$, then

$$F(z_1, z_2) = \left(\frac{\bar{w} + w}{1 + |w|^2}, \frac{\bar{w} - w}{1 + |w|^2}, \frac{1 - |w|^2}{1 + |w|^2} \right). \tag{B.14}$$

If $w = u + iv$, then

$$F(z_1, z_2) = \left(\frac{2u}{1 + u^2 + v^2}, \frac{2v}{1 + u^2 + v^2}, \frac{1 - u^2 - v^2}{1 + u^2 + v^2} \right). \tag{B.15}$$

It is easily to see that

$$(2u)^2 + (2v)^2 + (1 - u^2 - v^2)^2 = (1 + u^2 + v^2)^2,$$

so $F(z_1, z_2)$ maps every vector $[z_1, z_2]$ to a point (B.15) on the unit sphere S^2 in $\mathbb{R}^3$. Another vector $[z_3, z_4]$ defines the same line in the $\mathbb{C}P^1$ if and only if

$$z_4/z_3 = z_2/z_1 = w,$$

i.e., the map F (B.14) defines a bijection from the complex projective line $\mathbb{C}P^1$ and the unit sphere in $\mathbb{R}^3$:

$$F : \mathbb{C}P^1 \Longleftrightarrow S^2. \tag{B.16}$$

Let (x, y, z) be a point on S^2 distinct from the *north pole* $N = \{0, 0, 1\}$. The *stereographic projection* is the map

$$S : S^2 \setminus N \longrightarrow \mathbb{C} \tag{B.17}$$

defined by

$$S(x, y, z) := \frac{x}{1-z} + i \frac{y}{1-z}. \quad (\text{B.18})$$

see Fig. B.1. For details, see, for example, [Jen94, §2.9].




FIGURE B.1. The stereographic projection

Consider the composition $S(F(z_1, z_2))$. By (B.13), we have

$$1 - z = \frac{2\bar{z}_2 z_2}{z_1 \bar{z}_1 + \bar{z}_2 z_2}, \quad (\text{B.19})$$

and

$$\frac{x}{1-z} = \frac{z_1 \bar{z}_2 + \bar{z}_1 z_2}{2\bar{z}_2 z_2} = \frac{1}{2} \left(\frac{z_1}{z_2} + \frac{\bar{z}_1}{\bar{z}_2} \right), \quad (\text{B.20})$$

$$\frac{y}{1-z} = \frac{z_1 \bar{z}_2 - \bar{z}_1 z_2}{2\bar{z}_2 z_2} = \frac{1}{2} \left(\frac{z_1}{z_2} - \frac{\bar{z}_1}{\bar{z}_2} \right).$$

Let

$$\frac{z_1}{z_2} = w = u + iv.$$

Then

$$\frac{x}{1-z} = \frac{1}{2} (w + \bar{w}) = u, \quad \frac{y}{1-z} = \frac{1}{2} (w - \bar{w}) = v, \quad (\text{B.21})$$

and

$$S(F(z_1, z_2)) = u + iv = w = \frac{z_1}{z_2}. \quad (\text{B.22})$$

B.5. Prime spectrum, coordinate ring, orbit space

B.5.1. Hilbert's Nullstellensatz (Theorem of zeros). From now on, we assume that R is a commutative ring with identity.

A proper ideal $\mathfrak{m}$ of R is said to be *maximal* if $\mathfrak{m}$ is not a proper subset of any other proper ideal of R . An ideal $\mathfrak{m} \subset R$ is maximal if and only if the quotient ring $R/\mathfrak{m}$ is a field. For example, every ideal $p\mathbb{Z}$ is maximal in the ring of integers $\mathbb{Z}$, if p is a prime number, and, in this case, the quotient ring $\mathbb{Z}/p\mathbb{Z}$ is a field.

A proper ideal $\mathfrak{p}$ of a commutative ring R is called a *prime ideal* if the following condition holds:

$$\text{for any } a, b \in R, \text{ if } a \cdot b \in \mathfrak{p}, \text{ then either } a \in \mathfrak{p} \text{ or } b \in \mathfrak{p}. \quad (\text{B.23})$$

An ideal $\mathfrak{p} \subset R$ is prime if and only if the quotient ring $R/\mathfrak{p}$ is an *integral domain* (i.e., the commutative ring which has no divisors of 0). Examples of prime ideals:

$$\begin{aligned} &\text{ideal } (5, \sqrt{6}) \text{ in the ring } \mathbb{Z}[\sqrt{6}], \\ &\text{ideal } (x) \text{ in the ring } \mathbb{Z}[x], \\ &\text{ideal } (y + x + 1) \text{ in the ring } \mathbb{C}[x, y]. \end{aligned} \tag{B.24}$$

The maximal ideals are prime since the fields are integral domains, but not conversely:

$$\boxed{\text{maximal ideals}} \subset \boxed{\text{prime ideals}}. \tag{B.25}$$

The ideal $(x) \in \mathbb{Z}[x]$ is prime, but not maximal, since, for example:

$$(x) \subset (2, x) \subset \mathbb{Z}[x]. \tag{B.26}$$

Let k be an algebraically closed field (e.g., the complex field $\mathbb{C}$), and let I be an ideal in $k[x_1, \dots, x_n]$. Define $\mathbb{V}_k(I)$, the *zero set* of I , by

$$\mathbb{V}_k(I) = \{(a_1, \dots, a_n) \in k \mid f(a_1, \dots, a_n) = 0 \text{ for all } f \in I\}. \tag{B.27}$$

Denote by $\mathbb{A}_k^n$ (or just $\mathbb{A}^n$) the n -dimensional *affine space* over the field k . The *Zariski topology* on $\mathbb{A}_k^n$ is defined to be the topology whose closed sets are the zero sets $\mathbb{V}_k(I)$.

For any ideal I of the commutative ring R , the radical $\sqrt{I}$ of I , is the set

$$\{a \in R \mid a^n \in I \text{ for some integer } n > 0\}. \tag{B.28}$$

The radical of an ideal I is always an ideal of R . If $I = \sqrt{I}$, then I is called a *radical ideal*. The prime ideals are radical:

$$\boxed{\text{prime ideals}} \subset \boxed{\text{radical ideals}}. \tag{B.29}$$

THEOREM B.6. (Hilbert's Nullstellensatz, [Re88, Ch.2], or [Mum88, pp.9-11])

Let k be an algebraically closed field.

(1) The maximal ideals in the ring $A = k[x_1, \dots, x_n]$ are the ideals

$$\mathfrak{m}_P = (x_1 - a_1, \dots, x_n - a_n) \tag{B.30}$$

for some point $P = (a_1, \dots, a_n)$. The ideal $\mathfrak{m}_P$ coincides with the ideal $I(P)$ of all functions which are zero in P .

(2) If $\mathbb{V}_k(I) = \emptyset$, then $I = k[x_1, \dots, x_n]$.

(3) For any ideal $J \subset A$

$$I(\mathbb{V}_k(J)) = \sqrt{J}, \tag{B.31}$$

The set $X \subset \mathbb{A}_k^n$ is called the *affine variety* if $X = \mathbb{V}_k(I)$ for some ideal $I \subset A$, see (B.27). An affine variety $X \subset \mathbb{A}_k^n$ is called *irreducible affine variety* if there does not exist a decomposition

$$X = X_1 \cup X_2,$$

where X_1, X_2 are two proper subsets of X . For example, the affine variety

$$X = \{(x, y) \in \mathbb{A}_{\mathbb{C}}^2 \mid xy = 0\}$$

is decomposed to the sum of

$$X_1 = \{(x, y) \in \mathbb{A}_{\mathbb{C}}^2 \mid x = 0\} \text{ and } X_2 = \{(x, y) \in \mathbb{A}_{\mathbb{C}}^2 \mid y = 0\}.$$

PROPOSITION B.7. Let $X \subset \mathbb{A}_k^n$ be an affine variety and $I = I(X)$ the corresponding ideal, i.e., $X = \mathbb{V}_k(I)$, see (B.27). Then

$$X \text{ is irreducible} \iff I(X) \text{ is the prime ideal.}$$

For a proof of this proposition, see, e.g., [Re88, §3.7].

One of the important corollaries of Hilbert's Nullstellensatz is the following correspondence [Re88, §3.10] between subvarieties $X \subset \mathbb{A}_k^n$ and ideals $I \subset A$:

$$\begin{array}{ccc} \{ \text{radical ideals} \} & \Longleftrightarrow & \{ \text{affine variety} \} \\ \bigcup & & \bigcup \\ \{ \text{prime ideals} \} & \Longleftrightarrow & \{ \text{irreducible affine variety} \} \\ \bigcup & & \bigcup \\ \{ \text{maximal ideals} \} & \Longleftrightarrow & \{ \text{points} \} \end{array} \quad (\text{B.32})$$

B.5.2. Prime spectrum. The *prime spectrum* $\text{Spec}(R)$ of the commutative ring R is defined to be the set of proper prime ideals of R :

$$\{\mathfrak{p} \subset R \mid \mathfrak{p} \text{ is a prime ideal of } R\}. \quad (\text{B.33})$$

The ring R itself is not counted as a prime ideal, but (0) , if prime, is counted.

A topology is imposed on $\text{Spec}(R)$ by defining the sets of *closed sets*. The closed set $V(J)$ is the set of the form

$$\{\mathfrak{p} \mid \mathfrak{p} \supseteq J \text{ for some ideal } J \subset R\}. \quad (\text{B.34})$$

This set is denoted by $V(J)$. The *closure* $\overline{\mathcal{P}}$ of the subset $\mathcal{P}$ is the intersection of all closed sets containing $\mathcal{P}$:

$$\overline{\mathcal{P}} = \bigcap_{V(J) \supset \mathcal{P}} V(J). \quad (\text{B.35})$$

Consider the closure of points of the topological space $\text{Spec}(R)$. For a point of $\text{Spec}(R)$ which is the prime ideal $\mathfrak{p}$, we have the closure

$$\overline{\mathfrak{p}} = \bigcap_{V(J) \supset \mathfrak{p}} V(J) = V(\mathfrak{p}), \quad (\text{B.36})$$

which consists of all prime ideals $\mathfrak{p}' \supset \mathfrak{p}$. In particular, the closure $\overline{\mathfrak{p}}$ consists of one point if and only if the ideal $\mathfrak{p}$ is maximal. The point which coincides with its closure is called the *closed point*. Thus, closed points in $\text{Spec}(R)$ correspond one-to-one with maximal ideals.

In the conditions of Hilbert's Nullstellensatz every maximal ideal J is $\mathfrak{m}_P$ (B.30) defined by some point $P = (a_1, \dots, a_n) \in k^n$. Thus, in the finitely generated ring $A = k[x_1, \dots, x_n]$ over an algebraically closed field k the closed points in $\text{Spec}(A)$ correspond to the maximal ideals and correspond to the points $P = (a_1, \dots, a_n) \in k^n$.

There exist non-closed points in the topological space $\text{Spec}(R)$. Let R has no divisors of 0. Then the ideal (0) is prime and is contained in all other prime ideals. Thus, the closure $\overline{(0)}$ in (B.36) consists of all prime ideals, i.e. coincides with the space $\text{Spec}(R)$. The point (0) is everywhere dense point in $\text{Spec}(R)$. The everywhere dense is called the *generic point*.

DEFINITION B.8. Let $\mathcal{P}$ is an irreducible closed subset of $\text{Spec}(R)$. A point $a \in \mathcal{P}$ is called the *generic point* of $\mathcal{P}$ if the closure $\overline{a}$ coincides with $\mathcal{P}$.

PROPOSITION B.9. ([Mum88, p.126]) *If $x \in \text{Spec}(R)$, then the closure $\overline{\{x\}}$ of $\{x\}$ is irreducible and x is a generic point of $\overline{\{x\}}$. Conversely, every irreducible closed subset $\mathcal{P} \subset \text{Spec}(R)$ equals $V(J)$ for some prime ideal $J \subset R$ and J is its unique generic point.*

A *distinguished open set* of $\text{Spec}(R)$ is defined to be an open set of the form

$$\text{Spec}(R) := \{\mathfrak{p} \in \text{Spec}(R) \mid f \notin \mathfrak{p}\}. \quad (\text{B.37})$$

for any element $f \in R$.

EXAMPLE B.10. 1) If R is a field, then $\text{Spec}(R)$ has just one point (0) .

2) Let $R = k[X]$ be a polynomial ring in one variable x . Then $\text{Spec}(R)$ is the affine line A_k^1 over k . There exist two types of prime ideals: (0) and $(f(x))$, where f is an irreducible polynomial, corresponding to the prime ideal $(f(x))$. For algebraically closed k , the closed points are all of the form $(X - a)$. The point (0) is generic.

3) $\text{Spec}(\mathbb{Z})$ consists of closed points for every prime ideal (p) , plus ideal (0) .

B.5.3. Coordinate ring. Let $V \subset \mathbb{A}_k^n$ be an affine variety and $I(V)$ be an ideal of V . The *coordinate ring* of the affine variety V is the quotient ring denoted by $k[V]$ and defined as follows:

$$k[V] := k[x_1, \dots, x_n]/I(V). \quad (\text{B.38})$$

A *regular function* on the affine variety V is the restriction to V of a polynomial in $x_1, \dots, x_n$, modulo $I(V)$ (i.e., modulo functions vanishing on V). Thus, the regular functions on the affine variety V are elements of the coordinate ring $k[V]$.

EXAMPLE B.11. 1) Let $f(x, y)$ be a complex polynomial function from $\mathbb{C}^2$. The coordinate ring of a plane curve defined by the equation $f(x, y) = 0$ in $\mathbb{A}_{\mathbb{C}}^2$ is

$$\mathbb{C}[x, y]/(f(x, y)), \quad (\text{B.39})$$

where $(f(x, y))$ is the ideal generated by the polynomial $f(x, y)$.

2) Consider two algebraic curves: $y = x^r$ ($r \in \mathbb{N}$), and $y = 0$. The coordinate rings of these curves are isomorphic:

$$\begin{array}{ll} y = x^r & \text{corresponds to } \mathbb{C}[x, y]/(y - x^r) \simeq \mathbb{C}[x, x^r] \simeq \mathbb{C}[x], \\ y = 0 & \text{corresponds to } \mathbb{C}[x, y]/(y) \simeq \mathbb{C}[x, x] \simeq \mathbb{C}[x]. \end{array}$$

Thus, in the sense of algebraic geometry, curves $y = x^r$ and $y = 0$ are the same.

3) On the other hand

$$\mathbb{C}[x, y]/(y^2 - x^3) \not\simeq \mathbb{C}[x],$$

and the curves $y^2 = x^3$ and $y = 0$ are not the same (in the sense of algebraic geometry). Really, there exists the isomorphism

$$x \longmapsto T^2, \quad y \longmapsto T^3,$$

and

$$\mathbb{C}[x, y]/(y^2 - x^3) \simeq \mathbb{C}[T^2, T^3] \subset \mathbb{C}[T].$$

The affine variety $X = \{(x, y) \mid y^2 = x^3\}$ is called *Neile's parabola*, or the *semicubical parabola*.

REMARK B.12. Consider the coordinate ring $R = k[V]$ of the affine variety V over algebraically closed k . Then,

*the prime spectrum $\text{Spec}(R)$ contains exactly
the same information as the variety V .*

Really, according to Hilbert's Nullstellensatz, the maximal ideals of $k[V]$ correspond one-to-one with points of V :

$$v \in V \iff \mathfrak{m}_v \subset k[V].$$

Besides, according to Proposition B.7 every other prime ideal $\mathfrak{p} \subset k[V]$ is the intersection of maximal ideals corresponding to the points of some irreducible subvariety $Y \subset V$:

$$\mathfrak{p}_Y = \bigcap_{v \in Y} \mathfrak{m}_v.$$

B.5.4. Orbit space. Let G be a finite group acting on affine variety V and $k[V]$ be the coordinate ring of V . The problem here is that the quotient space V/G may not exist, even for very trivial group actions, see Example B.13. To find how this problem is solved in the geometric invariant theory (GIT), see [Dol03] or [Kr85].

EXAMPLE B.13. Consider the multiplicative group $G = k^*$ on the affine line A_k^1 . An orbit space for the group G consists of two orbits: $\{0\}$ and $A_k^1 - \{0\}$. The second orbit $A_k^1 - \{0\}$ is not closed subset in the Zariski's topology, and the first orbit $\{0\}$ is contained in the closure of the orbit $A_k^1 - \{0\}$. Thus, the point $\{0\}$ is the generic point (§B.5.2).

The following two definitions of the *categorical quotient* and the *orbit space* can be found, e.g., in the P. E. Newstead textbook [Newst78, p.39].

DEFINITION B.14. Let G be an algebraic group acting on a variety V . A *categorical quotient* of V by G is a pair (Y, φ) , where Y is a variety and φ is a morphism $\varphi : V \rightarrow Y$ such that

- (i) φ is constant on the orbits of the action;
- (ii) for any variety Y' and morphism $\varphi' : V \rightarrow Y'$ which is constant on orbits, there is a unique morphism $\psi : Y \rightarrow Y'$ such that $\psi \circ \varphi = \varphi'$.

DEFINITION B.15. A categorical quotient of V by G is called an *orbit space* if $\varphi^{-1}(y)$ consists of a single orbit for all $y \in Y$. The *orbit space* is denoted by V/G .

The *orbit space* $X = V/G$ is an affine variety whose points correspond one-to-one with orbits of the group action.

PROPOSITION B.16. Let G be a finite group of order n acting on affine variety V . Assume that the characteristic of k does not divide n . Set

$$X := \operatorname{Spec} k[V]^G. \quad (\text{B.40})$$

Then X is an orbit space for the action G on V .

For a proof of this theorem see, e.g., [Shaf88, Ch.1 §2, Ex.11] or [Rom00, Th.1.6].

B.6. The orbit structure of the Coxeter transformation

Let $\mathfrak{g}$ be the simple complex Lie algebra of type A, D or E , and $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$. Let $\mathfrak{h}^\vee$ be the dual space to $\mathfrak{h}$ and $\alpha_i \in \mathfrak{h}^\vee, i = 1, \dots, l$ be an ordered set of simple positive roots. Here, we follow B. Kostant's description [Kos84] of the orbit structure of the Coxeter transformation $\mathbf{C}$ on the highest root in the root system of $\mathfrak{g}$. We consider a bipartite graph and a bicolored Coxeter transformation from §3.1.1, §3.1.2. Let β be the highest root of $(\mathfrak{h}, \mathfrak{g})$, see §4.1. Between two bicolored Coxeter transformations (3.1) we select w_1, w_2 such that

$$w_2\beta = \beta. \quad (\text{B.41})$$

Consider, for example, the Dynkin diagram E_6 . Here,

$$w_1 = \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ 1 & 1 & 0 & -1 & & \\ 1 & 0 & 1 & & -1 & \\ 1 & 0 & 0 & & & -1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \\ y_1 \\ y_2 \\ y_3 \end{pmatrix}, \quad w_2 = \begin{pmatrix} -1 & & & & & \\ & -1 & & & & \\ & & -1 & & & \\ & & & 1 & 1 & 1 \\ & & & 1 & 0 & 0 \\ & & & & 1 & 0 \\ & & & & & 1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \\ y_1 \\ y_2 \\ y_3 \end{pmatrix}, \quad (\text{B.42})$$

where any vector $z \in \mathfrak{h}^\vee$ and the highest vector β are:

$$x_1 \quad y_1 \quad x_0 \quad y_2 \quad x_2, \quad \beta = \begin{matrix} 1 & 2 & 3 & 2 & 1 \\ & & 2 & & \end{matrix}, \quad \text{or} \quad \begin{pmatrix} 3 \\ 1 \\ 1 \\ 2 \\ 2 \\ 2 \end{pmatrix} \begin{matrix} x_0 \\ x_1 \\ x_2 \\ y_1 \\ y_2 \\ y_3 \end{matrix}, \quad (\text{B.43})$$

Further, following B.Kostant [Kos84, Th.1.5] consider the alternating products $\tau^{(n)}$:

$$\begin{aligned} \tau^{(1)} &= w_1, \\ \tau^{(2)} &= \mathbf{C} = w_2 w_1, \\ \tau^{(3)} &= w_1 \mathbf{C} = w_1 w_2 w_1, \\ &\dots, \\ \tau^{(n)} &= \begin{cases} \mathbf{C}^k = w_2 w_1 \dots w_2 w_1 & \text{for } n = 2k, \\ w_1 \mathbf{C}^k = w_1 w_2 w_1 \dots w_2 w_1 & \text{for } n = 2k + 1, \end{cases} \end{aligned} \quad (\text{B.44})$$

and the orbit of $\tau^{(n)}$ on the highest vector β :

$$\beta_n = \tau^{(n)} \beta, \quad (\text{B.45})$$

where $n = 1, \dots, h-1$ (h is the Coxeter number, see(4.1)).

THEOREM B.17. (B. Kostant, [Kos84, Theorems 1.3, 1.4, 1.5, 1.8])

1) *There exist $z_j \in \mathfrak{h}^\vee, j = 1, \dots, h-1$ and even integers $2 \leq a \leq b \leq h$ (see (5.58) and Table 5.1) such that the generating functions $P_G(t)$ (see (1.6), (1.21), or (5.52), (5.53)) are obtained as follows:*

$$[P_G(t)]_i = \begin{cases} \frac{1+t^h}{(1-t^a)(1-t^b)} & \text{for } i = 0, \\ \frac{\sum_{j=1}^{h-1} z_j t^j}{(1-t^a)(1-t^b)} & \text{for } i = 1, \dots, r. \end{cases} \quad (\text{B.46})$$

Indices $i = 1, \dots, r$ enumerate vertices of the Dynkin diagram and the coordinates of the vectors z_n ; $i = 0$ corresponds to the additional vertex, the one that extends the Dynkin diagram to the extended Dynkin diagram.

2) *Vectors z_n (we call these vectors the assembling vectors) are obtained by means of the orbit of $\tau^{(n)}$ on the highest vector β as follows:*

$$z_n = \tau^{(n-1)} \beta - \tau^{(n)} \beta. \quad (\text{B.47})$$

3) *On has*

$$z_g = 2\alpha_*, \quad \text{where } g = \frac{h}{2}, \quad (\text{B.48})$$

and α_ is the simple root corresponding to the branch point for diagrams D_n, E_n and to the midpoint for the diagram A_{2m-1} . In all these cases h is even, and g is integer. The diagram A_{2m} has been excluded.*

4) *The series of assembling vectors z_n is symmetry:*

$$z_{g+k} = z_{g-k} \quad \text{for } k = 1, \dots, g. \quad (\text{B.49})$$

TABLE B.1. Orbit of the Coxeter transformation on the highest root

β	$\tau^{(1)}\beta = w_1\beta$	$\tau^{(2)}\beta = \mathbf{C}\beta$
1 2 3 2 1	1 2 3 2 1	1 2 2 2 1
2	1	1
$\tau^{(3)}\beta = w_1\mathbf{C}\beta$	$\tau^{(4)}\beta = \mathbf{C}^2\beta$	$\tau^{(5)}\beta = w_1\mathbf{C}^2\beta$
1 1 2 1 1	0 1 1 1 0	0 0 1 0 0
1	1	0
$\tau^{(6)}\beta = \mathbf{C}^3\beta$	$\tau^{(7)}\beta = w_1\mathbf{C}^3\beta$	$\tau^{(8)}\beta = \mathbf{C}^4\beta$
0 0 -1 0 0	0 -1 -1 -1 0	-1 -1 -2 -1 -1
0	-1	-1
$\tau^{(9)}\beta = w_1\mathbf{C}^4\beta$	$\tau^{(10)}\beta = \mathbf{C}^5\beta$	$\tau^{(11)}\beta = w_1\mathbf{C}^5\beta$
-1 -2 -2 -2 -1	-1 -2 -3 -2 -1	-1 -2 -3 -2 -1
-1	-1	-2

TABLE B.2. Assembling vectors $z_n = \tau^{(n-1)}\beta - \tau^{(n)}\beta$

$z_1 = \beta - w_1\beta$	$z_2 = w_1\beta - \mathbf{C}\beta$	$z_3 = \mathbf{C}\beta - w_1\mathbf{C}\beta$
0 0 0 0 0	0 0 1 0 0	0 1 0 1 0
1	0	0
$z_4 = w_1\mathbf{C}\beta - \mathbf{C}^2\beta$	$z_5 = \mathbf{C}^2\beta - w_1\mathbf{C}^2\beta$	$z_6 = w_1\mathbf{C}^2\beta - \mathbf{C}^3\beta$
1 0 1 0 1	0 1 0 1 0	0 0 2 0 0
0	1	0
$z_7 = \mathbf{C}^3\beta - w_1\mathbf{C}^3\beta$	$z_8 = w_1\mathbf{C}^3\beta - \mathbf{C}^4\beta$	$z_9 = \mathbf{C}^4\beta - w_1\mathbf{C}^4\beta$
0 1 0 1 0	1 0 1 0 1	0 1 0 1 0
1	0	0
$z_{10} = w_1\mathbf{C}^4\beta - \mathbf{C}^5\beta$	$z_{11} = \mathbf{C}^5\beta - w_1\mathbf{C}^5\beta$	
0 0 1 0 0	0 0 0 0 0	
0	1	

In the case of the Dynkin diagram E_6 vectors $\tau^{(n)}\beta$ are given in Table B.1, and the assembling vectors z_n are given in Table B.2. Vector z_6 coincides with $2\alpha_{x_0}$, where α_{x_0} is the simple root corresponding to the vertex x_0 , see (B.43). From Table B.2 we see, that

$$z_1 = z_{11}, \quad z_2 = z_{10}, \quad z_3 = z_9, \quad z_4 = z_8, \quad z_5 = z_7. \quad (\text{B.50})$$

Denote by $z(t)_i$ the polynomial $\sum_{j=1}^{h-1} z_j t^j$ from (B.46). In the case of E_6 we have:

$$\begin{aligned} z(t)_{x_1} &= t^4 + t^8, \\ z(t)_{y_1} &= t^3 + t^5 + t^7 + t^9, \\ z(t)_{y_2} &= t^3 + t^5 + t^7 + t^9, \\ z(t)_{x_2} &= t^4 + t^8, \\ z(t)_{y_3} &= t + t^5 + t^7 + t^{11}. \end{aligned} \tag{B.51}$$

The Kostant numbers a, b (see Table 5.1) for E_6 are $a = 6, b = 8$. From (B.46) and (B.51), we have

$$\begin{aligned} [P_G(t)]_{x_1} &= [P_G(t)]_{x_2} = \frac{t^4 + t^8}{(1 - t^6)(1 - t^8)}, \\ [P_G(t)]_{y_1} &= [P_G(t)]_{y_2} = \frac{t^3 + t^5 + t^7 + t^9}{(1 - t^6)(1 - t^8)}, \\ [P_G(t)]_{y_3} &= \frac{t + t^5 + t^7 + t^{11}}{(1 - t^6)(1 - t^8)}. \end{aligned} \tag{B.52}$$

Since

$$1 - t^6 = \sum_{n=0}^{\infty} t^{6n}, \quad 1 - t^8 = \sum_{n=0}^{\infty} t^{8n}, \tag{B.53}$$

we have

$$\begin{aligned} [P_G(t)]_{x_1} &= [P_G(t)]_{x_2} = \sum_{i=0, j=0}^{\infty} (t^{6i+8j+4} + t^{6i+8j+8}), \\ [P_G(t)]_{y_1} &= [P_G(t)]_{y_2} = \sum_{i=0, j=0}^{\infty} (t^{6i+8j+3} + t^{6i+8j+5} + t^{6i+8j+7} + t^{6i+8j+9}), \\ [P_G(t)]_{y_3} &= \sum_{i=0, j=0}^{\infty} (t^{6i+8j+1} + t^{6i+8j+5} + t^{6i+8j+7} + t^{6i+8j+11}), \end{aligned} \tag{B.54}$$

Recall, that $m_\alpha(n), \alpha = x_1, x_2, y_1, y_2, y_3$ are the multiplicities of the indecomposable representations ρ_α of G (considered in the context of McKay correspondence, §(A.4)) in the decomposition of $\pi_n|G$ (5.45). These multiplicities are the coefficients of the Poincaré series (B.54), see (5.50), (5.52), (5.53). For example,

$$[P_G(t)]_{x_1} = [P_G(t)]_{x_2} = t^4 + t^8 + t^{10} + t^{12} + t^{14} + 2t^{16} + t^{18} + 2t^{20} + \dots \tag{B.55}$$

$$m_{x_1} = m_{x_2} = \begin{cases} 0 & \text{for } n = 1, 2, 3, 5, 6 \text{ and } n = 2k + 1, k \geq 3, \\ 1 & \text{for } n = 4, 8, 10, 12, 14, 18, \dots \\ 2 & \text{for } n = 16, 20, \dots \\ \dots & \end{cases} \quad (\text{B.56})$$

In particular representations $\rho_{x_1}(n)$ and $\rho_{x_2}(n)$ do not enter in the decomposition of π_n of $SU(2)$ (see §5.4.1) for all odd n .

B.7. Fixed and anti-fixed points of the Coxeter transformation

B.7.1. Chebyshev polynomials and the McKay-Slodowy matrix. We continue to use block arithmetic for $(m + k) \times (m + k)$ matrices as in §3.1.3, namely:

$$K = \begin{pmatrix} 2I & 2D \\ 2F & 2I \end{pmatrix}, \quad w_1 = \begin{pmatrix} -I & -2D \\ 0 & I \end{pmatrix}, \quad w_2 = \begin{pmatrix} I & 0 \\ -2F & -I \end{pmatrix}. \quad (\text{B.57})$$

Let

$$\beta = \frac{1}{2}K - I = \begin{pmatrix} 0 & D \\ F & 0 \end{pmatrix}. \quad (\text{B.58})$$

The matrix β coincides (up to a factor $-\frac{1}{2}$) with the McKay matrix from §A.4 or with the Slodowy matrix in the multiply-laced case, (A.101), (A.108).

Introduce two recursive series $f_p(\beta)$ and $g_p(\beta)$ of polynomials in β :

$$\begin{aligned} f_0(\beta) &= 0, \\ f_1(\beta) &= K = 2\beta + 2I, \\ f_{p+2}(\beta) &= 2\beta f_{p+1}(\beta) - f_p(\beta) \quad \text{for } p \geq 0 \end{aligned} \quad (\text{B.59})$$

and

$$\begin{aligned} g_0(\beta) &= 2I, \\ g_1(\beta) &= K - 2I = 2\beta, \\ g_{p+2}(\beta) &= 2\beta g_{p+1}(\beta) - g_p(\beta) \quad \text{for } p \geq 0. \end{aligned} \quad (\text{B.60})$$

For example,

$$\begin{aligned} f_2(\beta) &= 2\beta(2\beta + 2I) = (K - 2I)K = 4 \begin{pmatrix} DF & D \\ F & FD \end{pmatrix}, \\ f_3(\beta) &= (K - 2I)(K - 2I)K - K = 2 \begin{pmatrix} 4DF - I & 4DFD - D \\ 4FDF - F & 4FD - I \end{pmatrix}, \end{aligned} \quad (\text{B.61})$$

$$\begin{aligned} g_2(\beta) &= 2(2\beta^2 - I) = 2 \begin{pmatrix} 2DF - I & 0 \\ 0 & 2FD - I \end{pmatrix}, \\ g_3(\beta) &= 4\beta(2\beta^2 - I) - 2\beta = 2 \begin{pmatrix} 0 & 8DFD - 6D \\ 8FDF - 6F & 0 \end{pmatrix}. \end{aligned} \quad (\text{B.62})$$

REMARK B.18. Let us formally substitute $\beta = \cos t$. Then

1) The polynomials $g_p(\beta)$ are Chebyshev polynomials of the first kind (up to a factor $\frac{1}{2}$):

$$g_p(\cos t) = 2 \cos pt.$$

Indeed,

$$g_1(\beta) = 2\beta = 2 \cos t, \quad g_2(\beta) = 2(2\beta^2 - 1) = 2 \cos 2t,$$

and

$$g_{p+2}(\beta) = 2(2 \cos t \cos(p+1)t - \cos pt) = 2 \cos(p+2)t.$$

2) The polynomials $f_p(\beta)$ are Chebyshev polynomials of the second kind (up to a factor $2\beta + 2$):

$$f_p(\cos t) = 2(\cos t + 1) \frac{\sin pt}{\sin t}.$$

Indeed,

$$f_1(\beta) = 2\beta + 2I = 2 \cos t + 2, \quad f_2(\beta) = 2\beta(2\beta + 2I) = 4 \cos t(\cos t + 1),$$

and

$$f_{p+2}(\beta) = 2 \frac{\cos t + 1}{\sin t} (2 \cos t \sin(p+1)t - \sin pt) = 2(\cos t + 1) \frac{\sin(p+2)t}{\sin t}.$$

B.7.2. A theorem on fixed and anti-fixed points. The purpose of this section is to prove the following

THEOREM B.19 ([SuSt82]). 1) *The fixed points of the powers of the Coxeter transformation satisfy the following relations*

$$\mathbf{C}^p z = z \iff f_p(\beta)z = 0.$$

2) *The anti-fixed points of the powers of the Coxeter transformation satisfy the following relations*

$$\mathbf{C}^p z = -z \iff g_p(\beta)z = 0.$$

In the proof of Theorem B.19 we will need some properties of block matrices. Define the involution σ by setting

$$\sigma : A = \begin{pmatrix} P & Q \\ S & T \end{pmatrix} \mapsto \bar{A} = \begin{pmatrix} P & Q \\ -S & -T \end{pmatrix}. \quad (\text{B.63})$$

The map σ is a linear operator because $\overline{\alpha_1 A_1 + \alpha_2 A_2} = \alpha_1 \bar{A}_1 + \alpha_2 \bar{A}_2$.

The involution σ does not preserve products:

$$\overline{AR} \neq \bar{A}\bar{R},$$

but

$$\overline{AR} = \bar{A} R \quad \text{for any } R = \begin{pmatrix} X & Y \\ U & V \end{pmatrix}, \quad (\text{B.64})$$

and

$$R \bar{A} = -\overline{RA} \quad \text{for } R = \begin{pmatrix} 0 & Y \\ U & 0 \end{pmatrix}. \quad (\text{B.65})$$

The following relations are easy to check:

$$\mathbf{C} = w_1 w_2 = \begin{pmatrix} 4DF - I & 2D \\ -2F & -I \end{pmatrix}, \quad (\text{B.66})$$

$$\mathbf{C}^{-1} = w_2 w_1 = \begin{pmatrix} -I & -2D \\ 2F & 4FD - I \end{pmatrix},$$

$$w_2\mathbf{C} = \begin{pmatrix} 4DF - I & 2D \\ -8FDF + 4F & -4FD + I \end{pmatrix}, \quad (\text{B.67})$$

$$w_1\mathbf{C}^{-1} = \begin{pmatrix} -4DF + I & 4D - 8DFD \\ 2F & 4FD - I \end{pmatrix},$$

$$w_1 - w_2 = 4 \begin{pmatrix} -2I & -2D \\ 2F & 2I \end{pmatrix} = -\overline{K}, \quad w_1 + w_2 = -(K - 2I), \quad (\text{B.68})$$

$$\mathbf{C} - \mathbf{C}^{-1} = 4 \begin{pmatrix} DF & D \\ -F & -FD \end{pmatrix} = 4 \begin{pmatrix} \overline{DF} & \overline{D} \\ \overline{F} & \overline{FD} \end{pmatrix}, \quad (\text{B.69})$$

$$\mathbf{C} + \mathbf{C}^{-1} = 2 \begin{pmatrix} 2DF - I & 0 \\ 0 & 2FD - I \end{pmatrix},$$

$$\begin{aligned} w_2\mathbf{C} - w_1\mathbf{C}^{-1} &= 2 \begin{pmatrix} 4DF - I & 4DFD - D \\ -(4FDF - F) & -(4FD - I) \end{pmatrix} = \\ &= 2 \begin{pmatrix} \overline{4DF - I} & \overline{4DFD - D} \\ \overline{4FDF - F} & \overline{4FD - I} \end{pmatrix}, \end{aligned} \quad (\text{B.70})$$

$$w_2\mathbf{C} + w_1\mathbf{C}^{-1} = \begin{pmatrix} 0 & -8DFD + 6D \\ -8FDF + 6F & 0 \end{pmatrix}, \quad (\text{B.71})$$

$$2\beta w_2 + I = -\mathbf{C}, \quad 2\beta w_1 + I = -\mathbf{C}^{-1}. \quad (\text{B.72})$$

PROPOSITION B.20. *The following relations hold*

$$w_2 - w_1 = \overline{f_1(\beta)}, \quad (\text{B.73})$$

$$\mathbf{C} - \mathbf{C}^{-1} = \overline{f_2(\beta)}, \quad (\text{B.74})$$

$$w_2\mathbf{C} - w_1\mathbf{C}^{-1} = \overline{f_3(\beta)}, \quad (\text{B.75})$$

$$w_2 + w_1 = -g_1(\beta), \quad (\text{B.76})$$

$$\mathbf{C} + \mathbf{C}^{-1} = g_2(\beta), \quad (\text{B.77})$$

$$w_2\mathbf{C} + w_1\mathbf{C}^{-1} = -g_3(\beta). \quad (\text{B.78})$$

Proof. The proof of this theorem follows from comparing relations (B.59)-(B.60) and relations (B.69)-(B.71).

(B.73) follows from (B.59) and (B.68).

(B.74) follows from (B.61) and (B.69).

(B.75) follows from (B.61) and (B.70).

(B.76) follows from (B.60) and (B.68).

(B.77) follows from (B.62) and (B.69).

(B.78) follows from (B.62) and (B.71). $\square$

PROPOSITION B.21. *The following relations hold:*

$$\overline{f_{2m}(\beta)} = \mathbf{C}^m - \mathbf{C}^{-m}, \quad \overline{f_{2m+1}(\beta)} = w_2 \mathbf{C}^m - w_1 \mathbf{C}^{-m}, \quad (\text{B.79})$$

$$g_{2m}(\beta) = \mathbf{C}^m + \mathbf{C}^{-m}, \quad g_{2m+1}(\beta) = -(w_2 \mathbf{C}^m + w_1 \mathbf{C}^{-m}). \quad (\text{B.80})$$

Proof. 1) Relations (B.73)-(B.75) realize the basis of induction.

By (B.65), by the induction hypothesis, and by (B.72) we have

$$\begin{aligned} \overline{f_{2m+2}(\beta)} &= \overline{2\beta f_{2m+1}(\beta) - f_{2m}(\beta)} = -2\beta \overline{f_{2m+1}(\beta)} - \overline{f_{2m}(\beta)} = \\ &= -2\beta(w_2 \mathbf{C}^m - w_1 \mathbf{C}^{-m}) - (\mathbf{C}^m - \mathbf{C}^{-m}) = \\ &= -(2\beta w_2 + I)\mathbf{C}^m + (2\beta w_1 + I)\mathbf{C}^{-m} = \mathbf{C}^{m+1} - \mathbf{C}^{-(m+1)}. \end{aligned}$$

In the same way we have

$$\begin{aligned} \overline{f_{2m+3}(\beta)} &= -2\beta \overline{f_{2m+2}(\beta)} - \overline{f_{2m+1}(\beta)} = \\ &= -2\beta(\mathbf{C}^{m+1} - \mathbf{C}^{-(m+1)}) - w_2 \mathbf{C}^m - w_1 \mathbf{C}^{-m} = \\ &= -(2\beta w_1 + I)w_2 \mathbf{C}^m + (2\beta w_2 + I)w_1 \mathbf{C}^{-m} = \\ &= \mathbf{C}^{-1}w_2 \mathbf{C}^m - \mathbf{C}w_1 \mathbf{C}^{-m} = w_1 \mathbf{C}^{m+1} - w_2 \mathbf{C}^{-(m+1)}. \end{aligned}$$

2) Relations (B.76)-(B.78) realize the basis of induction. By the induction hypothesis and by (B.72) we have

$$\begin{aligned} g_{2m+2}(\beta) &= 2\beta g_{2m+1}(\beta) - g_{2m}(\beta) = \\ &= -2\beta(w_2 \mathbf{C}^m + w_1 \mathbf{C}^{-m}) - (\mathbf{C}^m + \mathbf{C}^{-m}) = \\ &= -(2\beta w_2 + I)\mathbf{C}^m - (2\beta w_1 + I)\mathbf{C}^{-m} = \mathbf{C}^{m+1} + \mathbf{C}^{-(m+1)}. \end{aligned}$$

By analogy, we have

$$\begin{aligned} g_{2m+3}(\beta) &= 2\beta g_{2m+2}(\beta) - g_{2m+1}(\beta) = \\ &= -2\beta(\mathbf{C}^{m+1} + \mathbf{C}^{-(m+1)}) - (w_2 \mathbf{C}^m + w_1 \mathbf{C}^{-m}) = \\ &= -(2\beta w_1 + I)w_2 \mathbf{C}^m - (2\beta w_2 + I)w_1 \mathbf{C}^{-m} = w_1 \mathbf{C}^{m+1} + w_2 \mathbf{C}^{-(m+1)}. \quad \square \end{aligned}$$

Theorem B.19 now follows from Proposition B.21.

Bibliography

- [ArGV86] V. I. Arnold, S. M. Gussein-Zade, A. N. Varchenko. *Singularities of differentiable maps* I, II. Birkhäuser, Basel, 1986, 1988.
- [AtMa69] M. F. Atiyah, I. G. MacDonald, *Introduction to Commutative Algebra*. Reading, Addison-Wesley, 1969.
- [AuRS95] M. Auslander, I. Reiten, S. O. Smalø, *Representation Theory of Artin Algebras*. Cambridge Studies in Advanced Mathematics 36, Cambridge, 1995.
- [AuPR79] M. Auslander, M. I. Platzcek, I. Reiten, *Coxeter Functors without Diagrams*. Trans. Amer. Math. Soc. 250 (1979), 1–46.
- [A'C75] N. A'Campo, *Le groupe de monodromie du déploiement des singularités isolées de courbes planes*. I. (French) Math. Ann. 213 (1975), 1–32.
- [A'C76] N. A'Campo, *Sur les valeurs propres de la transformation de Coxeter*. Invent. Math. 33 (1976), no. 1, 61–67.
- [Alv02] J. C. Álvarez Paiva *Interactive Course on Projective Geometry. Geometry of the Complex Projective Line*, 2000,
http://www.math.poly.edu/courses/projective_geometry .
- [Bak04] A. Baker, *Representations of Finite Groups*.
<http://www.maths.gla.ac.uk/~ajb/dvi-ps/groupreps.pdf> .
- [Bol96] A. Boldt, *Two Aspects of Finite-Dimensional Algebras: Uniserial Modules and Coxeter Polynomials*. University of California, Ph.D. thesis, 1996.
- [BT97] A. Boldt, M. Takane, *The spectral classes of unicyclic graphs*. J. Pure Appl. Algebra 133 (1998), no. 1-2, 39–49.
- [Bo] N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4,5,6*. Paris, Hermann, 1968.
- [BLM89] S. Berman, Y. S. Lee, R. V. Moody, *The spectrum of a Coxeter transformation, affine Coxeter transformations and the defect map*. J. Algebra 121 (1989), no. 2, 339–357.
- [BGP73] I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev, *Coxeter functors, and Gabriel's theorem*. Uspehi Mat. Nauk 28 (1973), no. 2(170), 19–33. English translation: Russian Math. Surveys 28 (1973), no. 2, 17–32.
- [BC55] A. Borel, C. Chevalley, *The Betti numbers of the exceptional groups*. Mem. Amer. Math. Soc. (1955). no. 14, 1–9.
- [Br35] R. Brauer, *Sur les invariants intégraux des variétés représentatives des groupes de Lie simples clos*, C. R. Acad. Sci. Paris, 201,(1935), 419–421.
- [Car70] R. Carter, *Conjugacy classes in the Weyl group*. 1970 Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, N.J., 1968/69) 297–318, Springer, Berlin.
- [Col58] A. J. Coleman, *The Betti numbers of the simple Lie groups*. Canad. J. Math. 10 (1958), 349–356.
- [Col89] A. J. Coleman, *Killing and the Coxeter transformation of Kac-Moody algebras*. Invent. Math. 95 (1989), no. 3, 447–477.
- [Con80] M. Conder, *Generators for alternating and symmetric groups*. J. London Math. Soc. (2) 22 (1980), no. 1, 75–86.
- [Con90] M. Conder, *Hurwitz groups: a brief survey*. Bull. Amer. Math. Soc. (N.S.) 23 (1990), no. 2, 359–370.
- [Con03] M. Conder, *Group actions on graphs, maps and surfaces with maximum symmetry*. invited paper in: Groups St Andrews 2001 in Oxford, London Math. Soc. Lecture Note Series, vol. 304, Cambridge University Press, 63–91, 2003.
- [Cox34] H. S. M. Coxeter, *Discrete groups generated by reflections*. Ann. of Math. (2) 35 (1934), no. 3, 588–621.
- [Cox40] H. S. M. Coxeter, *The binary polyhedral groups, and other generalizations of the quaternion group*. Duke Math. J. 7, (1940), 367–379.

- [Cox49] H. S. M. Coxeter, *Regular polytopes*. New York (1949).
- [Cox51] H. S. M. Coxeter, *The product of the generators of a finite group generated by reflections*. Duke Math. J. 18, (1951), 765–782.
- [CoxM84] H. S. M. Coxeter and W. O. J. Moser, *Generators and relations for discrete groups*. 4th ed., Springer-Verlag, NY, 1984.
- [Cr01] A. Craw, *The McKay correspondence and representations of the McKay quiver*. Warwick Ph.D. thesis, 2001.
- [Ch50] C. Chevalley, *The Betti numbers of the exceptional simple Lie groups*. Proceedings of the International Congress of Mathematicians, Cambridge, Mass., 1950, vol. 2, Amer. Math. Soc., Providence, R. I., 1952, 21–24.
- [Ch55] C. Chevalley, *Invariants of finite groups generated by reflections*. Amer. J. Math. 77 (1955), 778–782.
- [CR62] C. W. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*. Wiley (1962).
- [Dol03] I. Dolgachev, *Lectures on invariant theory January*. London Mathematical Society Lecture Note Series, 296. Cambridge University Press, Cambridge, (2003), xvi+220 pp.
- [DR76] V. Dlab, C. M. Ringel, *Indecomposable representations of graphs and algebras*. Mem. Amer. Math. Soc. 6 (1976), no. 173.
- [DR81] V. Dlab and C. M. Ringel, *Eigenvalues of Coxeter transformations and the Gelfand-Kirillov dimension of preprojective algebras*. Proc. Amer. Math. Soc. 83 (2) (1981), 228–232.
- [Drz74] Ju. A. Drozd, *Coxeter transformations and representations of partially ordered sets*. Funkcional. Anal. i Priložen. 8 (1974), no. 3, 34–42. English translation: Functional Anal. Appl. 8 (1974), 219–225 (1975).
- [Drz80] Ju. A. Drozd, *Tame and wild matrix problems*. Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), pp. 242–258, Lecture Notes in Math., 832, Springer, Berlin-New York, 1980.
- [DuVal34] P. Du Val, *On isolated singularities which do not affect the condition of adjunction*. Proc. Cambridge Phil. Soc. 30 (1934), 453–465.
- [Ebl02] W. Ebeling, *Poincaré series and monodromy of a two-dimensional quasihomogeneous hypersurface singularity*. Manuscripta Math. 107 (2002), no. 3, 271–282.
- [EbGu99] W. Ebeling, S. M. Gussein-Zade, *On the index of a vector field at an isolated singularity*. The Arnoldfest (Toronto, ON, 1997), 141–152, Fields Inst. Commun., 24, Amer. Math. Soc., Providence, RI, 1999.
- [Fr51] J. S. Frame, *Characteristic vectors for a product of n reflections*. Duke Math. J. 18, (1951), 783–785.
- [FSS96] J. Fuchs, B. Schellekens, C. Schweigert, *From Dynkin diagram symmetries to fixed point structures*. Comm. Math. Phys. 180 (1996), no. 1, 39–97.
- [Gab72] P. Gabriel, *Unzerlegbare Darstellungen I*. Manuscripta Math. 6 (1972), 71–103.
- [Gb02] A. Gabrielov, *Coxeter-Dynkin diagrams and singularities*. In: Selected Papers of E.B. Dynkin, AMS, 2000, p.367–369.
- [GH01] E. Ghatem, E. Hironaka, *The arithmetic and geometry of Salem numbers*. Bull. Amer. Math. Soc. 38 (2001), 293–314.
- [Ga90] F. Gantmakher, *The theory of matrices*. Chelsea, NY, 1990.
- [GorOnVi94] V. V. Gorbatsevich, A. L. Onishchik, È. B. Vinberg. *Structure of Lie groups and Lie algebras*. (Russian) Current problems in mathematics. Fundamental directions, Vol. 41, 5–259, Itogi Nauki i Tekhniki, Akad. Nauk SSSR, Moscow, 1990. Translation by V. Minachin. Translation edited by A. L. Onishchik and È. B. Vinberg. Encyclopaedia of Mathematical Sciences, 41. Springer-Verlag, Berlin, 1994. iv+248 pp.
- [GP69] I. M. Gelfand and V. A. Ponomarev, *Remarks on the classification of a pair of commuting linear transformations in a finite-dimensional space*. Funkcional. Anal. i Priložen. 3 1969 no. 4, 81–82. English translation: Functional Anal. Appl. 3, 325–326, 1969.
- [GP72] I. M. Gelfand, V. A. Ponomarev, *Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space*. Hilbert space operators and operator algebras (Proc. Internat. Conf., Tihany, 1970), pp. 163–237. Colloq. Math. Soc. Janos Bolyai, 5, North-Holland, Amsterdam, 1972.
- [GP74] I. M. Gelfand, V. A. Ponomarev, *Free modular lattices and their representations*. Collection of articles dedicated to the memory of Ivan Georgievic Petrovskii (1901–1973), IV. Uspehi Mat. Nauk 29 (1974), no. 6(180), 3–58. English translation: Russian Math. Surveys 29 (1974), no.6, 1–56.

- [GP76] I. M. Gelfand, V. A. Ponomarev, *Lattices, representations, and their related algebras. I.* Uspehi Mat. Nauk 31 (1976), no. 5(191), 71–88. English translation: Russian Math. Surveys 31 (1976), no.5, 67–85.
- [Gu76] S. M. Gussein-Zade, *The characteristic polynomial of the classical monodromy for series of singularities.* (Russian) Funkcional. Anal. i Priložen. 10 (1976), no. 3, 78–79. English translation: Functional Anal. Appl. 3, 229–231, 1976.
- [GV83] G. Gonzalez-Sprinberg, J. L. Verdier, *Construction geometrique de la correspondance de McKay.* Ann. Sci. Ecole Norm. Sup. (4) 16, (1983), no.3 409–449.
- [Ha89] M. Hamermesh, *Group Theory and Its Application to Physical Problems.* New York: Dover, 1989.
- [Har95] J. Harris, *Algebraic Geometry. A First Course.* Springer-Verlag, 1995.
- [HPR80] D. Happel, U. Preiser, C. M. Ringel, *Binary polyhedral groups and Euclidean diagrams.* Manuscripta Math. 31 (1980), no. 1-3, 317–329.
- [Hil1890] D. Hilbert, *Über die Theorie der Algebraischen Formen.* Math. Annalen 36. (1890), 473–534.
- [Hir02] E. Hironaka, *Lehmer’s problem, McKay’s correspondence, and 2, 3, 7.* Topics in algebraic and non-commutative geometry (Luminy/Annapolis, MD, 2001), 123–138, Contemp. Math., 324, Amer. Math. Soc., Providence, RI, 2003.
- [Hob02] J. van Hoboken, *Platonic solids, binary polyhedral groups, Kleinian singularities and Lie algebras of type A, D, E.* Master’s Thesis, University of Amsterdam, 2002, <http://home.student.uva.nl/joris.vanhoboken/scriptiejoris.ps>.
- [How82] R. Howlett, *Coxeter groups and M-matrices.* Bull. London Math. Soc. 14 (1982), no. 2, 137–141.
- [Ho41] H. Hopf, *Ueber die Topologie der Gruppenmannigfaltigkeiten und ihre Verallgemeinerungen.* Annals of Math. 42 (1941), 22–52.
- [Hur1893] A. Hurwitz, *Über algebraische Gebilde mit eindeutigen Transformationen in sich.* Math. Ann., 41 (1893), 403–442.
- [Hur1897] A. Hurwitz, *Ueber die Erzeugung der Invarianten durch Integration.* G”ott. Nachrichten, (1897), pp. 71–90.
- [JL2001] G. James, M. Leibeck, *Representations and characters of groups*, 2d edition, Cambridge University Press, 2001.
- [Jen94] G. Jennings, *Modern Geometry with Applications*, Springer-Verlag. New York, 1994.
- [Kac68] V. G. Kac, *Simple irreducible graded Lie algebras of finite growth.* (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 32 (1968) 1323–1367, English translation: Math. USSR-Izv. 2 (1968), 1271–1311.
- [Kac69] V. Kac, *Automorphisms of finite order of semisimple Lie algebras.* (Russian) Funkcional. Anal. i Priložen. 3 1969, 3, 94–96. English translation: Functional Anal. Appl. 3 (1969), 252–254.
- [Kac80] V. Kac, *Infinite root systems, representations of graphs and invariant theory.* Invent. Math. 56 (1980), no. 1, 57–92
- [Kac82] V. Kac, *Infinite root systems, representations of graphs and invariant theory. II.* J. Algebra 78 (1982), no. 1, 141–162.
- [Kac93] V. Kac, *Infinite-Dimensional Lie Algebras.* 3d edition, Cambridge University Press, 1993.
- [Kar92] G. Karpilovsky, *Group Representations: Introduction to Group Representations and Characters*, Vol 1 Part B, North-Holland Mathematics Studies 175, Amsterdam, 1992.
- [Kl1884] F. Klein, *Lectures on the Ikosahedron and the Solution of Equations of the Fifth Degree.* Birkhäuser publishing house Basel, 1993 (reproduction of the book of 1888).
- [Kn85] H. Knörrer, *Group representations and the resolution of rational double points.* in: Finite groups - coming of age. Cont. Math. 45, (1985) 175–222.
- [KMSS83] V. A. Kolmykov, V. V. Menshikh, V. F. Subbotin, M. V. Sumin, *Certain properties of acyclic graphs and polynomials related to them.* Voronezh, VINITI deposition, 12.12.1983, n. 6707-B83, 21 p, (in Russian).
- [KMSS83a] V. A. Kolmykov, V. V. Menshikh, V. F. Subbotin, M. V. Sumin. *Study of spectral properties of trees.* Voronezh, VINITI deposition, 12.12.1983, n. 6709-B83, 30 p, (in Russian).
- [Kos59] B. Kostant, *The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group.* Amer. J. Math. 81 (1959) 973–1032.
- [Kos84] B. Kostant, *The McKay correspondence, the Coxeter element and representation theory.* The mathematical heritage of Elie Cartan (Lyon, 1984), Asterisque 1985, Numero Hors Serie, 209–255.
- [Kosz50] J. L. Koszul, *Homologie et cohomologie des algèbres de Lie*, Bull. Soc. Math. France 78, (1950). 65–127.
- [Kr85] H. Kraft, *Geometrische Methoden in der Invariantentheorie.* (German) [Geometrical methods in invariant theory], Friedr. Vieweg & Sohn, Braunschweig, 1984. x+308 pp.

- [Lev66] J. Levine, *Polynomial invariants of knots of codimension two*. Ann. of Math. (2) 84, 1966, 537–554.
- [Leh33] D. H. Lehmer, *Factorization of certain cyclotomic functions*. Ann. of Math. 34 (1933), 461–469.
- [LuTW00] A. Lucchini, M. C. Tamburini and J. S. Wilson, *Hurwitz groups of large rank*. J. London Math. Soc. (2) 61 (2000), no. 1, 81–92.
- [LuT99] A. Lucchini, M. C. Tamburini, *Classical groups of large rank as Hurwitz groups*. J. Algebra 219 (1999), no. 2, 531–546.
- [Mum88] D. Mumford, *Introduction to algebraic geometry (preliminary version of first 3 chapters)* Reissued as The red book of varieties and schemes, Springer-Verlag, (1988), Lecture Notes in Mathematics, 1358.
- [McM02] C. McMullen, *Coxeter groups, Salem numbers and the Hilbert metric*. Publ. Math. Inst. Hautes Études Sci. no. 95 (2002), 151–183.
- [MeSu82] V. V. Menshikh, V. F. Subbotin, *Orientations of graphs and Coxeter transformations they generate*. Voronezh, VINITI deposition, 30.12.1982, n. 6479-B82, 26p, (in Russian).
- [Men85] V. V. Menshikh, *Conjugacy in the Weyl group and the regularity of the graph representation*. Applications of topology in modern calculus, Voronezh Univ. Press, Voronezh, 1985, 144–150 (in Russian).
- [MM64] M. Markus, H. Minc, *A Survey o Matrix Theory and Matrix Inequalities*. Allyn and Bacon, Boston, 1964.
- [McK80] J. McKay, *Graphs, singularities, and finite groups*. The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), pp. 183–186, Proc. Sympos. Pure Math., 37, Amer. Math. Soc., Providence, R.I., 1980.
- [Mo68] R. V. Moody, *A New Class of Lie Algebras*. J. Algebra 10 1968 211–230.
- [Mohr04] S. Mohr dieck *A Steinberg Cross-Section for Non-Connected Affine Kac-Moody Groups*, 2004, <http://arxiv.org/abs/math.RT/0401203>.
- [Mos98] M. Mossinghoff, *Polynomials with small mahler measure*. Mathematics of Computation, 67 (1998), no. 224, 1697–1705.
- [Mu01] Q. Mushtaq, *Parametrization of triangle groups as subgroups of $\mathrm{PSL}(2, q)$* . Southeast Asian Bull. Math. 25 (2001), no. 2, 309–312.
- [Nag59] M. Nagata. *On the 14th problem of Hilbert*. Amer. J. Math. 81 (1959), 766–772.
- [Naz73] L. A. Nazarova, *Representations of quivers of infinite type*. Izv. Akad. Nauk SSSR Ser. Mat. 37 (1973), 752–791, English translation: Math. USSR-Izv. 7 (1973), 749–792.
- [Newst78] P. E. Newstead. *Introduction to moduli problems and orbit spaces*. Tata Institute Lecture Note, 51, Springer, (1978).
- [OR77] S. A. Ovsienko, A. V. Roiter, *Bilinear forms and categories of representations*. Matrix problems, pp. 71–80. Akad. Nauk Ukrain. SSR Inst. Mat., Kiev, 1977.
- [OvTa77] V. Ovsienko, S. Tabachnikov, *Projective Differential Geometry Old and New From the Schwarzian Derivative to the Cohomology of Diffeomorphism Groups*. Series: Cambridge Tracts in Mathematics, 2005, No. 165.
- [OnVi90] A. L. Onishchik, È. B. Vinberg, *Lie groups and algebraic groups*. Translated from the Russian and with a preface by D. A. Leites. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1990. 328 pp.
- [Pon35] L. Pontrjagin, *Sur les nombres de Betti des groupes de Lie*, C. R. Acad. Sci. Paris 200, 1935, 1277–1280.
- [Pr94] V. V. Prasolov *Problems and theorems in linear algebra*. Translated from the Russian manuscript by D. A. Leites, American Mathematical Society, Providence, RI, 1994. xviii+225 pp.
- [PT90] J. A. de la Peña, M. Takane, *Spectral properties of Coxeter transformations and applications*. Arch. Math. 55 (1990), no. 2, 120–134.
- [PV94] V. L. Popov, È. B. Vinberg, *Invariant Theory*, Encycl. of Math. Sci., Algebraic Geometry. IV, Springer Verlag, Vol. 55, 1994, 123–284.
- [Re88] M. Reid, *Undergraduate Algebraic Geometry*. London Mathematical Society Student Texts 12, Cambridge Univ. Press, Cambridge, 1988
- [Re01] M. Reid, *Surface cyclic quotient singularities and Hirzebruch-Jung resolutions*. 2001, <http://www.maths.warwick.ac.uk/~miles/surf/more/cyclic.pdf>.
- [Rie02] O. Riemschneider, *Special representations and the two - dimensional McKay correspondence*. 2002, preprint, <http://www.math.uni-hamburg.de/home/riemschneider/hokmckay.ps>
- [Rin76] C. M. Ringel, *Representations of K -species and bimodules*. J. Algebra 41 (1976), no. 2, 269–302.

- [Rin94] C. M. Ringel, *The spectral radius of the Coxeter transformations for a generalized Cartan matrix*. Math. Ann. 300 (1994), no. 2, 331–339.
- [Rom00] T. Romstad, *Non-Commutative Algebraic Geometry Applied to Invariant Theory of Finite Group Actions*. Cand. Scient. Thesis. Institute of Mathematics, University of Oslo (2000), <http://folk.uio.no/romstad/thesis.pdf>.
- [Sam41] H. Samelson, *Beiträge zur Topologie der Gruppen-Mannigfaltigkeiten*. (German) Ann. of Math. (2) 42, (1941). 1091–1137.
- [Sat60] I. Satake, *On representations and compactifications of symmetric Riemannian spaces*. Ann. of Math. (2) 71 (1960), 77–110.
- [Sc24] I. Schur, *Neue Anwendung der Integralrechnung auf Probleme der Invariantentheorie*. Sitzungsber. Preuss. Akad. (1924), 189, 297, 346.
- [Shaf88] I. Shafarevich, *Osnovy algebraicheskoi geometrii*. (Russian) [Fundamentals of algebraic geometry.] Second edition. Nauka, Moscow, (1988) 352 pp.
- [ShT54] G. C. Shephard, J. A. Todd, *Finite unitary reflection groups*. Canad. J. Math. 6 (1954), 274–304.
- [Shi00] Jian-yi Shi, *Conjugacy relation on Coxeter elements*. Adv. Math. 161 (2001), no. 1, 1–19.
- [Sl80] P. Slodowy, *Simple singularities and simple algebraic groups*. Lecture Notes in Mathematics, 815. Springer, Berlin, 1980.
- [Sl83] P. Slodowy, *Platonic solids, Kleinian singularities, and Lie groups*. Algebraic geometry (Ann Arbor, Mich., 1981), 102–138, Lecture Notes in Math., 1008, Springer, Berlin, 1983.
- [Sp77] T. A. Springer, *Invariant Theory*, Lect. Notes in Math. 585, Springer-Verlag, NY 1977.
- [Sp87] T. A. Springer, *Poincaré series of binary polyhedral groups and McKay’s correspondence*. Math. Ann. 278 (1987), no. 1–4, 99–116.
- [Sp94] T. A. Springer, *Linear Algebraic Groups*, Encycl. of Math. Sci., Algebraic Geometry. IV, Springer Verlag, Vol. 55, 1994, 1–123.
- [Stn79] R. P. Stanley, *Invariants of finite groups and their applications to combinatorics*, Bull. Amer. Math. Soc., 1 (1979), 475–511.
- [Stb85] R. Steinberg, *Finite subgroups of SU_2 , Dynkin diagrams and affine Coxeter elements*. Pacific J. Math. 118 (1985), no. 2, 587–598.
- [Stb59] R. Steinberg, *Finite reflection groups*. Trans. Amer. Math. Soc. 91 (1959), 493–504.
- [St75] R. B. Stekolshchik, *Conditions for the regularity of representations of graphs*. In: Transactions of department of mathematics of Voronezh University. 16, Voronezh Univ. Press, Voronezh (1975), 58–61, (in Russian).
- [St81] R. B. Stekolshchik, *Coxeter numbers of the extended Dynkin diagrams with multiple edges*. Abstracts of the XVI National (the USSR) algebraic conference. Part 2, Leningrad, p.187, 1981, (in Russian).
- [St82] R. B. Stekolshchik, *Transforming elements connecting the Coxeter transforms, and conditions for the regularity of roots*. Funkt. Anal. i Prilozhen. 16 (1982), no. 3, 84–85. English translation: Functional Anal. Appl. 16 (1982), no. 3, 229–230 (1983).
- [St82a] R. B. Stekolshchik, *The Coxeter transformation associated with the extended Dynkin diagrams with multiple edges*. Kishinev, VINITI deposition, 02.09.1982, n. 5387-B82, 17p, (in Russian).
- [St85] R. B. Stekolshchik, *Study of Coxeter transformations and regularity problems associated with representation theory of graphs*. Ph.D.Thesis, Kiev, 95p, 1985, (in Russian).
- [SuSt75] V. F. Subbotin, R. B. Stekolshchik, *The spectrum of the Coxeter transformation and the regularity of representations of graphs*. In: Transactions of department of mathematics of Voronezh University. 16, Voronezh Univ. Press, Voronezh (1975), 62–65, (in Russian).
- [SuSt78] V. F. Subbotin, R. B. Stekolshchik, *The Jordan form of the Coxeter transformation, and applications to representations of finite graphs*. Funkcional. Anal. i Priložen. 12 (1978), no. 1, 84–85. English translation: Functional Anal. Appl. 12 (1978), no. 1, 67–68.
- [SuSt79] V. F. Subbotin, R. B. Stekolshchik, *Sufficient conditions of regularity of representations of graphs*. Applied analysis, Voronezh, (1979), 105–113, (in Russian).
- [SuSt82] V. F. Subbotin, R. B. Stekolshchik, *The relation between the Coxeter transformation, the Cartan matrix and Chebyshev polynomials*. Operator methods in nonlinear analysis, 118–121, Voronezh. Gos. Univ., Voronezh, 1982, (in Russian).
- [SuSum82] V. F. Subbotin, M. V. Sumin, *Study of polynomials related with graph representations*. Voronezh, VINITI deposition, 30.12.1982, n. 6480-B82, 37p, (in Russian).
- [SuU85] V. F. Subbotin, N. N. Udodenko, *On certain properties of characteristic polynomials, of the Coxeter transformation*. Voronezh, VINITI deposition, 07.08.1985, n. 5922-B85, 34p, (in Russian).

- [SuU88] V. F. Subbotin, N. N. Udodenko, *On certain results and problems of the representation theory of graphs and the spectral theory of the Coxeter transformation*. Voronezh, VINITI deposition, 12.07.1988, n. 6295-B88, 60p, (in Russian).
- [Ti66] J. Tits, *Classification of algebraic semisimple groups*. 1966 Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965) pp. 33–62 Amer. Math. Soc., Providence, R.I., 1966.
- [Vin85] È. B. Vinberg, *Hyperbolic reflection groups*. (Russian) Uspekhi Mat. Nauk 40 (1985), no. 1(241), 29–66, 255 English translation: Russian Math. Surveys 40 (1985), no.1, 31–75.
- [Weib] C. A. Weibel, *History of Homological Algebra*.
<http://www.math.rutgers.edu/~weibel/history.dvi>
- [Wi01] R. A. Wilson, *The Monster is a Hurwitz group*. J. Group Theory 4 (2001), no. 4, 367–374.
- [Zh89] Y. Zhang, *Eigenvalues of Coxeter transformations and the structure of regular components of an Auslander-Reiten quiver*. Comm. Algebra 17 (1989), no. 10, 2347–2362.
- [Zhe73] D. Zhelobenko, *Compact Lie groups and their representations*. AMS. Translations of Math. Monographs, 40 (1973).